

Time: 2:30 Hours

Total Marks: 75

N.B: 1) All questions are compulsory

2) From questions 1, 2 and 3 attempt any **One** from part (a) and any **Two** from part (b)

3) Attempt any **Three** from question 4

4) Figures to the right indicate marks

1. (a) i) Let $f: [a, b] \rightarrow \mathbb{R}$ be a bounded function. If P, Q are partitions of $[a, b]$, then Prove that 8
 $1) L(P, f) \leq U(P, f) \quad 2) L(P, f) \leq U(Q, f)$
- ii) Let $f, g: [a, b] \rightarrow \mathbb{R}$ be Riemann integrable on $[a, b]$. Prove that $f + g$ is Riemann 8
integrable on $[a, b]$ and hence show that $\int_a^b (f + g) = \int_a^b f + \int_a^b g$
- (b) i) Let $f: [a, b] \rightarrow \mathbb{R}$ be a continuous function on $[a, b]$. Show that f is Riemann integrable 6
on $[a, b]$.
- ii) Let $f(x) = \begin{cases} 1 - x & \text{if } 0 \leq x < \frac{1}{2} \\ \frac{x}{2} & \text{if } \frac{1}{2} \leq x \leq 1 \end{cases}$ 6
 $F: [0, 1] \rightarrow \mathbb{R}$ is defined by $F(x) = \int_0^x f(t) dt, x \in [0, 1]$. Show that F is differentiable
at $\frac{1}{2}$ and $F'(1/2) = f(1/2)$.
- iii) Let $f: [0, 1] \rightarrow \mathbb{R}$ be defined by $f(x) = x^2$. Using Riemann criterion show that f is 6
Riemann integrable on $[0, 1]$.
- iv) Express the sum $\sum_{r=0}^{n-1} \frac{1}{\sqrt{n^2 - r^2}}$ as a Riemann sum of a suitable function and evaluate 6
 $\lim_{n \rightarrow \infty} \sum_{r=0}^{n-1} \frac{1}{\sqrt{n^2 - r^2}}$
2. (a) i) State and Prove Fubini's theorem for a rectangular domain in \mathbb{R}^2 . 8
- ii) Define double integral of a bounded function $f: Q \rightarrow \mathbb{R}$ where $Q = [a, b] \times [c, d]$ is a 8
rectangle in \mathbb{R}^2 . Further show with usual notations
 $m(b - a)(d - c) \leq \iint_Q f \leq M(b - a)(d - c)$
- (b) i) State the change of variable formula for triple integrals. Stating clearly the conditions 6
under which it is valid. Express further, how will you use it to express the triple integral
in Spherical coordinates.
- ii) Evaluate $\int_0^1 \int_{y^2}^y x dx dy$ by reversing the order of integration. Sketch the region of 6
integration.
- iii) Introduce suitable change of variables and show that 6
 $\iint_S f(xy) dx dy = \ln 2 \int_1^2 f(u) du$, where S is the region in the 1st quadrant bounded by
the curves $xy = 1, xy = 2, y = x, y = 4x$.
- iv) Find the volume of the solid S by using triple integration where S is bounded by the 6
paraboloid $z = x^2 + y^2$ and the plane $z = 2$

- 3 (a) i) Let $\{f_n\}$ be a sequence of real valued R -integrable functions on $[a, b]$. If $\{f_n\}$ converges uniformly to f on $[a, b]$ then show that f is R -integrable on $[a, b]$ and $\int_a^b \lim_{n \rightarrow \infty} f_n(x) dx = \lim_{n \rightarrow \infty} \int_a^b f_n(x) dx$ 8
- ii) Let $\{f_n\}$ be a sequence of continuously differentiable real valued functions defined on $[a, b]$. If the series $\sum_{n=1}^{\infty} f_n$ converges pointwise to f on $[a, b]$ and the series $\sum_{n=1}^{\infty} f_n'$ converges uniformly on $[a, b]$, then show that $f'(x) = \sum_{n=1}^{\infty} f_n'(x)$ for $a \leq x \leq b$ 8
- (b) (i) Let $\{f_n\}$ be a sequence of real values functions defined on a non-empty subset S of \mathbb{R} . Show that $\{f_n\}$ converges uniformly to a function f if and only if for given $\epsilon > 0$, \exists a positive integer n_0 such that $|f_n(x) - f_m(x)| < \epsilon$ for $m, n \geq n_0$ and each $x \in S$ 6
- (ii) Discuss the pointwise and uniform convergence of the series of functions $\sum_{n=1}^{\infty} \frac{1}{(nx)^2}$, $x \neq 0$ 6
- (iii) By integrating a suitable power series term by term show that $\tan^{-1} x = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{2n+1}$ for $|x| < 1$ 6
- (iv) Let $f_n: [0, 1] \rightarrow \mathbb{R}$ be defined by $f_n(x) = \begin{cases} n(1-nx) & \text{for } 0 < x < \frac{1}{n} \\ 0 & \text{otherwise} \end{cases}$ 6
Check whether $\int_0^1 \lim_{n \rightarrow \infty} f_n(x) dx = \lim_{n \rightarrow \infty} \int_0^1 f_n(x) dx$. Does $\{f_n\} \rightarrow f$ uniformly? Justify.
- 4 i) Prove that if $f: [a, b] \rightarrow \mathbb{R}$ is Riemann integrable then $|f|$ is Riemann integrable on $[a, b]$. Is converse true? Justify 5
- ii) If $f, g: [a, b] \rightarrow \mathbb{R}$ are Riemann integrable and have antiderivatives F and G on $[a, b]$ then show that $\int_a^b F(x)g(x) = [f(b)G(b) - F(a)G(a)] - \int_a^b f(x)G(x)$ 5
- iii) Evaluate the following integral by using polar coordinates 5
 $\int_0^{\sqrt{2}} \int_y^{\sqrt{4-y^2}} \frac{dx dy}{1+x^2+y^2}$
- iv) Use spherical coordinates evaluate $\iiint_S x e^{(x^2+y^2+z^2)^2} dx dy dz$ where S is the solid that lies between the spheres $x^2 + y^2 + z^2 = 1$ and $x^2 + y^2 + z^2 = 4$ 5
- v) If a real power series $\sum_{n=0}^{\infty} a_n x^n$ has radius of convergence r , then show that it converges uniformly on $[-s, s]$ where $0 \leq s < r$. 5
- vi) Let $f_n(x) = \frac{x^n}{1+x^n}$ for $0 \leq x \leq 1$. Discuss the pointwise and uniform convergence of $\{f_n\}$ on $[0, 1]$. 5
