

N.B. 1. All questions are compulsory.

2. From Question 1, 2 and 3, Attempt any one from part(a) and any two from part(b).

3. From Question 4, Attempt any THREE

4. Figures to the right indicate marks for the respective parts.

- Q.1 a i Let $\langle f_n \rangle$ be sequence of differentiable real valued functions on $[a, b]$ (8)
such that $\langle f_n(x_0) \rangle$ converges for some $x_0 \in (a, b)$ and $\langle f'_n \rangle$
converges uniformly to function g on $[a, b]$. Prove that $\langle f_n \rangle$
converges uniformly on $[a, b]$ and if f is uniform limit of $\langle f_n \rangle$
then f is differentiable on (a, b) and $f' = g$ on (a, b) .
- ii State and prove Weierstrass M- test.
- b i State and prove Cauchy's criterion for uniform convergence of the (12)
sequence $\langle f_n \rangle$ of functions of real numbers.
- ii Examine whether $\int_0^1 \sum_{n=0}^{\infty} x^n (1 - 2x^n) dx = \sum_{n=0}^{\infty} \int_0^1 x^n (1 - 2x^n) dx$. Is the
series $\sum_{n=0}^{\infty} x^n (1 - 2x^n)$ uniformly convergent in $[0, 1]$? Justify.
- iii Find M_n , where $M_n = \sup \left\{ \frac{x}{(n+x^2)^2} : x \in [a, b] \right\}$, using Weierstrass
M- test. Evaluate $\int_a^b \sum_{n=0}^{\infty} \frac{x}{(n+x^2)^2} dx$.
- iv Let $f_n: [0, 1] \rightarrow \mathbb{R}$ be given by $f_n(x) = x^n$. Let f be pointwise limit of
 $\langle f_n \rangle$. Is f continuous on $[0, 1]$. Does $\langle f_n \rangle$ converge uniformly
on $[0, 1]$? Justify.
- Q.2 a i If a function f is continuous throughout a region R that is closed and (8)
bounded then show that there exists a non-negative integer M such that
 $|f(z)| \leq M \forall z \in R$. Also show that if $f'(z_0), g'(z_0)$ exist,
 $g'(z_0) \neq 0$, $f(z_0) = 0 = g(z_0)$ then $\lim_{z \rightarrow z_0} \frac{f(z)}{g(z)} = \frac{f'(z_0)}{g'(z_0)}$.
- ii Let $f(z) = u(x, y) + iv(x, y)$. If $f'(z)$ exists at a point $z_0 = x_0 + iy_0$
then prove that the first order partial derivatives of u and v exist at
 (x_0, y_0) and satisfy Cauchy- Riemann equations $u_x = v_y$, $u_y = -v_x$.
Show that the converse is not true. Also show that $f'(z) = (u_x)_{z=z_0} +$
 $i(v_x)_{z=z_0}$.
- b i Using the definition, discuss differentiability of the function $f(z) = z^2$ (12)
at any $z \in \mathbb{C}$.
- ii f is analytic throughout on a given domain D . If $|f(z)|$ is constant on
 D , show that $f(z)$ must be constant on D .
- iii If a function $f(z) = u(x, y) + iv(x, y)$ is analytic in a domain D then
show that its component functions u and v are harmonic in D .
- iv Find the image of the given set under the reciprocal map $w = \frac{1}{z}$ in the
extended complex plane: $\frac{1}{5} \leq |z| \leq 2$

(P.T.O)

- Q.3 a i State and prove Cauchy Integral Theorem. (8)
- ii Suppose that a function f is analytic throughout a disk $|z - z_0| < R_0$ centered at z_0 and with radius R_0 . Then prove that $f(z)$ has the power series representation $f(z) = \sum_{n=0}^{\infty} a_n(z - z_0)^n$, $|z - z_0| < R_0$ where $a_n = \frac{f^{(n)}(z_0)}{n!}$.

- b i If a function f is analytic at a given point then show that its derivatives of all orders are analytic at that point too. Further suppose that a function f is analytic inside and on a positively oriented circle C_R centered at z_0 and with radius R and if M_R denotes the maximum value of $|f(z)|$ on C_R then show that $|f^{(n)}(z_0)| \leq \frac{n! M_R}{R^n}$, $n = 1, 2, 3, \dots$ (12)
- ii Prove that a power series $\sum_{n=0}^{\infty} a_n(z - z_0)^n$ represents a continuous function $S(z)$ at each point inside its circle of convergence $|z - z_0| = r$.
- iii Show that any singular point of the function $f(z) = \frac{e^z}{z^2 + \pi^2}$ is a pole. Further determine the order m of each pole and find the corresponding residue of f .
- iv State Laurent's Theorem. For $f(z) = \frac{-1}{(z-1)(z-2)}$, write Laurent series expansion in the domains: $|z| < 1$, $2 < |z| < \infty$.

- Q.4 i Does the sequence $\langle f_n \rangle$, where $f_n(x) = \frac{nx}{1+nx^2}$ converges uniformly on $[0, \infty)$? Justify. (15)
- ii For $|x| < 1$, show that $\tan^{-1}x = \sum_{n=1}^{\infty} \frac{(-1)^{n-1} x^{2n-1}}{2n-1}$.
- iii Test differentiability of the function $f(z) = z \operatorname{Im} z$ at $(0, 0)$.
- iv Construct a linear fractional transformation that maps the points $i, \infty, 3$ to $\frac{1}{2}, -1, 3$ respectively.
- v Evaluate $\int_C \frac{1}{(z-z_0)^{n+1}} dz$ where C is the circle $|z - z_0| = r$, n is a non zero integer using a parameterisation of C .
- vi Evaluate $\int_C \frac{\sin^6 z}{(z - \frac{\pi}{2})^3} dz$ where $C : |z| = 2$.