

(3 Hours)

[Total Marks: 80]

N.B.: 1) Question No. 1 is Compulsory.

2) Answer any THREE questions from Q.2 to Q.6.

3) Figures to the right indicate full marks.

1) a) If $A = \begin{bmatrix} 2 & 4 \\ 0 & 3 \end{bmatrix}$ then find the eigen values of $6A^{-1} + A^3 + 2I$ [05]

b) Determine whether the given vectors $u = (-4, 6, -10, 1), v = (2, 1, -2, 9)$ are orthogonal with respect to the Euclidean inner product [05]

c) The probability density function of a random variable x is zero except at $x = 0, 1, 2$ and

$p(0) = 3\alpha^3, p(1) = 4\alpha - 10\alpha^2, p(2) = 5\alpha - 1$. Find α [05]

d) Evaluate $\oint_c \frac{z+6}{z^2-4} dz$ where c is (i) $|z| = 1$ (ii) $|z-2| = 1$. [05]

2) a) Using Rayleigh-Ritz method, find an appropriate solution for the extremal of the functional

$I = \int_0^1 [2xy - y^2 - y'^2] dx$ given $y(0)=y(1)=0$ [06]

b) Using Cauchy's Residue theorem evaluate $\int_0^{2\pi} \frac{d\theta}{5 + 4 \cos \theta}$ [06]

c) A random variable X has the probability distribution given below:

$X=x$	-2	3	1
$P(X=x)$	1/3	1/2	1/6

Find i) the moment generating function ii) the first four moments about the origin [08]

3) a) Compute $A^9 - 6A^8 + 10A^7 - 3A^6 + A + I$ where $A = \begin{bmatrix} 1 & 2 & 3 \\ -1 & 3 & 1 \\ 1 & 0 & 2 \end{bmatrix}$ [06]

b) Verify Cauchy-Schwartz inequality for the vectors $u = (-4, 2, 1)$ & $v = (8, -4, -2)$ [06]

c) Obtain Taylor's or Laurent's series expansion of the function $f(z) = \frac{1}{z^2 - 3z + 2}$ when

(i) $|z| < 1$ (ii) $1 < |z| < 2$ [08]

- 4) a) Obtain the equation of the line of regression of Y on X for the following data and estimate Y when X = 73

[06]

X	70	72	74	76	78	80
y	163	170	179	188	196	200

- b) Show that the functional $\int_{x_1}^{x_2} [y^2 + x^2 y'] dx$ assumes extreme values on the straight line $y = x$

[06]

- c) Let R^3 have the Euclidean inner product. Use the Gram-Schmidt process to transform

the basis vectors $u_1 = (1, 0, 0), u_2 = (3, 7, -2), u_3 = (0, 4, 1)$ into an orthonormal basis

[08]

- 5) a) Evaluate $\oint_c \frac{1}{z} \cos z \, dz$ where c is the ellipse $9x^2 + 4y^2 = 1$

[06]

- b) Seven dice are thrown 729 times. How many times do you expect at least four 10 dice to show three or five?

[06]

- c) Show that the matrix $A = \begin{bmatrix} -9 & 4 & 4 \\ -8 & 3 & 4 \\ -16 & 8 & 7 \end{bmatrix}$ is diagonalisable. Find the diagonal form D and the

diagonalising matrix M.

[08]

- 6) a) A continuous random variable X has the p.d.f. defined by $f(x) = A + Bx, 0 \leq x \leq 1$. If the mean of the distribution is $\frac{1}{3}$ find A and B

[06]

- b) Find e^A , if $A = \begin{bmatrix} 3 & 1 \\ 2 & 2 \\ 1 & 3 \\ 2 & 2 \end{bmatrix}$

[06]

- c) Evaluate $\int_{-\infty}^{\infty} \frac{x^2 dx}{(x^2 + a^2)(x^2 + b^2)}$ ($a > 0, b > 0$)

[08]
