Time Dura	tion: 3Hr	Total Marks: 80
	N.B.:1) Question no.1 is compulsory.2) Attempt any three questions from Q.2to Q.6.3) Figures to the right indicate full marks.	
Q1. a) b) c)	Find the Laplace transform of $e^{-4t}t \sin 3t$. Find the half-range cosine series for $f(x) = x$, $0 < x < 2$.	[5] [5] [5]
d)	Find $\nabla \cdot \left(r \nabla \frac{1}{r^3}\right)$. Show that the function $f(z) = \sin z$ is analytic and find $f'(z)$ in terms of z.	[5]
Q2. a)	Find the inverse Z-transform of $F(z) = \frac{1}{(z-5)^3}$, $ z < 5$.	[6]
b) c)	Find the analytic function whose imaginary part is $e^{-x}(y \sin y + x \cos y)$. Obtain Fourier series for the function $f(x) = x + x^2$, $-\pi \le x \le \pi$ a $f(x + 2\pi) = f(x)$. Hence deduce that $\frac{\pi^2}{6} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \cdots$ and $\frac{\pi^2}{8} = \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \cdots$	[6] nd [8]
Q3. a)	Find $L^{-1}\left[\frac{1}{(s-a)(s-b)}\right]$ using convolution theorem.	[6]
	Is $S = \left\{ \sin\left(\frac{\pi x}{4}\right), \sin\left(\frac{3\pi x}{4}\right), \sin\left(\frac{5\pi x}{4}\right), \dots \dots \right\}$ orthogonal in $(0, 1)$?	[6]
c)	Using Green's theorem in the plane evaluate $\int_c (xy + y^2)dx + (x^2)dy$ where is the closed curve of the region bounded by $y = x$ and $y = x^2$.	C [8]
Q4. a)	Find Laplace transform of $f(t) = \begin{cases} \sin 2t & \text{if } 0 < t \le \frac{\pi}{2} \\ 0 & \text{if } \frac{\pi}{2} < t < \pi \end{cases}$ and	[6]
b)	$f(t) = f(t + \pi)$. Prove that a vector field \overline{f} is irrotational and hence find its scalar potential	[6]
c)	$\bar{f}=(x^2+xy^2)i+(y^2+x^2y)j$. Find the Fourier expansion for $f(x)=\sqrt{1-\cos x}$ in $(0,2\pi)$. Hence deduthat $\frac{1}{2}=\sum_{1}^{\infty}\frac{1}{4n^2-1}$.	[8]
Q5.a)	Use Gauss's Divergence Theorem to show that $\iint_S \nabla r^2 ds = 6V$ where S is a	ny [6]
b) c)	closed surface enclosing a volume V. Find the Z-transform of $f(k) = b^k$, k< 0. i) Find $L^{-1}\left[\frac{s}{(s-2)^6}\right]$. ii) Find $L^{-1}\left[\log\left(1+\frac{a^2}{s^2}\right)\right]$.	[6] [8]
Q6.a)	Solve using Laplace transform $(D^2 + 9)y = 18t$, given that $y(0) = 0$ and $y(\frac{\pi}{2}) = 0$	[6]
b)	Find the bilinear transformation which maps the points $Z=\infty$, i, 0 onto $W=0$, i, ∞ .	[6]
c)	Find Fourier integral representation of $f(x) = e^{- x } - \infty < x < \infty$.	[8]

67960