

[Time: 2 ½ Hours ]

[ Total Marks : 60 ]

**N.B.:(1) All questions are compulsory.****(2) Figures to the right indicate full marks.****(3) Symbols have their usual meanings unless otherwise stated.****(4) Use of log tables / non-programmable calculator is allowed.**1. (a) Attempt any **one** : -

8

(i) a) Derive Cauchy Riemann equations in cartesian form.

b) Determine whether  $\frac{1}{z}$  is analytic or not?(ii) If  $f(z)$  is analytic in a closed curve  $C$ , except at a finite number of poles within  $C$ , then

$$\int_C f(z) dz = 2\pi i (\text{sum of residues})$$

Hence Evaluate  $\int_C \frac{4-3z}{z(z-1)(z-2)} dz$  where  $C$  is the circle  $|z| = \frac{3}{2}$ (b) Attempt any **one** : -


4

(i) Prove that  $u = x^2 - y^2 - 2xy - 2x + 3y$  is harmonic. Find a function  $v$  such that  $f(z) = u + iv$  is analytic.(ii) Expand  $f(z) = \frac{1}{(z+1)(z+3)}$  in Laurent series fora)  $1 < |z| < 3$ b)  $|z| > 3$ 2. (a) Attempt any **one** : -

8

(i) Find the inertia tensor about the origin for the mass distribution consisting of mass 1 at  $(0, 1, 1)$  and mass 2 at  $(1, -1, 0)$ . Find the principal moment of inertia and the principal axis.(ii) Find the characteristic frequencies and characteristic mode of vibration for the system of masses ( $m$  &  $4m$ ) and springs having spring constants ( $3k$  &  $k$ ) as shown in the figure the motion is along a vertical line.

(Neglect mass of the spring.)b



(b) Attempt any **one** : -

4

(i) Using the Levi-Civita symbol, show that,  $A \times (B \times C) = B(A \cdot C) - C(A \cdot B)$ (ii) Find the eigenvalues of the matrix  $\begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & -2 \\ 2 & -2 & 1 \end{bmatrix}$ 3. (a) Attempt any **one** : -

8

(i) Solve the Laguerre equation using Frobenius method

$$x \frac{d^2 y}{dx^2} + (1 - x) \frac{dy}{dx} + ny = 0$$

(ii) Solve the Bessel equation

$$x^2 \frac{d^2 y}{dx^2} + x \frac{dy}{dx} + (x^2 - n^2) y = 0 \text{ using Frobenius method.}$$

(b) Attempt any **one** : -

4

(i) Using the generating function for Hermite polynomials

$$G(x, t) = e^{2tx - t^2} = \sum_{n=0}^{\infty} H_n(x) \frac{t^n}{n!} \text{ prove that,}$$

$$2xH_n(x) = 2n H_{n-1}(x) + H_{n+1}(x)$$

(ii) Using Green's Theorem, Find the area of the region in the first quadrant bounded by the curves  $y = x$ ,  $y = \frac{1}{x}$ ,  $y = \frac{x}{4}$ .4. (a) Attempt any **one** : -

8

(i) State the expression of Fourier transform and its inverse in three-dimensional space. Find the Fourier transform of Yukawa potential  $\frac{e^{-ar}}{r}$ 

(ii) Solve the differential equation using Laplace transform

$$y'' - y = t \quad \text{subject to initial conditions} \quad y(0) = 1 \quad \& \quad y'(0) = 1$$

(b) Attempt any **one** : -

4

- (i) Find the Laplace transform of  $f(t) = t \sinh at$ .
- (ii) State and prove Fourier Convolution Theorem.

5. Attempt any **four** : -

12

- (a) Determine the poles of the function and residue at the poles:  $f(z) = \frac{z}{\sin z}$
- (b) Use Cauchy's integral formula to calculate  

$$\oint_C \frac{2z + 1}{z^2 + z} dz \quad \text{where } C \text{ is } |z| = \frac{1}{2}$$
- (c) What is the rank of the tensor  $T_{ijkl}$  ? Write its transformation equation.
- (d) For matrices M, C and D, show that  $M^n = C D^n C^{-1}$  where  $C^{-1} M C = D$  and D is diagonal.
- (e) Prove that  $P_n(1) = 1$
- (f) Find regular singular point of the differential equation  

$$2x^2 \frac{d^2 y}{dx^2} + 3x \frac{dy}{dx} + (x^2 - 4)y = 0$$
- (g) Find the inverse Laplace transform of  $F(s) = \frac{7}{(s-1)^3}$
- (h) Show Fourier transform property:  $[\nabla f(\vec{r})]^T(\vec{k}) = -i\vec{k}g(\vec{k})$

\*\*\*\*\*