UNIT I

1

THE MEAN, MEDIAN, MODE, AND OTHER MEASURES OF CENTRAL TENDENCY

Unit structure

- 1.0 Objectives
- 1.1 Introduction
- 1.2 Index or Subscript, Notation, Summation Notation
- 1.3 Averages or Measures of Central Tendency
- 1.4 Arithmetic Mean
 - 1.4.1 Arithmetic Mean Computed from Grouped Data
 - 1.4.2 Properties of the Arithmetic Mean
- 1.5 Weighted Arithmetic Mean
- 1.6 Median
- 1.7 Mode
- 1.8 Empirical Relation between the Mean, Median, and Mode
- 1.9 Geometric Mean
- 1.10 Harmonic Mean
- 1.11 Relation between the Arithmetic, Geometric, and Harmonic Means
- 1.12 Root Mean Square
- 1.13 Quartiles, Deciles and Percentiles
- 1.14 Software and Measures of Central Tendency
- 1.15 Summary
- 1.16 Exercise
- 1.17 References

1.0 OBJECTIVES

After going through this chapter, students will able to learn

- To present huge data in a summarized form
- To calculate and interpret the mean, the median and the mode,
- To facilitate comparison
- To calculate geometric mean, harmonic mean
- To trace precise relationship

- To calculate Quartiles, deciles and percentiles
- To help in decision-making

1.1 INTRODUCTION

A measure of central tendency is a single value that describes a set of data by identifying the central position within that set of data. Mean, median and mode are different measures of central tendency in a numerical data. The word average is commonly used in day to day conversation like we often talk about average height of the girls, average student of the class, and average run rate of the match. When we say average means neither too good nor bad. However, in statistics the term average has different meaning. Average is a single value which representing a group of values so such a value easy to understand, easy to compute and based on all observations.

1.2 INDEX OR SUBSCRIPT, NOTATION, SUMMATION NOTATION

Let the symbol X_i (read 'X subscript i) denote any of the N values $X_1, X_2, X_3, \ldots, X_N$ assumed by a variable X. The letter i in X_i which can stand for any of the numbers 1, 2, 3,, N is called a subscript or index. Any letter other than i such as j, k, p, q or r could be used also.

Summation Notation:

The symbol $\sum_{i=1}^{N} X_i$ is used to denote the sum of all the X_i 's from i = 1 to N.

 $\sum_{i=1}^{N} X_i = X_1 + X_2 + X_3 + \dots + X_N$

We generally denote this sum simply by $\sum X$, $\sum X_i$.

The symbol \sum is the Greek capital letter *sigma* denoting sum.

Ex.
$$\sum_{i=1}^{N} aX_i = aX_1 + aX_2 + aX_3 + \dots + aX_N$$

= a $(X_1 + X_2 + X_3 + \dots + X_N)$ = a $\sum_{i=1}^N X_i$, where a

is a constant.

1.3 AVERAGES OR MEASURES OF CENTRAL TENDENCY

There are different ways of measuring the central tendency of a set of values.

Various authors defined Average differently.

"Average is an attempt to find one single figure to describe whole of figures." – Clark

"An average is a single value selected from a group of values to represent them in some way- a value which is supposed to stand for whole group, of which it is a part, as typical of all the values in the group." – A. E. Waugh

"An average is a typical value in the sense that it is sometimes employed to represent all the individual values in the series or of a variable." – **Ya-Lun-Chou**

Types of Averages:

Arithmetic Mean: a. Simple, b. Weighted Median Mode Geometric Mean Harmonic Mean

1.4 ARITHMETIC MEAN

The most popular and widely used measure of representing the entire data by one value is mean or Average.

It simply involves taking the sum of a group of numbers, then dividing that sum by the total number of values in the group.

Arithmetic mean can be of two types.

- a. Simple arithmetic mean
- b. Weighted arithmetic mean

A. Simple Arithmetic Mean – Individual Observations:

Calculation of mean in case of individual observations [i. e. when frequencies are not given] is very simple. Here, we add all values of the variable and divide the total by the number of items.

$$\overline{X} = \frac{X_1 + X_2 + X_3 + \dots + X_n}{N} = \frac{\sum X}{N}$$

 \overline{X} = Arithmetic Mean; N = number of observations;

 $\sum X =$ sum of all the values of the variable X i. e. $X_1, X_2, X_3, \dots, X_n$

Ex 1. Find the Arithmetic mean of following five values 8, 45, 49, 54, 79. Sol: We know that, $\overline{X} = \frac{X_1 + X_2 + X_3 + \dots + X_n}{N}$ $\overline{X} = \frac{8+45+49+54+79}{5} = \frac{235}{5} = 47$

Ex 2. Find the Arithmetic mean of following values. 4350, 7200, 6750, 5480, 7940, 3820, 5920, 8450, 4900, 5350.

Sol: We know that,
$$\overline{X} = \frac{X_1 + X_2 + X_3 + \dots + X_n}{N}$$

 $\bar{X} = \frac{8350 + 7200 + 6750 + 5480 + 7940 + 3820 + 5920 + 8450 + 4900 + 5350}{10} = \frac{64160}{10} = 6416$

Short cut method: $\overline{X} = A + \frac{\sum d}{N}$

Where A is assumed mean and d is deviation of items from assumed mean i. e. d = (X - A).

Ex 3. Calculate arithmetic mean from following data. 2690, 3670, 4580, 5660, 2750, 2830, 4100, 5720, 5040, 4840 Sol:

X	d = (X-A)
2690	-2310
3670	-1330
4580	-420
5660	660
2750	-2250
2830	-2170
4100	-900
5720	720
5040	40
4840	-160
	$\sum d = -8120$

Consider assumed mean, A = 5000 $\bar{X} = A + \frac{\sum d}{N} = 5000 - \frac{8120}{10} = 4188$

1.4.1 Arithmetic Mean Computed from Grouped Data:

Simple Arithmetic Mean – Discrete series:

Calculation of mean in case of frequencies are given,

$$\bar{X} = \frac{\sum f X}{N}$$

f = Frequency;

X = the variable

N = Total number of observations i.e. $\sum f$

Here, first multiply the frequency of each row with variable and obtain the total $\sum f X$ and then divide the total by number of observations, i.e. total frequency.

Ex 4. Following are the marks obtained by 60 students. Calculate arithmetic mean.

Marks	15	30	45	60	70	80
No. of students	6	14	15	15	4	6

Sol: Let the marks denoted by X and number of students denoted by f.

Marks	No. of Students	fX
X	f	
15	6	90
30	14	420
45	15	675
60	15	900
70	4	280
80	6	480
	N = 60	$\sum f X = 2845$

$$\bar{X} = \frac{\sum f X}{N} = \frac{2845}{60} = 47.42$$

Short cut method: $\bar{X} = A + \frac{\sum fd}{N}$ Where A is assumed mean and d is deviation of items from assumed mean

i. e. d = (X - A), N = $\sum f$

Ex 5. Calculate arithmetic mean by the short cut method using data from Ex. 4

0 1	
SA	•
501	٠

Marks X	No. of Students f	$\mathbf{d} = (\mathbf{X} - \mathbf{A})$	fd
15	6	-30	-180
30	14	-15	-210
45	15	0	0
60	15	15	225
70	4	25	100
80	6	35	210
	N = 60		$\sum f X = 145$

Assumed mean, A = 45 $\bar{X} = A + \frac{\sum fd}{N} = 45 + \frac{145}{60} = 47.4166$

Simple Arithmetic Mean – Continuous Series:

$$\bar{X} = \frac{\sum fm}{N}$$

m = mid-point of various classes; f = the frequency of each class;

N = the total frequency

Here, first obtain the mod-point of each class and denote it by m.

Multiply these mid-points by the respective frequency of each class and obtain the total $\sum fm$

Divide the total by the sum of the frequency, i.e. N.

Ex 5. From the following data compute arithmetic mean.

Marks	0-10	10-20	20-30	30-40	40-50	50-60
No. of students	5	10	25	30	20	10

Sol:

Marks	Mid- point m	No. of Students f	fm
0-10	5	5	25
10-20	15	10	150
20-30	25	25	625
30-40	35	30	1050
40-50	45	20	900
50-60	55	10	550
		N = 100	$\sum f m = 3300$
$\overline{I} = \frac{\sum f m}{N} =$	$=\frac{3300}{100}=33$		

 $\bar{X} = \frac{\sum f m}{N} = \frac{3300}{100} = 33$

Ex 6. From the following data compute arithmetic mean.

Class Intervals	0-10	10-20	20-30	30-40	40-50	50-60	60-70
Frequency	4	4	7	10	12	8	5

Sol:

Marks	Mid- point m	No. of Students f	f m
0-10	5	4	20
10-20	15	4	60
20-30	25	7	175
30-40	35	10	350
40-50	45	12	540
50-60	55	8	440
50-60	65	5	325
		N = 50	$\sum f m = 1910$

$$\bar{X} = \frac{\sum f m}{N} = \frac{1910}{50} = 38.2$$

Short cut method: $\overline{X} = A + \frac{\sum fd}{N}$

Where A is assumed mean and d is deviation of items from assumed mean i. e. d = (m - A), m= mid point , N = $\sum f$

Ex 7. Calculate arithmetic mean by the short cut method using data from Ex. 5.

S	ol:					
	Marks	Mid-	point	No. of Students f	d = (m - A)	fd
		m				
	0-10	5		5	-30	-150
	10-20	15		10	-20	-200
	20-30	25		25	-10	-250
	30-40	35		30	0	0
	40-50	45		20	10	200
	50-60	55		10	20	200
				N = 100		$\sum f m = 200$

Assumed mean, A = 35 $\bar{X} = A + \frac{\sum fd}{N} = 35 - \frac{200}{100} = 33$

1.4.2 Properties of the Arithmetic Mean:

1. The sum of deviation from their arithmetic mean is always equal to zero.

Symbolically, $\sum (X - \overline{X}) = 0$

Ex 8:

Х	10	20	30	40	50	$\sum X = 150$
$X - \overline{X}$	-20	-10	0	10	20	$\sum X - \overline{X} = 0$

 $\bar{X} = \frac{\sum X}{N} = \frac{150}{5} = 30$

When we calculate the deviations of all the items from their arithmetic mean ($\overline{X} = 30$), we find that the sum of the deviations from the arithmetic mean i. e. $\sum (X - \overline{X}) = 0$

2. The sum of squared deviations of the items from arithmetic mean is minimum, that is, less than the sum of squared deviations of the items from any other value.

Ex	9:

Х	X - \overline{X}	$(X - 4)^2$
2	-2	4
3	-1	1
4	0	0

5	1	1
6	2	4
$\sum X = 20$	$\sum X - \overline{X} = 0$	$\sum (X - \overline{X})^2 = 0$

$$\bar{X} = \frac{\sum X}{N} = \frac{20}{5} = 4$$

The sum of the squared deviations is equal to 10 in the above example. If the deviations are taken from any other value the sum of the squared deviations are taken from any other value the sum of the squared deviations would be greater than 10.

Let us calculate the squares of the deviations of item from the value less than the arithmetic mean, say 3

Х	X - 3	$(X - 3)^2$
2	-1	1
3	0	0
4	1	1
5	2	4
6	3	9
$\sum X = 20$		$\sum (X - 3)^2 = 0$

3. Arithmetic mean is NOT independent of change of origin.

If each observation of a series is increased (or decreased) by a constant, then the mean of these observations is also increased (or decreased) by that constant.

4. Arithmetic mean is NOT independent of change of scale.

If each observation of a series is multiplied (or divided) by constant, then the mean of these observations is also multiplied (or divided) by that constant.

5. If arithmetic mean and number of items of two or more related groups are given, then we can compute the combined mean using the formula given below.

$$\bar{X}_{12} = \frac{N1\overline{X1} + N2\overline{X2}}{N1 + N2},$$

Where

 \bar{X}_{12} = Combined mean of two groups;

N1 = Number of items in the first group; N2 = Number of items in the second group

 $\overline{X1}$ = Arithmetic mean of the first group; $\overline{X2}$ = Arithmetic second group

 $\overline{X2}$ = Arithmetic mean of the

1.5 WEIGHTED ARITHMETIC MEAN

Arithmetic mean gives equal importance to all the items. When importance of the items are not same, in these cases we compute weighted arithmetic mean. The term weighted represents to the relative importance to the item.

$$\bar{X}_{w} = \frac{W_{1}X_{1} + WX_{2} + W_{3}X_{3} + \dots + W_{n}X_{n}}{W_{1} + W_{2} + \dots + W_{n}} = \frac{\sum WX}{\sum W}$$

Where

 \overline{X}_{w} represent the weighted arithmetic mean; X represent the variable values i. e. $X_{1}, X_{2}, \dots, X_{n}$

W represent the weights attached to the variable values i. e. $w_1, w_2 \dots w_n$ respectively.

To calculate weighted arithmetic mean, multiply the weight by the variable X and obtain the total $\sum WX$. Then divide this total by the sum of the weights, i.e. $\sum W$

In case of frequency distribution, if f_1, f_2, \ldots, f_n are the frequencies of the variable values X_1, X_2, \ldots, X_n respectively then the weighted arithmetic mean is given by

$$\bar{X}_{w} = \frac{W_{1}(f_{1} X_{1}) + W_{2}(f_{2} X_{2}) + \dots + W_{n}(f_{n} X_{n})}{W}$$
$$\bar{X}_{w} = \frac{\Sigma w(fX)}{\Sigma w}$$

Note: Simple arithmetic mean shall be equal to the arithmetic mean if the weights are equal.

Ex. 10 Calculate the weighted mean for following data.

X	1	2	5	7
W	2	14	8	32

Sol:

Х	W	WX
1	2	2
2	14	28
5	8	40
7	32	224
	$\sum W = 56$	$\sum WX = 294$

$$\bar{X}_{\rm w} = \frac{\sum WX}{\sum W} = \frac{294}{56} = 5.25$$

Ex. 11 Calculate the weighted mean for following data.

Wages per Day (X)	200	150	85
No. of workers (W)	25	20	10

Sol:

Х	W	WX
200	25	5000
150	20	3000
85	10	850
	$\sum W = 55$	$\sum WX = 8850$

 $\bar{X}_{\rm w} = \frac{\Sigma WX}{\Sigma W} = \frac{8850}{55} = 160.90$

Ex. 12. Calculate the weighted mean for following data and compare it with arithmetic mean

Subject	Weight	Student		
		X	Y	Ζ
Physics	2	72	42	52
Chemistry	3	75	52	62
Biology	5	58	88	68

Sol: For Student X,

Arithmetic Mean, $\bar{X}_{X} = \frac{\Sigma X}{3} = \frac{72+75+58}{3} = \frac{203}{3} = 67.67$ Weighted Arithmetic Mean, $\bar{X}_{WX} = \frac{\Sigma WX}{\Sigma W} = \frac{(2*72) + (3*75) + (5*58)}{2+3+5} = \frac{144+225+290}{10} = \frac{659}{10} = 65.9$

For Student Y, Arithmetic Mean, $\overline{X}_{y} = \frac{\sum X}{3} = \frac{42+52+88}{3} = \frac{200}{3} = 60.67$ Weighted Arithmetic Mean, $\overline{X}_{wY} = \frac{\sum WX}{\sum W} = \frac{(2*42) + (3*52) + (5*88)}{2+3+5} = \frac{84+156+440}{10} = \frac{680}{10} = 68$

For Student Z, Arithmetic Mean, $\bar{X}_Z = \frac{\sum X}{3} = \frac{52+62+68}{3} = \frac{182}{3} = 60.67$ Weighted Arithmetic Mean, $\bar{X}_{wZ} = \frac{\Sigma WX}{\Sigma W} = \frac{(2*52) + (3*62) + (5*68)}{2+3+5} = \frac{104+186+340}{10} = \frac{630}{10} = 63$

1.6 MEDIAN

Median is a middle value in the distribution. Median is a numeric value that separates the higher half of a set from the lower half. It is the value that the number of observations above it is equal to the number of observations below it. The median is thus a positional average.

For example, if the salary of five employees is 6100, 7150, 7250, 7500 and 8500 the median would be 7250.

When odd number of observations are there then the calculations of median is simple. When an even number of observations are given, there is no single middle position value and the median is taken to be the arithmetic mean of two middlemost items.

In the above example we are given the salary of six employees as 6100, 7150, 7250, 7500, 8500 and 9000, the median salary would be Median $=\frac{7250+7500}{2}=\frac{14750}{2}=7375$

Hence, in case of even number of observations median may be found by averaging two middle position values.

Calculations of Median – Individual Observations:

Arrange the data in ascending or descending order of magnitude.

In a group composed of an odd number of values, add 1 to the total number of values and divide by 2 gives median value. Median = size of $\frac{N+1}{2}$ th item

Ex. 13 From the following data, compute the median: 15, 9, 7, 23, 25, 25, 42, 25, 16, 14, 58, 25, 31

Sol: Arrange the numbers in ascending order 7, 9, 14, 15, 16, 23, 25, 25, 25, 31, 42, 58 Median = = size of $\frac{N+1}{2}$ th item = $\frac{13+1}{2}$ = 7th item = 25 \therefore Median = 25

The procedure for calculating median of an even numbered of items is not as above. The median value for a group composed of an even number of items is the arithmetic mean of the two middle values - i.e. adding two values in the middle and dividing by 2

Ex. 14 From the following data, compute the median: 451, 502, 523, 512, 622, 612, 754, 732, 701, 721 Sol: Arrange the numbers in ascending order 451, 502, 512, 523, 612, 622, 701, 721, 732, 754 Median = = size of $\frac{N+1}{2}$ th item = $\frac{10+1}{2}$ = 5.5th item Size of 5.5th item = $\frac{612 + 622}{2} = \frac{1234}{2}$ ∴ Median = 617

Calculations of Median – Discrete Series: Steps:

- 1. First arrange the data in ascending or descending order.
- 2. Find out the cumulative frequencies.
- 3. Apply formula: Median = size of $\frac{N+1}{2}$
- 4. Find out total in the cumulative frequency column which is equal to $\frac{N+1}{2}$ or next higher to that value and determine the value of the variable corresponding to it. That gives the median value.

Ex. 15 From the following data, find the value of median.

Income (Rs.)	450	500	630	550	710	580
No. of persons	29	31	21	25	11	35

Sol:

Income (Rs.)	No. of persons	Cumulative Frequency
Ascending order	f	c.f.
450	29	29
500	31	60
550	25	85
580	35	120
630	21	141
710	11	152

Median = = size of $\frac{N+1}{2}$ th item = $\frac{152+1}{2}$ = 76.5th item Size of 76.5th item = Rs. 550 It is median income.

Calculations of Median – Continuous Series:

The following formula is used to calculate median for continuous series.

Median = L +
$$\frac{N/2 - c.f.}{f}$$
 x i

L = Lower limit of median class; f = Simple freq. of the median class; c.f. = Cumulative freq. of the preceding the median class;

i= Class interval of the median class

Ex. 16 From the following data, find the value of median.

Marks		70-80	60-70	50-60	40-50	30-40	20-30	10-20
No.	of	10	15	26	30	42	31	24
students								

Sol: Arrange the data in ascending order

Marks	f	c.f.
10-20	24	24
20-30	31	55
30-40	42	97
40-50	30	127
50-60	26	153
60-70	15	168
70-80	10	178

Median = size of $\frac{N}{2}$ item = $\frac{178}{2}$ = 89th item Median lies in the class 30-40 (marked in pink) Median = L + $\frac{N/2 - c.f.}{f}$ x i L = 30. $\frac{N}{2}$ = 89, c.f. = 55, f = 42, i = 10 Median = 30 + $\frac{89-55}{42}$ x 10 = 30 + 8.09 = 38.09

1.7 MODE

The mode or the modal value is that value in a series of observations which occurs with the greatest frequency.

For example, the mode of the values 4, 6, 9, 6, 5, 6, 9, 4 would be 6.

Calculations of Mode – Discrete Series:

Ex. 17 From the following data, find the value of mode.

Size of cloth	28	29	30	31	32	33
No. of persons wearing	15	25	45	70	55	20

Sol: The mode or modal size is 31 because the value 31 occurred maximum number of times.

Calculations of Mode – Continuous Series:

The following formula is used to calculate mode for continuous series.

Mode = $L + \frac{f_1 - f_0}{2f_1 - f_0 - f_2} \mathbf{x} \mathbf{i}$, L = Lower limit of modal class; f_1 = freq. of the modal class; f_0 = freq. of the class preceding the modal class;

 f_2 = freq. of the class succeeding the modal class;

i= Class interval of the modal class

Ex. 18 From the following data, find the value of mode.

Marks	0-10	10-	20-	30-	40-	50-	60-	70-	80-	90-
		20	30	40	50	60	70	80	90	100
No. of	3	5	7	10	12	15	12	6	2	8
students										

Sol: After observing the table, modal class is 50-60

Mode =
$$L + \frac{f_1 - f_0}{2f_1 - f_0 - f_2} \times i$$
,
= $50 + \frac{15 - 12}{2x15 - 12 - 12} \times 10$
= $50 + \frac{3}{6} \times 10 = 55$

1.8 EMPIRICAL RELATION BETWEEN THE MEAN, MEDIAN, AND MODE

Karl Pearson has expressed the relationship between mean, median and mode as follows:

Mode = Mean - 3 [Mean - Median]

Mode = 3 Median - 2 Mean

If we know any of the two values out of the three, we can compute third from these relationships.

1.9 GEOMETRIC MEAN

Geometric mean of a set of n observations is the nth root of their product.

G. M. = $\sqrt[n]{(X_1)(X_2)(X_3) \dots \dots (X_n)}$. G. M. of 3 values 2, 4, 8 would be

G. M. = $\sqrt[3]{2X4X8} = \sqrt[3]{64} = 4$

For calculation purpose, take the logarithm of both sides

$$\log G. M = \frac{\log X_1 + \log X_2 + \dots + \log X_n}{N}$$
$$\log G. M = \frac{\sum \log X}{N} \quad \therefore G. M = \text{Antilog} \left[\frac{\sum \log X}{N}\right]$$
$$\text{In Discrete series, G. M. = Antilog} \left[\frac{\sum f \log X}{N}\right]$$
$$\text{In Continuous series, G. M. = Antilog} \left[\frac{\sum f \log m}{N}\right]$$

Calculations of Geometric Mean – Discrete/Individual Series:

Ex. 19 Daily income of ten families of a particular place is below. Calculate Geometric Mean.

85	70	15	75	500	8	45	250	40	36
----	----	----	----	-----	---	----	-----	----	----

Sol:

X	log X
85	1.9294
70	1.8451
15	1.1761
75	1.8751
500	2.6990
8	0.9031
45	1.6532
250	2.3979
40	1.6021
36	1.5563
	$\sum \log X 17.6373$

G. M. = Antilog $\left[\frac{\sum \log X}{N}\right]$

= Antilog
$$\left[\frac{\sum 17.6373}{10}\right]$$
 = Antilog (1.7637) = 58.03

Calculations of Geometric Mean – Continuous Series:

Ex 20. Calculate Geo	met	ric Me	ean from	n follow	ing data.

Marks	4-8	8-	12-	16-	20-	24-	28-	32-	36-
		12	16	20	24	28	32	36	40
Frequency	8	12	20	30	15	12	10	6	2

Sol:

Marks	m.p (m)	f	log m	f log m
4-8	6	8	0.7782	6.2256
8-12	10	12	1.0000	12.0000
12-16	14	20	1.1461	22.922
16-20	18	30	1.2553	37.6590
20-24	22	15	1.3424	20.1360
24-28	26	12	1.4150	16.9800
28-32	30	10	1.4771	14.7710
32-36	34	6	1.5315	9.1890
36-40	38	2	1.5798	3.156
		N=115		$\sum f \log m = 143.0386$

G. M. = Antilog
$$\left[\frac{\sum f \log m}{N}\right]$$

= Antilog $\left[\frac{\sum 143.0386}{115}\right]$
= Antilog (1.2438) = 17.53

1.10 HARMONIC MEAN

Harmonic mean of a number of observations, none of which is zero, is the reciprocal of the arithmetic mean of the reciprocals of the given values. Thus, harmonic mean (H. M.) of n observations x_i , i = 1, 2, ..., n is given by,

H. M.
$$= \frac{1}{\frac{1}{n} \sum_{1}^{n} \frac{1}{x_{i}}} = \frac{N}{\frac{1}{x_{1}} + \frac{1}{x_{2}} + \frac{1}{x_{3}} + \frac{1}{x_{n}}}$$

Calculations of Harmonic Mean – Individual Observations:

Ex. 21 Find the harmonic mean of 4, 36, 45, 50, 75.

Sol: H. M. =
$$\frac{N}{\frac{1}{x_1} + \frac{1}{x_2} + \frac{1}{x_3} + \frac{1}{x_n}} = \frac{5}{\frac{1}{4} + \frac{1}{36} + \frac{1}{45} + \frac{1}{50} + \frac{1}{75}} = \frac{5}{\frac{1}{3}} = 15$$

Calculations of Harmonic Mean – Discrete Series:

Formula for harmonic mean in Discrete series, H. M. = $\frac{N}{\sum \left[\int x \frac{1}{x} \right]} = \frac{N}{\sum \left[\frac{f}{x} \right]}$

Ex. 22 From the following data, Find the harmonic mean.

Marks	10	20	30	40	50
No. of students	20	40	60	30	10

Sol:

501.		
Marks X	f	f/X
10	20	2
20	40	2
30	60	2
40	30	0.75
50	10	0.20
	N = 160	$\sum f / X = 6.95$

H. M. $= \frac{N}{\sum \left[\frac{f}{X}\right]} = \frac{160}{6.95} = 23.0215$

Calculations of Harmonic Mean – Continuous Series:

Formula for harmonic mean in continuous series,

H. M. =
$$\frac{N}{\sum \left[\frac{f}{m}\right]}$$

Ex. 23 From the following data, compute the value of harmonic mean.

Class interval	10-20	20-30	30-40	40-50	50-60
Frequency	6	8	12	9	5

Sol:

Class Interval	Mid point (m)	f	f/m
10 - 20	15	6	0.40
20-30	25	8	0.32
30-40	35	12	0.3428
40 - 50	45	9	0.2
50 - 60	55	5	0.0909
		N = 40	$\sum f/m = 1.3537$

H. M.
$$= \frac{N}{\sum \left[\frac{f}{m}\right]} = \frac{40}{1.3537} = 29.54$$

1.11 RELATION BETWEEN THE ARITHMETIC, GEOMETRIC AND HARMONIC MEAN

Arithmetic mean is greater than geometric mean and geometric mean is greater than harmonic mean.

 $A.M. \ge G.M \ge H.M.$

The quality signs hold only if all the numbers X_1 , X_2 , X_3 ,..., X_n are identical.

1.12 ROOT MEAN SQUARE

The root mean square (RMS) is defined as the square root of the mean square (the arithmetic mean of the squares of a set of numbers). It is also called as the Quadratic average. Sometimes it is denoted by $\sqrt{\overline{X}^2}$ and given by,

$$\text{RMS} = \sqrt{\bar{X}^2} = \sqrt{\frac{\sum_{i=1}^n X_i^2}{N}} = \sqrt{\frac{\sum X^2}{N}}$$

It is very useful in fields that study sine waves like electrical engineering.

Ex. 24 Find RMS of 1, 3, 5, 7 and 9
Sol: RMS =
$$\sqrt{\frac{\Sigma X^2}{N}}$$

= $\sqrt{\frac{1^2 + 3^2 + 5^2 + 7^2 + 9^2}{5}} = \sqrt{\frac{1^2 + 3^2 + 5^2 + 7^2 + 9^2}{5}} = \sqrt{53} = 7.28$

1.13 QUARTILES, DECILES AND PERCENTILES

From the definition of median that it's the middle point which divides the set of ordered data into two equal parts. In the same way we can divide the set into four equal parts and this called quartiles. These values denoted by Q_1 , Q_2 and Q_3 , are called the first, second and the third quartile respectively. In the same way the values that divide the data into 10 equal parts are called deciles and are denoted by D_1 , D_2 , ..., D_9 whereas the values dividing the data into 100 equal parts are called percentiles and are denoted by P_1 , P_2 , ..., P_{99} . The fifth decile and the 50th percentile corresponds to median.

Formulae:

Quartile:

For individual observations, $Q_i = (\frac{i}{4})$. No. of observation, i = 1, 2, 3For discrete series, $Q_i = (\frac{i}{4})$. N, $N = \sum f$ and i = 1, 2, 3For continuous series, $Q_i = L + \frac{iN}{4} - cf}{f}$. c,

Where, i = 1, 2, 3, c = size of class interval.

L = Lower limit of the class interval in which lower quartile lies,

f = freq. of the interval in which lower quartile lies,

cf = cumulative freq. of the class preceding the quartile class,

Deciles:

For individual observations, $D_i = (\frac{i}{10})$. No. of observation, i = 1, 2, ..., 9For discrete series, $D_i = (\frac{i}{10})$. N, $N = \sum f$ and i = 1, 2, ..., 9For continuous series, $D_i = L + \frac{\frac{iN}{10} - cf}{f}$. c, i = 1, 2, ..., 9

Percentiles:

For individual observations, $P_i = (\frac{i}{100})$. value of observation, i = 1, 2, ..., 99For discrete series, $P_i = (\frac{i}{100})$. N, $N = \sum f$ and i = 1, 2, ..., 99For continuous series, $P_i = L + \frac{\frac{iN}{100} - cf}{f}$. c, i = 1, 2, ..., 99 **Ex. 25** Find the quartiles Q_1 , Q_3 , D_1 , D_5 , D_8 , P_8 , P_{50} and P_{85} of the following data 20, 30, 25, 23, 22, 32, 36.

Sol: Arrange data in ascending order, n = 7 i.e. odd number 20, 22, 23, 25, 30, 32, 36

$q_1 = (\frac{1}{4}).7 = 1.75$	$\therefore q_1 = 2$	$\therefore Q_1 = 22$
$q_3 = (\frac{3}{4}).7 = 5.75$	$\therefore q_2 = 6$	$\therefore Q_3 = 32$
$d_1 = (\frac{1}{10}).7 = 0.7$	\therefore d ₁ = 1	$\therefore D_1 = 20$
$d_5 = (\frac{5}{10}).7 = 3.5$	\therefore d ₅ = 4	∴D ₅ = 25
$d_8 = (\frac{8}{10}).7 = 5.6$	\therefore d ₈ = 6	$\therefore D_8 = 32$
$p_8 = (\frac{8}{100}).7 = 0.56$	$\therefore p_8 = 1$	$\therefore P_8 = 20$
$p_{50} = (\frac{50}{100}).7 = 3.5$	$\therefore p_{50} = 4$	$\therefore P_{50}=25$
$p_{85} = (\frac{85}{100}).7 = 5.95$	$\therefore p_{85} = 6$	$\therefore P_{85}=32$

Ex. 26 Find Q₁, Q₃, D₄, P₂₇ for the following data.

Х	0	1	2	3	4	5	6	7	8
f	1	9	26	59	72	52	29	7	1
c.f.	1	10	36	95	167	219	248	255	256

Sol. We know that, $Q_i = (\frac{i}{4})$. N

 $Q_1 = (\frac{1}{4}).256 = 64$ and c.f. just greater than 64 is 95. Hence $Q_1 = 3$ $Q_3 = (\frac{3}{4}).256 = 192$ and c.f. just greater than 192 is 219. Hence $Q_3 = 5$

 $\mathbf{D}_4 = (\frac{4}{10}).256 = 102.4$ and c.f. just greater than 102.4 is 167. Hence $\mathbf{D}_4 = \mathbf{4}$ $\mathbf{P}_{27} = (\frac{27}{100}).256 = 69.12$ and c.f. just greater than 69.12 is 95. Hence $\mathbf{P}_{27} = \mathbf{3}$

100^y 5 C

Ex. 27 Find Q_1 , Q_3 , D_2 , P_{90} for the following data.

Marks	Below	10-20	20-40	40-60	60-80	Above
	10					80
No. of students	8	10	22	25	10	5

Sol: We know that, $Q_i = L + \frac{\frac{iN}{4} - cf}{f}$. c,

Marks	Below 10	10-20	20-40	40-60	60-80	Above 80
f	8	10	22	25	10	5
cf	8	18	40	65	75	80

 Q_1 = Size of (N/4)th item = size of (80/4)= 20th item. Q_1 lies in the class 20-40.

L=20, N/4 = 20, cf = 18, f = 22 and c = 20

 $Q_1 = 20 + \{(20 - 18)/22\} * 20 = 20 + 1.82 = 21.82$

 Q_3 = Size of (3N/4)th item = size of (3*80/4)= 60th item. Q_3 lies in the class 40-60.

L=40, 3N/4 = 60, cf = 40, f = 25 and c = 20

 $Q_3 = 40 + \{(60 - 40)/25\} * 20 = 56$

 D_2 = Size of (2N/10)th item = size of (2*80/10)= 16th item. D_2 lies in the class 10-20

L=10,
$$2N/10 = 16$$
, cf=8, f = 10 and c=10

 $D_2 = 10 + \{(16 - 8)/10\} * 10 = 18$

 P_{90} = Size of (90N/100)th item = size of (90*80/10)= 72th item. P_{90} lies in the class 60-80.

L=60, 90N/100 = 72, cf=65, f = 10 and c=20

 $P_{90} = 60 + \{(72 - 65)/10\} * 20 = 74.$

1.14 SOFTWARE AND MEASURES OF CENTRAL TENDENCY

There are many software available to calculate measures of central tendency. We can use Excel to calculate the standard measures of central tendency (mean, median and mode). In Microsoft Excel, the mean can calculated by using one of the functions like AVERAGE, AVERAGEA, AVERAGEIF, AVERAGEIFS. The mean can be calculated by using the MEDIAN function. We can calculate a mode by using the MODE function, GEOMEAN to calculate geometric mean and HARMEAN to calculate harmonic mean.

We can use SPSS to calculate the standard measures of central tendency (mean, median and mode). We can get SPSS to compute mean, median and mode in the command submenu. Go to the Statistics menu, select the Analyse submenu, and then the Descriptive Statistics submenu and then the Frequencies option. We can use MINITAB to calculate the standard measures of central tendency using the functions Mean, Median, Mode ang GMEAN. To compute these go to Stat-Tables-Descriptive statistics. Using R software one could easily obtain the value of the mean using summary function.

We could find median value using summary function in R. The randomForest library can be used to impute the missing values using Median for numeric variables. Mode is used for missing value imputation for categorical variables using randomForest library in R. Model can be easily located graphically. You shouldn't be surprised that the R's mode function (mode ()) does not provide a model value. It shows the datatype of the particular variable which does not comply with our standard expectation. So how one would find mode using R software? We need to use table function for finding mode. As you know the table function in R provides frequency distribution of the variable. Thus the value with highest frequency is a modal value.

Geometric mean is the only average that is recommended for finding average growth (decline) rates. It is defined as the nth root of the product of n terms. Since it is defined in product terms so the observation shouldn't be having zero or negative values. We don't have a built-in function in R for its computation but one could find it by using its formula directly in R platform.

1.15 SUMMARY

A measure of central tendency is a measure that tells us where the middle of a group of data lies. Mean, median and mode are the most important measures of central tendency. The complete dataset may be represented by these values. It is not necessary for mean, median and mode to have the same values. Mean is sensitive to extreme data values. Median is a better way to understand skewed distribution than mean. It is possible that there is no mode in the data. Mean and median cannot be zero unless all data values are zero.

1.16 EXERCISE

X	10	30	50	70	89
f	7	8	10	15	10

- 1. Find the arithmetic mean of the following distribution:
- 2. Find the arithmetic mean of the following distribution:

Х	3	9	12	14	15	17
f	1	3	4	1	4	2

3. Find the arithmetic mean of the following data.

Class Interval	15-	25-	35-	45-	55-	65-	75-
	25	35	45	55	65	75	85
Frequency	6	11	7	4	4	2	1

4. Find the arithmetic mean of the following data.

Class Interval	10-20	20-30	30-40	40-50	50-60
Frequency	30	27	14	17	2

5. Obtain the median for the following frequency distribution:

Х	1	2	3	4	5	6	7	8	9
f	8	10	11	16	20	25	15	9	6

[Ans: Median = 5]

6. Obtain the median from the following data.

Х	20-25	25-30	30-35	35-40	40-45	45-50	50-55	55-60
f	35	45	70	105	90	74	51	30

7. Find the mode for the following distribution.

Marks	0-10	10-20	20-30	30-40	40-50	50-60	60-70	70-80
No. of	5	8	7	12	28	20	10	10
students								

[Ans: Mode = 46.67]

8. Calculate Geometric Mean from following data.

125	1462	38	7	0.22	0.08	12.75	0.5
[Ans: 6.	952]						

- 9. Find the geometric mean, harmonic mean and root mean square of the numbers 3, 5, 6, 6, 7, 10 and 12.
 [Ans: G. M. = 6.43, H. M. = 5.87, RMS = 7.55]
- 10. Find the arithmetic mean, geometric mean, harmonic mean of numbers 2, 4 and 8. Check the relation between them.
- 11. Calculate Quartile 3, Deciles -7 and Percentiles 20 from following data.

Class	2 - 4	4 – 6	6 – 8	8 - 10
Frequency	3	4	2	1

12. Calculate Q₁, Q₂, Q₃ D₁, D₅, D₉, P₁₁, P₆₅ from following data.

Wages	No. of employees
250.00 - 259.99	8
260.00 - 269.99	10

270.00 - 279.99	16
280.00 - 289.99	14
290.00 - 299.99	10
300.0 - 309.99	5
310.00 - 319.99	2

1.16 REFERENCES

- FUNDAMENTAL OF MATHEMATICAL STATISTICS by S. C. Gupta and V. K. Kapoor
- Statistical Methods by S. P. Gupta
- STATISTICS by Murray R. Spiegel, Larry J. Stephens



THE STANDARD DEVIATION AND OTHER MEASURES OF DISPERSION

Unit structure

- 2.0 Objectives
- 2.1 Introduction
- 2.2 Dispersion, or Variation
- 2.3 Range
- 2.4 Semi-Interquartile Range
- 2.5 Mean Deviation
- 2.6 10–90 Percentile Range
- 2.7 Standard Deviation
- 2.8 Short Methods for Computing the Standard Deviation
- 2.9 Properties of the Standard Deviation
- 2.10 Variance
- 2.11 Charlier's Check
- 2.12 Sheppard's Correction for Variance
- 2.13 Empirical Relations between Measures of Dispersion
- 2.14 Absolute and Relative Dispersion
- 2.15 Coefficient of Variation
- 2.16 Standardized Variable and Standard Scores
- 2.17 Software and Measures of Dispersion
- 2.18 Summary
- 2.19 Exercise
- 2.20 Reference

2.0 OBJECTIVES

After going through this chapter, students will able to learn

- To provide the importance of the concept of dispersion
- To calculate range, semi-Interquartile range, mean deviation
- To explain why measures of dispersion must be reported in addition to measures of central tendency
- To calculate standard deviation, variance, standard scores
- To trace precise relationship
- To compare two or more series with regard to their variability

2.1 INTRODUCTION

The measures of central tendency or Averages give us an idea of the concentration of the observations about the central part of distribution. But the average alone cannot adequately describe a set of observations. They must be supported and supplemented by some other measures, called Dispersion.

2.2 DISPERSION OR VARIATION

Literal meaning of dispersion is 'scatteredness'. In two or more distributions the central value may be the same but still there can be wide differences in the formation of distribution. Measures of dispersion help us in studying this important characteristic of a distribution.

Definitions of Dispersion:

- 1. "Dispersion is the measure of the variation of the items." A. L. Bowley
- "Dispersion is the measure of extent to which individual item vary." –
 L. R. Connor
- 3. "The degree to which numerical data tend to spread about an average value is called variation or dispersion of the data". Spiegel

2.3 RANGE

Range is the difference between two extreme observations of the distribution. Symbolically,

Range = L - S, where L = Largest item, S = smallest item

The relative measure corresponding to range, called the coefficient of range.

Coefficient of range $=\frac{L-S}{L+S}$

Since range is based on two extreme observations, it is not at all a reliable measure of dispersion.

Ex 1. From the following data, calculate range and coefficient of range.							
Day	Mon	Tues	Wed	Thurs	Fri	Sat	
Price	20	21	18	16	22	25	

Ex 1. From the following data, calculate range and coefficient of range.

Sol: Range = L - S = 25 - 16 = 9

Coefficient of range = $\frac{L-S}{L+S}$

 $=\frac{25-16}{25+16}=\frac{9}{41}=0.21$

For continuous series, find the difference between the upper limit of the highest class and the lower limit of the lowest class.

Ex 2. From the following data coefficient of range.

Marks	10-20	20 - 30	30-40	40-50	50-60
No. of Students	10	12	14	8	6

Sol: Coefficient of range $= \frac{L-S}{L+S}$

$$=\frac{60-10}{60+10}=\frac{50}{70}=0.21$$

2.4 SEMI-INTERQUARTILE RANGE OR QUARTILE DEVIATION

Semi-Interquartile Range Or Quartile Deviation is given by,

Q. D.
$$=\frac{Q_3 - Q_1}{2}$$

Quartile Deviation is a better measure than a range as it makes use of 50% of the data. But since it ignores the other 50% of the data, it cannot be considered as a reliable measure.

Q. D.
$$=\frac{Q_3 - Q_1}{2}$$

The relative measure corresponding to Q. D., called the coefficient of Q. D.

Coefficient of Q. D. = $\frac{Q_{3}-Q_{1}/2}{Q_{3}+Q_{1}/2} = \frac{Q_{3}-Q_{1}}{Q_{3}+Q_{1}}$

Coefficient of Q. D. can be used to compare the degree of variation in different distributions.

Computation of Quartile Deviation - Individual Observations:

Ex. 3 Find out Quartile Deviation and Coefficient of Quartile Deviation from following data.

25 33 45 17 35 20 55

Sol: Arrange the data in ascending order:

17 20 25 33 35 45 55

$$Q_1 = \text{size of } \left[\frac{N+1}{4}\right]$$
 th item = size of $\left[\frac{7+1}{4}\right]$ th item = 2nd item

 $\therefore Q_{1} = 20$ $Q_{3} = \text{size of } 3\left[\frac{N+1}{4}\right] \text{ th item} = \text{size of } 3\left[\frac{7+1}{4}\right] \text{ th item} = 6^{\text{th}} \text{ item}$ $\therefore Q_{3} = 45$ $Q. D. = \frac{Q_{3} - Q_{1}}{2} = \frac{45 - 20}{2} = 12.5$ Coefficient of Q. D. = $\frac{Q_{3} - Q_{1}}{Q_{3} + Q_{1}} = \frac{45 - 20}{45 + 20} = \frac{25}{65} = 0.455$

Computation of Quartile Deviation -Discrete Series:

Ex. 4 Find out Quartile Deviation and Coefficient of Quartile Deviation from following data.

Marks	10	20	30	40	50	60
No. of Students	7	10	18	12	10	6

Sol:

Marks	10	20	30	40	50	60
Frequency f	7	10	18	12	10	6
cf	7	17	35	47	57	63

 $Q_1 = \text{size of } \left[\frac{N+1}{4}\right]$ th item = size of $\left[\frac{63+1}{4}\right]$ th item = 16th item $\therefore Q_1 = 20$

 $Q_3 = \text{size of } 3\left[\frac{N+1}{4}\right]$ th item = size of $3\left[\frac{63+1}{4}\right]$ th item = 48^{th} item

$$\therefore Q_3 = 50$$

Q. D. $=\frac{Q_3 - Q_1}{2} = \frac{50 - 20}{2} = 15$

Coefficient of Q. D. $= \frac{Q_3 - Q_1}{Q_{3+Q_1}}$ $= \frac{50 - 20}{50 + 20} = \frac{30}{70} = 0.4285$

Computation of Quartile Deviation - Continuous Series:

Ex. 5 Find out Quartile Deviation and Coefficient of Quartile Deviation from following data.

Marks	35-44	45 - 54	55-64	65 - 74	75 - 84
No. of Students	12	40	33	13	12

Sol:

Marks	35-44	45 - 54	55-64	65 - 74	75 - 84
Frequency f	12	40	33	13	12
cf	12	52	75	88	100

Q₁ = size of
$$[\frac{N}{4}]$$
 th item = size of $[\frac{100}{4}]$ th item = 25th item
∴ Q₁ lies in the class 45 – 54
Q₁ = L + $\frac{N/4 - c.f.}{f}$ * i
L = 45, $N/4$ = 25, c.f. = 12 [c.f. of previous class], f= 40, i = 9
Q₁ = 45 + $\frac{25 - 12}{40}$ * 9 = 47.925
Q₃ = size of 3[$\frac{N}{4}$] th item = size of 3[$\frac{100}{4}$] th item = 75th item
∴ Q₃ lies in the class 55-64
Q₃ = L + $\frac{3N/4 - c.f.}{f}$ * i
L = 55, $\frac{3N}{4}$ = 75, c.f. = 52 [c.f. of previous class], f= 33, i = 9
Q₃ = 55 + $\frac{75 - 52}{33}$ * 9 = 61.2727
Q. D. = $\frac{Q_3 - Q_1}{2}$
= $\frac{61.2727 - 47.925}{2}$ = 6.67
Coefficient of Q. D. = $\frac{Q_3 - Q_1}{Q_{3+Q_1}}$
= $\frac{61.2727 - 47.925}{61.2727 + 47.925}$ = $\frac{6.67}{109.1977}$ = 0.061

2.5 MEAN DEVIATION

Mean deviation is also known as the average deviation.

If xi | fi, i = 1, 2, ..., n is the frequency distribution, then mean deviation from the average A (usually mean, median or mode).

Since mean deviation is based on all the observations, it is a better measure of dispersion than range and quartile deviation.

Note: Mean deviation is least when taken from median

The relative measure corresponding to the mean deviation called the coefficient of mean deviation and is obtained by,

Coefficient of M. D. = $\frac{M.D.}{Median}$

Computation of Mean deviation – Individual observations

M. D. = $\frac{1}{n} \sum |X - A|$

 $=\frac{1}{N}\sum |D|$, where |D| = |X - A| is the modulus value or absolute value of the deviation ignoring plus and minus signs.

Ex. 6 Calculate mean deviation and coefficient of mean deviation from following data:

600, 620, 640, 660, 680

Data	Deviation from median 640 D
600	40
620	20
640	0
660	20
680	40
N= 5	$\Sigma D = 120$

Sol: From above data, Median = 640

M. D. =
$$\frac{1}{N} \sum |D| = \frac{120}{5} = 24$$

Coefficient of M. D. = $\frac{M.D.}{Median}$

$$=\frac{24}{640}$$

= 0.037

Computation of Mean deviation – Discrete series:

M. D. =
$$\frac{1}{N} \sum f |D|$$
, where $|D| = |X - A|$

Ex. 7 Calculate mean deviation from following data.

Х	20	21	22	23	24
f	6	15	21	15	6

Sol:

Х	f	c.f	D	f D
20	6	6	2	12

21	15	21	1	15
22	21	42	0	0
23	15	57	1	15
24	6	63	2	12
	N = 63			$\sum f D = 54$

Median = size of $\frac{N+1}{2}$ th item = size of $\frac{63+1}{2}$ th item = 32th item

Size of 32th item is 22, hence Median = 22

M. D.
$$=\frac{1}{N}\sum f|D|$$

 $=\frac{54}{63}=0.857$

Computation of Mean deviation – Continuous series:

Here we have to obtain the mid-point of the various classes and take deviations of these points from median. Formula is same.

M. D. =
$$\frac{1}{N} \sum f |D|$$

Ex. 8 Calculate mean and mean deviation from following data.

Size	0-10	10-20	20-30	30-40	40-50	50-60	60-70
Frequency	7	12	18	25	16	14	8

Sol:

Size	f	c.f.	m.p (m)	m - 35.2	f D
				D	
0-10	7	7	5	30.2	211.4
10-20	12	19	15	20.2	242.4
20-30	18	37	25	10.2	183.6
30-40	25	62	35	0.2	5.0
40-50	16	78	45	9.8	156.8
50-60	14	92	55	19.8	277.2
60-70	8	100	65	29.8	238.4
	N=100				$\sum f D =1314.8$

Median = size of $\frac{N}{2}$ th item = size of $\frac{100}{2}$ th item = 50th item Median lies in the class 30 - 40

Median = L + $\frac{N_{2}-c.f.}{f}$ * i L= 30, N_{2} = 50, c.f. = 37, f = 25, i = 10 Median = 30 + $\frac{50-37}{25}$ *10 = 35.2 M. D. = $\frac{1}{N} \sum f |D|$ $=\frac{1314.8}{100}=13.148$

2.6 10-90 PERCENTILE RANGE:

The 10-90 percentile range of a set of data is defined by, 10 - 90 percentile range = $P_{90} - P_{10}$ Where P_{10} and P_{90} are the 10^{th} and 90^{th} for the data. Semi 10-90 percentile range = $\frac{P_{90} - P_{10}}{2}$

2.7 STANDARD DEVIATION:

Standard deviation is the positive square root of the arithmetic mean of the squares of the deviations of the given values from their arithmetic mean.

Standard deviation is also known as root mean square deviation as it is the square root of the mean of the standard deviation from arithmetic mean. Standard deviation is denoted by the small Greek letter σ (read as sigma).

Calculation of Standard Deviation - Individual Observations:

$$\sigma = \sqrt{\frac{\sum x^2}{N}}$$
, where $x = (X - \overline{X})$

Calculation of Standard Deviation - Discrete Series:

$$\sigma = \sqrt{\frac{\sum f x^2}{N}}$$
, where $x = (X - \overline{X})$

Calculation of Standard Deviation: Continuous Series:

$$\sigma = \sqrt{\frac{\sum f d^2}{N} - \left(\frac{\sum f d}{N}\right)^2} * i, \text{ where } d = \frac{(m-A)}{i}, i = \text{class interval}$$

Ex. 9 Calculate mean and standard deviation from the following data.

Size	0-10	10-20	20-30	30-40	40-50	50-60	60-70
Frequency	7	10	32	43	50	35	23

501:						
Marks	m. p. (m)	f	d = (m-35)/10	d^2	fd	fd ²
0-10	5	7	-3	9	-21	63
10-20	15	10	-2	4	-20	40
20-30	25	32	-1	1	-32	32
30-40	35	43	0	0	0	0
40-50	45	50	1	1	50	50
50-60	55	35	2	4	70	140

Sale

60-70	65	23	3	9	69	207
		N = 200			$\sum fd$ =116	$\sum_{532} fd^2 =$

Assumed mean, A=35

$$\overline{X} = A + \frac{\sum fd}{N} * i$$

$$= 35 + \frac{116}{200} * 10 = 40.8$$

$$\sigma = \sqrt{\frac{\sum fd^2}{N} - \left(\frac{\sum fd}{N}\right)^2} * i$$

$$= \sqrt{\frac{532}{200} - \left(\frac{116}{200}\right)^2} * 10$$

$$= \sqrt{2.66 - 0.3364} * 10 = 1.5243 * 10 = 15.243$$

2.8 SHORT METHODS FOR COMPUTING THE STANDARD DEVIATION:

Calculation of Standard Deviation- Individual Observations:

When actual mean is in fractions eg 568.245, it would be too bulky to do calculations. In such case either the mean may be approximated or the deviations be taken from assumed mean A. Following is formula if we take deviations from assumed mean A:

$$\sigma = \sqrt{\frac{\sum d^2}{N} - \left(\frac{\sum d}{N}\right)^2}$$
, where d = (X - A)

Ex. 10 Calculate standard deviation with the help of assumed mean.

340, 360, 390, 345, 355, 388, 372, 363, 277, 351

Sol: Consider assumed mean = 364

Х	d = (X - 364)	d^2
340	-24	576
360	-4	16
390	26	676
345	-19	361
355	-9	81
388	24	576
372	8	64
363	-1	1
377	13	169
351	-13	169
	$\sum d = 1$	$\sum d^2 = 2689$

$$\sigma = \sqrt{\frac{\sum d^2}{N} - \left(\frac{\sum d}{N}\right)^2} = \sqrt{\frac{2689}{10} - \left(\frac{1}{10}\right)^2} = 16.398$$

Calculation of Standard Deviation - Discrete Series:

Assumed mean method: $\sigma = \sqrt{\frac{\sum f d^2}{N} - \left(\frac{\sum f d}{N}\right)^2}$, where d = (X - A)

Ex. 11 Calculate standard deviation from the following data.

Size	3.5	4.5	5.5	6.5	7.5	8.5	9.5
Frequency	4	8	21	60	85	30	9

Sol:

Size	f	d = (X-6.5)	d^2	fd	fd^2
3.5	4	-3	9	-12	36
4.5	8	-2	4	-16	32
5.5	21	-1	1	-21	21
6.5	60	0	0	0	0
7.5	85	1	1	85	85
8.5	30	2	4	60	120
9.5	9	3	9	27	81
	N = 217			\sum fd = 123	$\Sigma fd^2 = 375$

Assumed mean, A= 6.5

$$: \sigma = \sqrt{\frac{\Sigma f d^2}{N} - \left(\frac{\Sigma f d}{N}\right)^2}$$
$$= \sqrt{\frac{375}{217} - \left(\frac{123}{217}\right)^2} = \sqrt{1.7281 - 0.3212} = 1.1861$$

2.9 PROPERTIES OF THE STANDARD DEVIATION

1. Combined standard deviation: We can compute combined standard deviation of two or more groups. It is denoted by σ_{12} and given by

$$\sigma_{12} = \sqrt{\frac{N_1 \sigma_1^2 + N_2 \sigma_2^2 + N_1 d_1^2 + N_2 \sigma_2^2}{N_1 + N_2}}$$

Where σ_{12} = combined standard deviation;

 σ_1 = standard deviation of first group;

 σ_2 = standard deviation of second group;

$$d_1 = |\overline{X_1} - \overline{X_{12}}|;$$

$$d_2 = |\overline{X_2} - \overline{X_{12}}|$$

2. The standard deviation of the first n natural numbers can obtained by, $\sigma = \sqrt{\frac{1}{12} (N^2 - 1)}$

Thus the standard deviation of natural numbers 1 to 20 will be

$$\sigma = \sqrt{\frac{1}{12} (20^2 - 1)} = \sqrt{\frac{1}{12} 399} = 5.76$$

3. Standard deviation is always computed from the arithmetic mean because the sum of the squares of the deviations of items from their arithmetic mean is minimum.

4. For normal distribution,

Mean $\pm 1 \sigma$ covers 68.27% of the items.

Mean $\pm 2 \sigma$ covers 95.45% of the items.

Mean $\pm 3 \sigma$ covers 99.73% of the items.

2.10 VARIANCE

The square of standard deviation is called the variance and is given by,

Variance = $\frac{\sum (X - \overline{X})^2}{N}$

i.e. Variance = σ^2 or $\sigma = \sqrt{Variance}$

In the frequency distribution where deviations are taken from assumed mean,

Variance =
$$\left\{\frac{\sum f d^2}{N} - \left(\frac{\sum f d}{N}\right)^2\right\}^*$$
 i², where d = $\frac{(X-A)}{i}$ and i = class interval

Ex. 12 Calculate standard deviation from the following data.

Marks	10-20	20-30	30-40	40-50	50-60	60-70
No. of students	2	6	8	12	7	5

Sol:

Marks	m.p (m)	f	d=(m-35)/10	d^2	fd	fd^2
10-20	15	2	-2	4	-4	8
20-30	25	6	-1	1	-6	6
30-40	35	8	0	0	0	0
40-50	45	12	1	1	12	12
50-60	55	7	2	4	14	28
60-70	65	5	3	9	15	45
		N =			$\sum fd =$	$\sum fd^2 = 99$
		40			31	

Variance =
$$\left\{\frac{\sum f d^2}{N} - \left(\frac{\sum f d}{N}\right)^2\right\}$$
* i²
= $\left\{\frac{99}{40} - \left(\frac{31}{40}\right)^2\right\}$ *10²
= (2.475 - 0.6006)*100 = 187.44

2.11 CHARLIE'S CHECK

Some error may be made while calculating the value of mean and standard deviations using different method. The accuracy of calculations can be checked by using following formulae.

$$\sum f (\mathbf{u}+1) = \sum f \mathbf{u} + \sum f = \sum f \mathbf{u} + \mathbf{N}$$

$$\sum f (\mathbf{u}+1)^2 = \sum f (\mathbf{u}^2 + 2\mathbf{u}+1) = \sum f \mathbf{u}^2 + 2 \sum f \mathbf{u} + \sum f = \sum f \mathbf{u}^2 + 2 \sum f \mathbf{u} + \mathbf{N}$$

$$\sum f (\mathbf{u}+1)^3 = \sum f \mathbf{u}^3 + 2 \sum f \mathbf{u}^2 + 2 \sum f \mathbf{u} + \mathbf{N}$$

$$\sum f (u+1)^{3} = \sum f u^{3} + 3 \sum f u^{2} + 3 \sum f u + N$$

Ex. 13 Use Charlier's check to verify mean and the standard deviation.

Size	20-30	30-40	40-50	50-60	60-70	70-80	80-90
Freq	9	12	8	10	11	35	15

Sol:

Х	f	m. p.	u=	u+1	f(u+1)	u ²	fu	fu ²
		(m)	(m-					
			55)/i					
20-30	9	25	-3	-2	-18	9	-27	81
30-40	12	35	-2	-1	-12	4	-24	48
40-50	8	45	-1	0	0	1	-8	8
50-60	10	55	0	1	10	0	0	0
60-70	11	65	1	2	22	1	11	11
70-80	35	75	2	3	105	4	70	140
80-90	15	85	3	4	60	9	45	135
	N=∑f				Σ		∑fu	$\sum fu^2$
	=100				f(u+1)=		=67	=423
					167			

 $\sum f (u+1) = 167$ $\sum f u + N = 67 + 100 = 167$ $\therefore \sum f (u+1) = \sum f u + N$

This provides the required check on the mean.

Х	f	m. p. (m)	u= (m-55)/i	u+1	f(u+1)	$f(u+1)^2$
20-30	9	25	-3	-2	-18	36
30-40	12	35	-2	-1	-12	12
40-50	8	45	-1	0	0	0

50-60	10	55	0	1	10	10
60-70	11	65	1	2	22	44
70-80	35	75	2	3	105	315
80-90	15	85	3	4	60	240
	N=				Σ	Σ
	∑f				f(u+1)	$f(u+1)^2$
	=10				= 167	==657
	0					

 $\sum f (u+1)^2 = 657$ $\sum f u^2 + 2 \sum f u + N = 423 + 2*67 + 100 = 657$ $\therefore \sum f (u+1)^2 = \sum f u^2 + 2 \sum f u + N$

This provides the required check on the standard deviation.

2.12 SHEPPARD'S CORRECTION FOR VARIANCE

The computation of the standard deviation is somewhat in error as a result of grouping the data into classes (grouping error). To adjust for grouping error, we use the formula,

Corrected variance = variance from grouped data - $\frac{i^2}{12}$

Where i is the class interval size. The correction $\frac{i^2}{12}$ is called Sheppard's correction. It is used for distribution of continuous variables where the tails tends to zero in both direction.

Ex. 14 Apply Sheppard's Correction to determine the standard deviation of the data in Ex. 8

Sol: $\sigma = 15.243 \div \sigma^2 = 232.349$ and i= 10.

Corrected variance = variance from grouped data - $\frac{i^2}{12}$ = 232.349 - $\frac{10^2}{12}$ = 224.016

Corrected Standard deviation = $\sqrt{224.016} = 14.9671$

2.13 EMPIRICAL RELATIONS BETWEEN MEASURES OF DISPERSION:

There is a fixed relationship between the three measures of dispersion in normal distribution.
Q. D. $=\frac{2}{3}\sigma$ or $\sigma = \frac{3}{2}$ Q. D and M. D. $=\frac{4}{5}\sigma$ or $\sigma = \frac{4}{5}$ M. D

The quartile deviation is smallest, the mean deviation next and the standard deviation is largest.

2.14 ABSOLUTE AND RELATIVE DISPERSION:

Measures of dispersion may be either absolute or relative. Absolute measures of dispersion are expressed in the same statistical unit in which the original data are given such as kilograms, tons, rupees etc. These values may be used to compare the variations in two distributions provided the variables are expressed in the same units and of the same average size. In case the two sets of data are expressed in different units such as quintals of sugar versus tons of sugarcane, the absolute measures of dispersion are not comparable. In such cases measures of relative dispersion is used.

A measure of relative dispersion is the ratio of a measure of absolute dispersion to an appropriate average. It is sometimes called coefficient of dispersion.

Relative dispersion = $\frac{absolute \ dispersion}{average}$

2.15 COEFFICIENT OF VARIATION:

Coefficient of is used in problems where we want to compare the variability of two or more than two series. That series or group for which the coefficient of variation is greater is said to be more variable or less consistent, less uniform, less stable or less homogeneous. The series for which the coefficient of variation is less is said to be less variable or more consistent, more uniform, more stable or more homogeneous.

If the absolute dispersion is standard deviation σ and if average is the mean \overline{X} , then relative dispersion is called coefficient of variation, it is denoted by C. V. and is given by,

Coefficient of variation (C.V.) = $\frac{\sigma}{\bar{x}} \times 100$

Ex. 15 Calculate arithmetic mean, standard deviation and coefficient of variation.

Class	23-	28-	33-	38-	43-	48-	53-	58-	63-	68-
	27	32	37	42	47	52	57	62	67	72
Freq	2	6	7	12	18	13	9	7	4	2

Class	m. p. (m)	f	d = (m-	d^2	fd	fd ²
			50)/5			
23-27	25	2	-5	25	-10	50
28-32	30	6	-4	16	-24	96
33-37	35	7	-3	9	-21	63
38-42	40	12	-2	4	-24	48
43-47	45	18	-1	1	-18	18
48-52	50	13	0	0	0	0
53-57	55	9	1	1	9	9
58-62	60	7	2	4	14	28
63-67	65	4	3	9	12	36
68-72	70	2	4	16	8	32
		N = 80			$\sum fd = -44$	$\sum fd^2 = 380$

Mean
$$(\bar{X}) = A + \frac{\sum fd}{N} x i$$

 $= 50 + \frac{-44}{80} x 5 = 47.25$
S. D. $(\sigma) = \sqrt{\frac{\sum fd^2}{N} - (\frac{\sum fd}{N})^2} x i$
 $= \sqrt{\frac{380}{80} - (\frac{-44}{80})^2} x 5$
 $= \sqrt{4.75 - 0.3025} x 5 = 22.23$
C. V. $= \frac{\sigma}{\pi} x 100$

$$= \frac{22.2375}{47.25} \times 100 = 47.06\%$$

2.16 STANDARDIZED VARIABLE AND STANDARD SCORES

75

The variable that measures the deviation from the mean in units of the standard deviation is called standardized variable, is independent of the units used and is given by,

$$Z = \frac{X - \bar{X}}{\sigma}$$

If the deviations from the mean are given in units of the standard deviation, they are said to be expressed in standard units or standard scores. These are of great value in the comparison of distribution. The variable z is often used in educational testing, where it is called as a standard score.

Ex. 16 Your test score is 160 while the test has a mean of 120 and standard deviation of 15. If the distribution is normal, what is your z score? Explain the meaning of the result.

Sol:
$$z = \frac{X - \bar{X}}{\sigma} = \frac{160 - 120}{15} = 2.7$$

The score is 2.7 standard deviations above the mean.

Ex. 17 A student received a grade of 84 on a final examination in English for which mean grade was 76 and the standard deviation was 10. On a final examination in Science for which mean grade was 82 and the standard deviation was 16, she received a grade of 90. In which subject was her relative standing higher?

Sol: Standardized variable $z = \frac{X - \bar{X}}{\sigma}$ For English, $z = \frac{84 - 76}{10} = 0.8$ For Science, $z = \frac{90 - 82}{16} = 0.5$

Thus, the student had a grade of 0.8 of a standard deviation above the mean in English but only 0.5 of a standard deviation above the mean in science. Thus her relative standing was higher in English.

2.17 SOFTWARE AND MEASURES OF DISPERSION:

The statistical software gives a variety of measures for dispersion. The dispersion measures are usually given in descriptive statistics. EXCEL and MINITAB allows for the computation of all the measures discussed above. The output from MINITAB and STATISTIX has helped clarify some of the statistical concepts which are hard to understand without some help from the graphics involved.

Calculating Range In Excel: Excel does not offer a function to compute range. However, we can easily compute it by subtracting the minimum value from the maximum value. The formula would be =MAX()-MIN() where the dataset would be the referenced in both the parentheses. The =MAX() and =MIN() functions would find the maximum and the minimum points in the data. The difference between the two is the range. Microsoft Excel has two functions to compute quartiles. The inter-quartile range has to be calculated as the difference between the quartile 3 and quartile 1 values Quartiles can be calculated using =QUARTILE.INC() or =QUARTILE.EXC(). Both functions calculate the quartiles by calculating the percentiles on the data. Excel offers two functions, =STDEV.S() for sample standard deviation, and =STDEV.P() for population standard deviation.Excel with two different functions: =VAR.P() for population variance, and =VAR.S() for sample variance. Minitab may be used to

compute descriptive statistics for numeric variables, including the mean, median, mode, standard deviation, variance and coefficient of variance. To compute these go to Stat-Tables-Descriptive statistics.

You can use SPSS to calculate the measures of dispersion such as range, semi-interquartile range, standard deviation and variance. We can get SPSS to compute these in the command submenu. Go to the Statistics menu, select the Analyse submenu, and then the Descriptive Statistics submenu and then the Frequencies option. We can use MINITAB to calculate the measures of dispersion the functions Q1, Q3, Range StDev, Variance and CorfVar

2.18 SUMMARY

A measure of dispersion indicates the scattering of data. Dispersion is the extent to which values in a distribution differ from the average of the distribution. The measure of dispersion displays and gives us an idea about the variation and the central value of an individual item. The range and interquartile range are generally ineffective to measure the dispersion of set of data. The useful measure that describes the dispersion of all the values is standard deviation or variance. Dispersion can prove very effective in association with central tendency in making any statistical decision.

2.19 EXERCISE

1. Calculate Quartile deviation (Q. D.), Mean Deviation (M. D.) from mean for the following data.

Marks	0-10	10-20	20-30	30-40	40-50	50-60	60-70
No. of Students	6	5	8	15	7	6	3

[Ans: Q.D. = 11.23, Mean = 33.4, M.D from mean = 13.184]

2. Calculate Mean Deviation (M. D.) from mean for the following data

Size	2	4	6	8	10	12	14	16
f	2	2	4	5	3	2	1	1

[Ans: Mean = 8, M.D from mean = 2.8]

3 Calculate Mean Deviation and its coefficient from mean for the following data.

Size	0-10	10-20	20-30	30-40	40-50	50-60	60-70	70 -80
Freq	5	8	12	15	20	14	12	6

[Ans: Median = 43, M.D = 15.37, Coe. Of M. D. = 0.357]

- 4. Find the standard deviation of the following data.
- i. 12, 6, 7,3,15, 10, 18, 5
- ii. 9, 3, 8, 8, 9, 8, 9, 18

[Ans: i. St. dev. σ = 4.87, ii. St. dev. σ = 3.87]

5. Find the standard deviation of the following data.

Age	20-25	25-30	30-35	35-40	40-45	45-50
No. of persons	170	110	80	45	40	35

Take assumed average = 32.5

[Ans: Standard deviation σ = 7.936]

6. Calculate the standard deviation from the following data by short method.

240.12, 240.13, 240.15, 240.12, 240.17, 240.15, 240.17, 240.16, 240.22, 240.21

7. Calculate standard deviation from the following data by short method.

Salary	45	50	55	60	65	70	75	80
No. of persons	3	5	8	7	9	7	4	7
[A no: Standard deviation $= 10.25$]								

[Ans: Standard deviation = 10.35]

8. Calculate arithmetic mean, standard deviation and coefficient of variation.

Class	20-25	25-30	30-35	35-40	40-45
Frequency	1	22	64	10	3

[Ans: \overline{X} = 32.1, S. D. (σ) =3.441, C. V. = 10.72]

2.20 REFERENCE

- FUNDAMENTAL OF MATHEMATICAL STATISTICS by S. C Gupta and V. K. Kapoor
- Statistical Methods by S. P. Gupta
- STATISTICS by Murray R. Spiegel, Larry J. Stephens

INTRODUCTION TO R

Unit structure

- 3.0 Objectives
- 3.1 Introduction
- 3.2 Basic syntax
- 3.3 Data types
- 3.4 Variables
- 3.5 Operators
- 3.6 Control statements
- 3.7 R-functions
- 3.8 R-Vectors
- 3.9 R lists
- 3.10 R Arrays
- 3.11 Summary
- 3.12 Exercise
- 3.13 References

3.0 OBJECTIVES

After going through this chapter, students will able to learn

- 1. Understand the different data types, variables in R.
- 2. Understand the basics in R programming in terms of operators, control statements
- 3. Use of built-in and user defined function
- 4. Understand the different data structures in R.

3.1 INTRODUCTION

R is programming language and software environment for statistical computing and graphics. It is an open source programming language. It was designed by Ross Ihaka and Robert Gentleman at the University of Auckland, New Zealand.in 1993. It was released on 31-Oct-2014 by the R Development Core Team. It is widely used by researchers from diverse disciplines to estimate and display results and by teachers of statistics and research methods. Today, millions of analysts, researchers, and brands such as Facebook, Google, Bing, Accenture, and Wipro are using R to solve complex issues. The applications of R are not limited to just one sector, we can see the use of R in banking, e-commerce, finance, and many more sectors

It is freely available on <u>www.r-project.org</u> or can also download from CRAN (Comprehensive R Archive Network) website <u>http://CRAN.R-project.org</u>.

R Command Prompt:

We will be using RStudio. Once we have R environment setup, then it's easy to start R command prompt by just typing the following command at command prompt – $\$

\$ R

This will launch R interpreter and you will get a prompt > where you can start typing your program

> "Hello, World!"

[1] "Hello, world!"

Usually, we will write our code inside scripts which are called RScripts in R.

Write the below given code in a file

1 print ("Hello, World!")

and save it as myfirstprogrm.R and then run it in console by writing:

Rscript myfirstprogram.R

It will produce following output

[1] "Hello, World!"

3.2 BASIC SYNTAX

Any program in R is made up of three things: Variables, Comments, and Keywords. Variables are used to store the data.

Comments are used to improve code readability. They are like helping text in your R program. Single comment is written using # in the beginning of the statement.

Eg. # This is my first R program

Keywords are reserved words that hold a specific meaning to the compiler. Keyword cannot be used as a variable name, function name.

Following are the Reserved words in R: if, else, while, repeat, for, function, in, next, break, TRUE, FALSE, NULL, Inf, NaN, NA, NA_integer_, NA_real_, NA_complex_, NA_character etc

We can view these keywords by using either help(reserved) or ?reserved R is case sensitive language.

3.3 DATA TYPES

Variables can store data of different types and different types can do different things. Variables are the reserved memory location to store values. As we create a variable in our program some space is reserved in memory.

Data type	Example	Description
Numeric	50, 25.65, 999	Decimal values
Logical	True, False	Data with only two possible values
		which can be constructed as true/false
Character	'A', "Excellent",	A character is used represent string
	'50.50'	values.
Integer	5L, 70L, 9876L	L tells R to store the value as an
		integer.
Complex	X = 5 + 4i	A complex value in R defined as the
		pure imaginary value i.
Raw		A raw data type is used to holds raw
		bytes.

Following are the data types used in R programming.

We can use the class () function to check the data type of a variable.

numeric a < - 25.5 class (a) # complex a < - 10+5i class (a) # integer a < - 100L class (a) # logical/boolean a < - TRUE class (a) # character/string a < - "I am doing R programming" class (a) output: [1] "numeric"

[1] "complex"

[1] "integer"

[1] "logical"

[1] "character"

3.4 VARIABLES

Variables are used to store the information to be manipulated in the R program. A variable in R can store an atomic vector, group of atomic vectors or a combination of many R-objects. A valid variable name consists of letters, numbers and the dot or underline characters. The variable name must start with a letter or the dot not followed by a number. Ex - valid - a, a, b, ab, a1, a1, a.c

Invalid - 2a, _a

R does not have a command for declaring a variable. A variable is created the moment you first assign a value to it.

In R, the assignment can be denoted in three ways:

- 1. = (Simple Assignment)
- 2. <- (Leftward Assignment)
- 3. -> (Rightward Assignment)

name = "Ajay"

gender < - "Male"

age < - 25

Here, name, gender and age are variables and "Ajay", "Male", 25 are values.

To print/output variable, you do not need any function. You can just type the name of the variable.

name = "Ajay" O/P:

[1] "Ajay" #auto print the value of name variable

However, R have a print() and cat() functions which are used to print the value of the variable. The cat() function combines multiple values into a continuous print output.

Cat ("My name is", name, "\n")

Cat ("my age is", age, "\n")

O/P: My name is Ajay

My age is 25

ls() function: To know all the variables currently available in the workspace, use the ls() function.

using equal to operator a = "Good morning" # using leftward operator b < - "Good morning" # using leftward operator "Good morning - > c print(ls()) O/P: "a" "b" "c" # List the variables starting with the pattern "var". > print(ls(pattern="var")) The variables starting with dot (.) are hidden, they can be listed using "all.names=TRUE" argument to ls() function.

```
> print(ls(all.name=TRUE))
```

rm() function: This is a built in function used to delete an wanted variables.

> rm(variable)

using equal to operator

a = "Good morning"

using leftward operator

b < - "Good morning"

```
# using leftward operator
```

```
"Good morning - > c
```

Removing variable

rm(a)

print(a)

O/P: Error in print(a) : object 'a' not found

All the variables can be deleted by using the **rm()** and **ls()** function together.

```
>rm(list=ls())
```

> print(ls())

3.5 OPERATORS

Operators are the symbols directing the compiler to perform various kinds of operations between the operands. There are different types of

operator, and each operator performs a different task. Operators simulate the various mathematical, logical, and decision operations performed on a set of Complex Numbers, Integers, and Numericals as input operands.

Types of Operators used in R programming:

- Arithmetic Operators
- Relational Operators
- Logical Operators
- Assignment Operators
- Miscellaneous Operators

Arithmetic Operators:

Arithmetic operators are used with numeric values to perform common mathematical operations

<-,=	Assignment	A < - 5 ; b=10
+	Addition	$x \le c(2,5.5,6); y \le c(8,3,4);$
		print(x+y) # O/P [1] 10.0 8.5 10.0
-	Subtraction	$x \le c(2, 5.5, 6); y \le c(8, 3, 4);$ print(x - y)
*	Multiplication	x <- c(2,5.5,6); y <- c(8, 3, 4); print(x*y)
/	division	$x \le c(2,5.5,6); y \le c(8,3,4);$ print(x/y)
%%	remainder	x<- c(2,5.5,6); y <- c(8, 3, 4); print(x%%y) #O/P [1] 2.0 2.5 2.0
%/%	gives quotient	x <- c(2,5.5,6); y <- c(8, 3, 4) ;print(x%/%y) # O/P 0 1 1
<u>∧ **</u>	exponent	x <- c(2,5.5,6) ; y <- c(8, 3, 4); print(x^y) #O/P 256.000 166.375 1296.000

Relational Operators: Relational/Comparison operators are used to compare two values

>	Greater than	$x \le c(2,5.5,6,9); y \le c(8,2.5,14,9);$
		print(x>y)
		# O/P [1] FALSE TRUE FALSE FALSE
<	Less than	$x \le c(2,5.5,6,9); y \le c(8,2.5,14,9);$
		print(x < y);
		#O/P [1] TRUE FALSE TRUE FALSE

<=	Less than equal to	$x \le c(2,5.5,6,9); y \le c(8,2.5,14,9)$
		print(x<=y)
		#O/P [1] TRUE FALSE TRUE TRUE
>=	Greater than	$x \le c(2,5.5,6,9); y \le c(8,2.5,14,9)$
	equal to	print(x>=y)
		#O/P [1] FALSE TRUE FALSE TRUE
==	Equal	x < -c(2,5.5,6,9); y < -c(8,2.5,14,9)
		print(x==y)
		#O/P [1] FALSE FALSE FALSE TRUE
!=	Not equal	$x \le c(2,5.5,6,9); y \le c(8,2.5,14,9)$
		print(x!=y)
		# O/P [1] TRUE TRUE TRUE FALSE

Logical Operators: Logical operators are used to combine conditional statements.

&	Element wise Logical AND	x <- c(3, 1, TRUE, 2+3i); y <- c(4, 1, FALSE, 2+3i) print(x&y); # O/P [1] TRUE TRUE FALSE TRUE
	Element wise Logical OR	x <- c(3, 0, TRUE, 2+2i); y <- c(4, 5, FALSE, 2+3i) print(x y) # O/P [1] TRUE TRUE TRUE TRUE
!	Element wise Logical NOT	x<- c(8, 0, FALSE, 4+4i); print(!x) # O/P [1] FALSE TRUE TRUE FALSE
&&	Takes first element of both the vectors and gives the TRUE only if both are TRUE.	x <- c(3,0,TRUE, 8+9i); y<- c(1,3,TRUE, 3+4i) print(x&&y) # O/P [1] TRUE
	Logical OR operator. It returns TRUE if one of the statement is TRUE.	x <- c(4, 0,TRUE, 8+9i); y<- c(3, 5, TRUE, 2+3i) print(x y) # O/P [1] TRUE

Miscellaneous Operators: Miscellaneous operators are used to anipulate data:

:	Create a series of numbers in sequence	x <- 2:9 print(x) # [1] 2 3 4 5 6 7 8 9
%in%	Find out if an element belongs to a vector	x <- 8; y <- 12; z <- 1:10 print(x %in% z); print(y %in% z) # O/P [1] TRUE [1] FALSE

3.6 CONTROL STATEMENTS

Control statements are expressions used to control the execution and flow of the program based on the conditions provided in the statements.

if condition: if statement checks the expression provided in the parenthesis is true or not true. The block of code inside if statement will be executed only when the expression evaluates to be true.

Syntax:

```
if (expression ) {
```

// statements will execute if expression is true.

```
}
a < - 500
if (a > 100) {
print((x, "is greater than 100"))
}
O/P: [1] "500 is greater than 100"
```

If else condition: If expression evaluates to be true, then the if block of code will be executed, otherwise else block of code will be executed.

Syntax:

```
if(expression){
```

// statements will execute if expression is true.

}

 $else\{$

// statements will execute if expression is false.

```
}
```

```
a < - 500
```

if (a > 100) {

print(a, "is greater than 100")

} else {

print(a, "is smaller than 100")

}

O/P: [1] "500 is greater than 100"

Repeat loop: Repeat loop executes the same code again and again until stop condition met.

Syntax: repeat { commands if (condition){ break } } a <- 1 repeat { print(a) a = a+1if (a>5) break } } O/P: [1] 1 [1] 2 [1] 3 [1] 4 [1] 5

return statement: return statement is used to return the result of an executed function and returns control to the calling function. Syntax:

```
return(expression)
```

Example: func < - function(a) {</pre>

if(a > 0){

return ("POSITIVE")

```
else if (a < 0)
```

```
return("NEGATIVE")
```

}else{
return("ZERO")
}
fun(1)
fun(0)

fun(-1)O/P : "POSITIVE" "NEGATIVE"

"ZERO"

next statement: next statement is useful when we want to skip the current iteration of a loop without terminating it.

Syntax:

next

Example:

a < - 1:8

#Print even numbers

for(i in a){

```
if(i%%2 !=0){
```

next

```
}
```

print(i)

O/P:

[1] 2

}

[1] 4

[1] 6

[1] 8

break statement: The break keyword is a jump statement that is used to terminate the loop at a particular iteration

Syntax:

if (test expression) {

break

}

switch Statement: A switch statement is a selection control mechanism. Switch case is a multiway branch statement. It allows a variable to be tested for equality against a list of values. If there is more than one match for a specific value, then the switch statement will return the first match found of the value matched with the expression.

Syntax:

switch(expression, case1, case2, case3,)

Example:

```
a < - switch( 2, "Nagpur", "Mumbai", "Delhi", "Raipur")
```

print(a)

O/P:

[1] "Mumbai"

while loop: The while loop executes the same code again and again until stop condition is met.

Syntax:

```
While (test_expression) {
Statement
}
Example:
a < - c("Hello", "World")
count < - 1
while (count < 5) {
print(a)
count = count + 1
}
O/P
[1] "Hello" "World"
[1] "Hello" "World"</pre>
```

[1] "Hello" "World"

for loop: The for loop can be used to execute a group of statements repeatedly depending upon the number of elements in the object. It is an entry controlled loop, in this loop the test condition is tested first, then the body of the loop executed, the loop body would not be executed if the test condition is false.

```
Syntax:
for (value in vector) {
 statements
}
Example:
v < - LETTERS[1:5]
for ( x in v
}) {</pre>
```

print(x) O/P: [1] "A" [1] "B" [1] "C" [1] "D" [1] "E" Example: for (x in c(-5, 8, 9, 11)) { print(x) } O/P: [1] -5 [1] 8 [1] 9 [1] 11

Nested for-loop: Nested loops are similar to simple loops. Nested means loops inside loop. R programming allows using one loop inside another loop. In loop nesting, we can put any type of loop inside of any other type of loop. For example, a if loop can be inside a for loop or vice versa. Moreover, nested loops are used to manipulate the matrix. for (i in 1:3)

```
{
```

}

```
for (j in 1:i)
         {
                print( i * j)
        }
O/P:
[1] 1
[1] 2
[1] 4
[1] 3
[1] 6
[1] 9
```

3.7 R-FUNCTIONS

A set of statements which are organized together to perform a specific task is known as a function. A function is a set of statements organized together to perform a specific task. R has a large number of in-built functions and the user can create their own functions. An R function is created by using the keyword **function**.

```
Syntax :
```

```
Function_name < - function(arg1, arg2, .....)
{ function body
```

The different components of function are -

- Function name is the actual name of the function.
- An argument is placeholder. In function, argument are optional means a function may or may not contain arguments, and these arguments have default values also.
- The function body contains a set of statements which defines what the function does.
- Return value is the last expression in the function body which is to be evaluated.
- R also has two types of function, i.e. Built in function and user defined function.

Built-in function: The functions which are already defined in the programming framework are known as built in functions. Simple examples of built-in functions are seq(), mean(), amx(), sum(), paste(...) etc. They are directly called by user written programs.

print(seq(50, 60)) O/P: [1] 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60 print(mean(30, 40)) O/P: [1] 35

User defined Function: R allows us to create our own function in our program. They are specific to what a user wants and once created they can be used like built-in functions.

Example:

```
areaofCircle < - function(radius){
area = pi*radius^2
return(area)
}
print(areaofCircle(2))
O/P: [1] 12.56637</pre>
```

Example:

```
# create a function without an argument.
a.function < - function (){
for(i in 1:5) {
       b < -i^{2}
print(b)
}
}
# call the function a function without supplying an argument
a.function()
O/P
[1] 1
[1] 4
[1] 9
[1] 16
[1] 25
# create a function with an argument.
a.function < - function (a){
for(i in 1:a) {
       b < -i^{2}
print(b)
}
}
# call the function a function without supplying 5 as an argument
a.function(5)
O/P
[1] 1
[1] 4
[1] 9
[1] 16
[1] 25
```

Calling a function with argument values:

```
#Create a function with argument
a.function < - function(x,y,z) {
esult < - x * y + z
print( result)
}
# call the function by position of arguments
a.function(4, 2, 10)
# call the function by names of the arguments
a.function(x=10, y=4, z=2)
O/P:
[1] 18
```

[1] 42

Calling a function with default argument:

#Create a function with argument a.function < - function(x = 5, y= 7) { result < - x * y print(result) } # call the function without giving any arguments a.function() # call the function with giving new values of the argument. a.function(10, 6) O/P: [1] 35 [1] 60

3.8 R –VECTORS

A vector is a basic data structure. In R, a sequence of elements which share the same data type is known as vector. A vector supports logical, integer, double, character, complex, or raw data type. A vector length is basically the number of elements in the vector, and it is calculated with the help of the length() function.

Vector is classified into two parts, i.e., Atomic vectors and Lists. There is only one difference between atomic vectors and lists. In an atomic vector, all the elements are of the same type, but in the list, the elements are of different data types. The elements which are contained in vector known as components of the vector. We can check the type of vector with the help of the typeof() function.

Creation of atomic vector

Single Element Vector: when you write just one value, it becomes a vector of length 1.

print("xyz") print(25.5) print(TRUE) O/P [1] "xyz" [1] 25.5 [1] TRUE

Multiple Elements vector:

1. Using the colon(:) operator: # Creating a sequence from 1 to 8 v<-1:8 print(v) # Creating a sequence from 1.5 to 8.5 v<-1.5:8.5 print(v) O/P: [1] 1 5 2 3 4 6 7 8 [1] 1.5 2.5 3.5 4.5 5.5 6.5 7.5 8.5 2. Using sequence (seq) operator: # Create a vector from 1 to 5 incrementing by 0.6 print(seq(1, 5, by = 0.6))O/P [1] 1.0 2.8 3.4 4.0 4.6 1.6 2.2 3. Using the c() function: The non character values are converted to character type if one of the elements is a character. x < - c('mango', 'yellow', 10, TRUE) print(x) O/P "10". [1] "mango" "yellow", "TRUE" Accessing Vector Elements: Elements of a Vector are accessed using indexing. The [] brackets are used for indexing. Indexing starts with position 1. Giving a negative value in the index drops that element from result. TRUE, FALSE or 0 and 1 can also be used for indexing. # Accessing vector elements using position x < - c("Jan", "Feb", "Mar", "April", "May", "Jun", "July", "Aug", "Sept", "Oct", "Nov", "Dec") a < -x[c(2,4,8)]print(a) # Accessing vector elements using logical indexing. b < -x[c(TRUE, TRUE, FALSE, FALSE, TRUE, FALSE, FFALSE, TRUE, TRUE, FALSE, FALSE)] print (b) # Accessing vector elements using negative indexing. c < -x[c(-1, -3, -4, -8, -9, -10, -11)]print(c) # Accessing vector elements using 0/1 indexing. d < -x[c(0,1,0,0,1,0,0,0,0,0,0,1)]print(d) O/P [1] "Feb" "April" "Aug" "Feb" "May" "Sept" "Oct" [1] "Jan" [1] "Feb" "May" "Jun" "July" "Dec"

```
[1] "Feb" "May" "Dec"
```

Vector Manipulation:

Vector arithmetic: Two vectors of same length can be added, subtracted, multiplied or divided giving the result as a vector output.

#create two vectors. x < -c(2,7,3,4,0,10)y < -c(3,10,0,7,1,1)add.result < -x + yprint(add.result) multi.result < -x * yprint(multi.result) O/P: [1] 5 3 11 17 11 1 70 0 28 0 10 [1] 6 Vector Element Recycling: If we apply arithmetic operations to two vectors of unequal length, then the elements of the shorter vector are recycled to complete the operations. x < -c(2,7,3,4,0,10)y < -c(3,10)# y becomes c(3,10,3,10,3,10) add.result < -x + yprint(add.result) 20O/P: [1] 5 14 3 17 6 Vector Element Sorting: Elements in a vector can be sorted using the sort() function. x < -c(2,7,3,-11,4,0,210)sort.result < - sort(x)print(sort.result) resort.result < - sort(x, decreasing - TRUE) print(resort.result) O/P: [1] -11 0 2 3 4 7 210 4 2 210 7 3 0 -11 [1]

3.9 R – LISTS

Lists are heterogeneous data structures. Lists are the R objects which contain elements of different types. These are also one-dimensional data structures. A list can be a list of vectors, list of matrices, a list of characters and a list of functions and so on. List is created using list() function.

Creating a List:

#Create a list containing strings, numbers, vectors)
list_1 < - list("Apple", "Mango", 25.25, 60.5, c(16,25,36))
print(list_1)</pre>

O/P: [[1]] "Apple" [[2]] [1] "Mango" [[3]] [1] 25.25 [[4]] [1] 60.5 [[5]] [1] 16 25 36

Naming List Element: The list elements can be given and they can be accessed using these names.

list_1 < - list(c("Mon", "Tues", "Wed"), matrix(c(1,2,3,4,5,6), nrow = 2)

```
#Give names to the elements in the list.
```

```
names(list_1) < - c("Days of Week", "Matrix")
print(list_1)</pre>
```

O/P

\$Days of Week [1] "Mon" "Tues" "Wed" **\$Matrix** [,1] [,2] [,3] 3 [1,] 1 2 5 6 [2,] 4

Accessing List Elements:

Elements of the list can be accessed by the index of the element in the list. In case of named lists it can also be accessed using the names.

list_1 < - list(c("Mon", "Tues", "Wed"), matrix(c(1,2,3,4,5,6), nrow = 2)
names(list_1) < - c("Days of Week", "Matrix")
print(list_1[1])</pre>

print(list_1\$Matrix)
O/P:

\$Days of Week

[1] "N	Mon"	"Tues"		"Wed"
	[,1]	[,2]	[,3]	
[1,]	1	2	3	
[2,]	4	5	6	

Manipulating List Elements:

We can add, delete and update list elements as shown below. We can add and delete elements only at the end of a list. But we can update any element.

list_1 < - list(c("Mon", "Tues", "Wed"), matrix(c(1,2,3,4,5,6), nrow = 2)
names(list_1) < - c("Days of Week", "Matrix")
 #add element at the end of the list
 list_1[3] < - "Add Element"
 print(list_1[4])
 O/P</pre>

[[1]]

[1] "Add Element"

Merging Lists:

You can merge many lists into one list by placing all the lists inside one list() function.

```
list_a < - list(1,2)
list_b < - list("Ankit", "Pooja")
#merge tow lists
merged.list < - c(list_a, list_b)
print(merged.list)</pre>
```

```
[[1]]
[1] 1
```

[[2]] [1] 2

[[3]]

[1] Ankit

[4]]

[1] Pooja

Converting List to vector:

A list can be converted to a vector so that the elements of the vector can be used for further manipulation. All the arithmetic operations on vectors can be applied after the list is converted into vectors.

```
list_a < - list(10:13)
print(list_a)
list_b < - list(20:23)
print(list_b)
#Convert the lists to vectors
x1 < - unlist(list_a)
x2 < - unlist(list_b)
print(x1)
print(x2)
add < -x1 + x2</pre>
```

	print(O/P	add)	
[[1]] [1] 10 [[2]]	11	12	13
[1] 20 [1] 10 [1] 20 [1] 20 [1] 30	21 11 21 32	22 12 22 34	23 13 23 36

3.10 R ARRAYS

Arrays are the R data objects which can store data in more than two dimensions. In R, an array is created with the help of the array() function. This array() function takes a vector as an input and to create an array it uses vectors values in the **dim** parameter.

#create two vectors of different lengths

v1 < -c(1,2,3)v2 < -c(4,5,6,7,8,9)

#Take these vectors as input to the array

array1 < - (c(v1,v2), dim = c(3,3,2))

print(array1)

O/P

, , 1

		[,1]	[,2]	[,3]
	[1,]	1	4	7
	[2,]	2	5	8
	[3,]	3	6	9
, , 2				×
		[,1]	[,2]	[,3]
	[1,]	1	4	7
	[2,]	2	5	8
	[3,]	3	6	9

Naming Columns and Rows: We can give names to the rows, columns and matrices in the array by using the dimnames parameter. #create two vectors of different lengths

v1 <- c(1,2,3) v2 <- c(4,5,6,7,8,9) column.names <- c("Col1", "Col2", "Col3") row.names <- c("Row1", "Row2", "Row3") matrix.names <- c("Matrix1", "Matrix2") array1 < - array(c(v1,v2),dim = c(3,3,2), dimnames = list(row.names, column.names, matrix.names)) print(array1) O/P: ,, Matrx1

		Col1	Col2	Col3
	Row1	1	4	7
	Row2	2	5	8
	Row3	3	6	9
, , Mat	rix2			
		Col1	Col2	Col3
	Row1	1	4	7
	Row2	2	5	8
	Row3	3	6	9

Accessing Array Elements:

#create two vectors of different lengths v1 < -c(1,2,3)v2 < -c(4,5,6,7,8,9)column.names <- c("Col1", "Col2", "Col3") row.names <- c("Row1", "Row2", "Row3") matrix.names <- c("Matrix1", "Matrix2") array1 < - array(c(v1,v2),dim = c(3,3,2), dimnames = list(row.names,column.names, matrix.names)) #Print the second row of the second matrix of the array. print(array1[2,2]) #Print the element in the first row and 3rd column of the first matrix. print(array1[1,3,1]) #Print the first Matrix print(array1[, ,1]) O/P Col1 Col2 Col3 2 5 8 [1] 7 Col1 Col2 Col3 Row1 4 1 7 5 2 8 Row2

Manipulating Array Element: As array is made up matrices in multiple dimensions, the operations on elements of array are carried out by accessing elements of the matrices. #create two vectors of different lengths

9

v1 < -c(5,9,3) v2 < -c(10,11,12,13,14,15)#Take these vectors as input to the array array1 < -array(c(v1,v2),dim = c(3,3,2))#create two vectors of different lengths V3 < -c(9,1,0)

Row3

3

6

```
V4 < -c(6,0,11,3,14,1,2,6,9)
array2 < - array(c(v3,v4),dim = c(3,3,2))
#create matrices from these arrays
matrix 1 < - array 1[, , 2]
matrix 2 < - array 2[, , 2]
#add the matrices
add1 < - matrix1+matrix2
print(add1)
O/P:
               [,1]
                       [,2]
                               [,3]
       [1,]
                10
                        20
                               26
       [2,]
                18
                        22
                               28
                6
                        24
                               30
       [3,]
```

3.11 SUMMARY

R is world's most widely used statistics programming language. It is the 1 choice of data scientists R is taught to solve critical business applications. In addition, R is a full-fledged programming language, with a rich complement of mathematical functions, matrix operations and control structures. It is very easy to write your own functions. In this chapter we covered basic programming to different types of data objects of R with suitable examples in simple and easy steps.

3.12 EXERCISE

```
1. Find the output of following code.
1) b= "15"
a = switch(b)
"5"="Hello A",
          "10"="Hello B",
          "15"="Hello C",
      "20"="Hello D")
          print (a)
2) a= 1
b = 2
y = switch (a+b, "Hello, A", "Hello B", "Hello C", "Hello D")
print (y)
3) # Create vegetable vector
vegetable <- c('Potato', 'Onion', 'Brinjal', 'Pumpkin')
for (x in vegetable) {
print(x)
}
```

```
4) for ( i in c (5, 10, 15, 20, 0, 25)
        if (i == 0)
        ł
               break
print (i)
}
print("outside loop")
5) for ( i in c (5, 10, 15, 20, 0, 25)
{
        if (i == 0)
               next
print (i)
}
print("outside loop")
6) a < - 10
b<- 14
count=0
if(a<b)
{
  cat(a,"is a smaller number\n")
  count=1
}
if(count==1){
  cat("Block is successfully execute")
}
7) a <-1
b<-24
count=0
while(a < b){
               cat(a,"is a smaller number\n")
                a=a+2
                if(x = 15)
        break
8) a < -24
if(a%%2==0){
               cat(a," is an even number")
}
if(a%%2!=0){
                       cat(a," is an odd number")
9) x <- c("Hardwork", "is", "the", "key", "of", "success")
```

```
if("key" %in% x) {
    print("key is found")
} else {
    print("key is not found")
}
10) Rectangle = function(l=6, w=5){
    area = 1 * w
    return(area)
}
print(Rectangle(3, 4))
print(Rectangle(w = 9, 1 = 3))
print(Rectangle())
```

3.13 REFERENCES

- The Art of R Programming: A Tour of Statistical Software Design by Norman Matloff
- Beginning R The Statistical Programming Language by Mark Gardener
- https://www.javatpoint.com/
- <u>https://www.ict.gnu.ac.in/content/r-programming</u>

• <u>https://www.geeksforgeeks.org/</u>

UNIT II

4

MOMENTS, SKEWNESS, AND KURTOSIS

Unit Structure

- 4.1 Objective
- 4.2 Introduction
- 4.3 Moments

4.3.1 Moments for Grouped Data.

4.3.2 Relations between Moments.

4.3.3 Computation of Moments for Grouped Data.

- 4.4 Charlie's Check and Sheppard's Corrections.
- 4.5 Moments in Dimensionless Form
- 4.6 Skewness
- 4.7 Kurtosis
- 4.8 Software Computation of Skewness and Kurtosis.
- 4.9 Summary
- 4.10 Exercise
- 4.11 List of References

4.1 **OBJECTIVE**

After going through this unit, you will able to :

- Define Moments and calculate for ungroup and group data.
- Explain types of moments.
- Find relation between raw and central moment.
- Used Charlier's check method in computing moments by coding method.
- Define Sheppard's correction for moments.
- Define moments in dimensional form.
- Define Skewness and Kurtosis.
- Calculate moments, Skewness and Kurtosis using software.

4.2 INTRODUCTION

The measure of central tendency (location) and measure of dispersion (variation) both are useful to describe a data set but both of them fail to tell anything about the shape of the distribution. We need some other certain measure called the moments to identify the shape of the distribution known as Skewness and Kurtosis.Moments are statistical measures that give certain characteristics of the distribution. Moments provide sufficient information to reconstruct a frequency distribution function. Moments are a set of statistical parameters to measure a distribution. Four moments are commonly used:1st moments for Average, 2nd for Variance, 3rd for Skewness and 4th moment for Kurtosis.

4.3 MOMENTS

The arithmetic mean of the rth power of deviations taken either from mean, zero or from any arbitrary origin is called moments. Assume there is sequence of random variables $x_1, x_2, x_3, \dots, x_n$. The first sample moment, usually called the average is defined by first moments. Three types of moments are defined as follow:

When the deviations are computed from the arithmetic mean, then such moments are called moments about mean (mean moments) or sometimes called central moments, denoted by μ_r and given as follows:Hence for ungroup data,

- i) The first moment about A, as $\mu_1 = \frac{\sum (x \bar{x})}{n}$.
- ii) The second moment about A, $as\mu_2 = \frac{\sum (x-\bar{x})^2}{n}$.
- iii) The third moment about A, $as\mu_3 = \frac{\sum (x \bar{x})^3}{n}$.
- iv) The fourth moments about A, as $\mu_4 = \frac{\sum (x \bar{x})^4}{n}$

When the deviations of the values are computed from any arbitrary value (provisional mean) say A, then such moments are called moments about arbitrary or provisional mean denoted by $\mu_r(a)$. Hence for ungroup data,

- i) The first moment about A, as $\mu_1(a) = \frac{\sum (x-A)}{n}$.
- ii) The second moment about A, $as\mu_2(a) = \frac{\sum (x-A)^2}{n}$.
- iii) The third moment about A, $as\mu_3(a) = \frac{\sum (x-A)^3}{n}$.
- iv) The fourth moments about A, as $\mu_4(a) = \frac{\sum (x-A)^4}{n}$

When the deviations of the values are computed from the origin or zero, then such moments are called moments about origin or raw moments denoted by $\mu_r(a)$

i) The first moment about origin, as $\mu'_1 = \frac{\Sigma(x)}{n}$.

ii) The second moment about origin, $as\mu'_2 = \frac{\sum (x)^2}{n}$.

iii) The third moment about origin, $as\mu'_3 = \frac{\Sigma(x)^3}{n}$.

iv) The fourth moments about origin, as $\mu'_4 = \frac{\sum(x)^4}{n}$ Example 1: Find raw moments for the following data: 5, 8, 12, 4, 6. Solution:

x	<i>x</i> ²	<i>x</i> ³	<i>x</i> ⁴
5	25	125	625
8	64	512	4096
2	4	8	16
4	16	64	256
6	36	216	1296
$\sum x = 25$	$\sum x^2 = 145$	$\sum x^3 = 925$	$\sum x^4 = 6289$

i) The first moment about origin, as $\mu'_1 = \frac{\sum(x)}{n} = \frac{25}{5} = 5$.

ii) The second moment about origin,
$$as\mu'_2 = \frac{\Sigma(x)^2}{n} = \frac{145}{5} = 29.$$

iii) The third moment about origin,
$$as\mu'_3 = \frac{\sum(x)^3}{n} = \frac{925}{5} = 185$$
.

iv) The fourth moments about origin, as
$$\mu'_4 = \frac{\Sigma(x)^4}{n} = \frac{6289}{5} = 1257.8$$

4.3.1 Moments for Grouped Data:

1. Moments about arbitrary point:Let x represents a variable occurring with frequency f, in a given distribution, then the i^{yh} moment $\mu_i(a)$ about A is defined as

$$\mu_i(a) = \frac{\sum f(x-A)^i}{N}$$
, where $N = \sum f$.

We generally find moments up to i = 4. \therefore we can write :

- i) The first moment about A, as $\mu_1(a) = \frac{\sum f(x-A)}{N}$.
- ii) The second moment about A, as $\mu_2(a) = \frac{\sum f(x-A)^2}{N}$.
- iii) The third moment about A, as $\mu_3(a) = \frac{\sum f(x-A)^3}{N}$.
- iv) The fourth moments about A, as $\mu_4(a) = \frac{\sum f(x-A)^4}{N}$.

Example 2: For the following distribution find all four moments about 5.

Х	2	4	6	8	10
F	4	6	12	5	3

Solution: let prepared table first,

x	f	(x - 5)	f(x - 5)	$f(x-5)^2$	$f(x-5)^{3}$	$f(x-5)^{5}$
2	4	-3	-12	36	-108	324
4	6	-1	-6	6	-6	6
6	12	1	12	12	12	12
8	5	3	15	45	135	405
10	3	5	15	75	375	1875
Total	30		24	174	408	2622

Moments about arbitrary A = 5 is given by The first moment about A, as $\mu_1(a) = \frac{\sum f(x-A)}{N} = \frac{24}{30} = 0.8$. The second moment about A, as $\mu_2(a) = \frac{\sum f(x-A)^2}{N} = \frac{174}{30} = 5.8$. The third moment about A, as $\mu_3(a) = \frac{\sum f(x-A)^3}{N} = \frac{408}{30} = 13.6$. The fourth moments about A, as $\mu_4(a) = \frac{\sum f(x-A)^4}{N} = \frac{2622}{30} = 87.4$.

2. Moments about mean (Central moments):

These are moments about the Arithmetic Mean \bar{x} . Hence when A is taken as \bar{x} , we obtain these moments. Thus it is given by

i) The first moment about \bar{x} , as

$$\mu_1 = \frac{\sum f(x - \bar{x})}{N}$$

ii) The second moment about \bar{x} , as

$$\mu_2 = \frac{\sum f(x - \bar{x})^2}{N}$$

iii) The third moment about \bar{x} , as

$$\mu_3 = \frac{\sum f(x - \bar{x})^3}{N}$$

iv) The fourth moments about \bar{x} , as

$$\mu_4 = \frac{\sum f(x - \bar{x})^4}{N}$$

From the definition of the mean \bar{x} and the standard deviation σ , it immediately follows that $\mu_1 = 0$, $\mu_2 = \sigma^2$ and μ_3 measure the asymmetry of the curve. These moments are important study the nature of the distribution.

Example 3: Find the central moments for the following distribution:

X	1	2	3	4	5
F	2	5	6	5	2

Solution:

x	f	fx	$(x-\bar{x})$	f	f	f(x	f(x)
				$(x-\bar{x})$	(<i>x</i>	$(-\bar{x})^{3}$	$(-\bar{x})^4$
					(x^{2})		
1	2	2	-2	-4	8	-16	32
2	5	10	-1	-5	5	-5	5
3	6	18	0	0	0	0	0
4	5	20	1	5	5	5	5
5	2	10	2	4	8	16	32
Total	20	60		0	26	0	74

Here, $\bar{x} = \frac{\sum fx}{N} = \frac{60}{20} = 3.$

Therefore, the central moments are given by

i) The first moment about
$$\bar{x}$$
, as

$$\mu_1 = \frac{\sum f(x - \bar{x})}{N} = \frac{0}{20} = 0 \; .$$

ii) The second moment about \bar{x} , as

$$\mu_2 = \frac{\sum f(x-\bar{x})^2}{N} = \frac{26}{20} = 1.3 \; .$$

iii) The third moment about \bar{x} , as

$$\mu_3 = \frac{\sum f(x - \bar{x})^3}{N} = \frac{0}{20} = 0 \; .$$

iv) The fourth moments about \bar{x} , as

$$\mu_4 = \frac{\sum f(x-\bar{x})^4}{N} = \frac{74}{20} = 3.7.$$

3. Moments about origin(Raw moments):

As the name suggests, taking A as the origin (A = 0), we get these moments. Thus it is given by

- i) The first moment about Origin, as $\mu_1' = \frac{\sum fx}{N} \, .$
- ii) The second moment about Origin, as $\mu_2' = \frac{\sum f x^2}{N}.$

$$\iota_2' = \frac{\Delta J^{\chi}}{N}$$

- iii) The third moment about Origin, as $\mu_3' = \frac{\sum fx^3}{N}$. iv) The fourth moments about Origin, as
- $\mu_4' = \frac{\sum fx^4}{N}$. Note that for first moment about origin is mean of the data.

Example 4: Find	the ra	w mom	ents for the	e following o	lata:

X	-1	0	1	2	3	4
F	2	4	3	7	3	1

Solution : lets prepared table

	<u> </u>	1	1		1
x	f	fx	fx^2	fx^3	fx^4
-1	2	-2	2	-2	2
0	4	0	0	0	0
1	3	3	3	3	3
2	7	14	28	56	112
3	3	9	27	81	243
4	1	4	16	64	256
Total	20	28	76	202	616

Therefore, the raw moments are given by

i) The first moment about Origin, as

$$\mu_1' = \frac{\sum fx}{N} = \frac{28}{20} = 1.4$$
.

ii) The second moment about Origin, as

$$\mu_2' = \frac{\sum f x^2}{N} = \frac{76}{20} = 3.8$$
.

iii) The third moment about Origin, as

$$\mu_3' = \frac{\sum fx^3}{N} = \frac{202}{20} = 10.1$$

iv) The fourth moments about Origin, as

$$\mu_4' = \frac{\sum fx^4}{N} = \frac{616}{20} = 30.8.$$

4.3.2 Relations between Moments:

We studied three different types of moments. Now it is very useful to simplifying relation between them. We will now give inter-relation between various moments and solve example using these relations.

Relation between moments about arbitrary point and the central moment:

i)
$$\mu_1 = \mu_1(a) - \mu_1(a) = 0$$

ii)
$$\mu_2 = \mu_2(a) - \mu_1(a)$$

ii) $\mu_2 = \mu_2(a) - \mu_1(a)^2$ iii) $\mu_3 = \mu_3(a) - 3\mu_1(a)\mu_2(a) + 2\mu_1(a)^3$

iv)
$$\mu_4 = \mu_4(a) - 4\mu_1(a)\mu_3(a) + 6\mu_1(a)^2\mu_2(a) - 3\mu_1(a)^4$$

Conversely the moments $\mu_i(a)$'s about A in term of μ_i 's are given as follows:

i)
$$\mu_1(a) = \bar{x} - A$$

ii) $\mu_2(a) = \mu_2 + \mu_1(a)^2$
iii) $\mu_3(a) = \mu_3 + 3\mu_2\mu_1(a) + \mu_1(a)^3$
iv) $\mu_4(a) = \mu_4 + 4\mu_3\mu_1(a) + 6\mu_2\mu_1(a)^2 + \mu_1(a)^4$

Relation between Raw moments and central moments:

Recall that, the raw moments μ'_i are obtained from the general moments $\mu_i(a)$ when A is taken as '0'.

Hence taking A as '0' and replacing $\mu_i(a)$ by corresponding μ'_i in the formula, we get

i)
$$\mu_1 = \mu'_1 - \mu'_1 = 0$$

ii) $\mu_2 = \mu_2' - {\mu_1}'^2$
iii) $\mu_3 = {\mu_3}' - 3\mu_1'\mu_2' + 2{\mu_1}'^3$
iv) $\mu_4 = {\mu_4}' - 4{\mu_1}'{\mu_3}' + 6{\mu_1}'^2{\mu_2}' - 3{\mu_1}'^4$

Conversely the moments μ'_i in term of μ_i are given as follows:

i) $\mu_1' = \bar{x}$ ii) $\mu_2' = \mu_2 + {\mu_1}'^2$ iii) $\mu_3' = \mu_3 + 3\mu_2\mu_1' + {\mu_1}'^3$ iv) $\mu_4' = \mu_4 + 4\mu_3\mu_1' + 6\mu_2{\mu_1}'^2 + {\mu_1}'^4$

Example 5:The first four central moments of a distribution are 0, 3, 5, 10. If the mean of the distribution is 2, find the moments about 3.

Solution: We have A = 3, $\bar{x} = 2$, $\mu_1 = 0$, $\mu_2 = 3$, $\mu_3 = 5$, $\mu_4 = 10$. Using the relation between central moments and arbitrary moments,

- i) $\mu_1(a) = \bar{x} A = 2 3 = -1.$
- ii) $\mu_2(a) = \mu_2 + \mu_1(a)^2 = 3 + (-1)^2 = 4$
- iii) $\mu_3(a) = \mu_3 + 3\mu_2\mu_1(a) + \mu_1(a)^3 = 5 + 3(3)(-1) + (-1)^3 = -5.$

iv)
$$\mu_4(a) = \mu_4 + 4\mu_3\mu_1(a) + 6\mu_2\mu_1(a)^2 + \mu_1(a)^4$$

= 10 + 4(5) + 6(3)(-1)^2 + (-1)^4 = 49.

Example 6: The first four raw moments about the origin are 2, 12, 74 and 384. Find the mean \bar{x} and the first four central moments.

Solution: We already define the raw moments about the origin i.e. $\mu_i(a)$'s with A = 0. Given that $\mu_1(a) = 2$, $\mu_2(a) = 12$, $\mu_3(a) = 74$, and $\mu_4(a) = 384$, with A = 0.

Therefore, Mean = $\bar{x} = \mu_1(a) + A = 2 + 0 = 2$.

Using the relation between raw moments and central moments

- i) $\mu_1 = \mu_1(a) \mu_1(a) = 2 2 = 0$
- ii) $\mu_2 = \mu_2(a) \mu_1(a)^2 = 12 2^2 = 8$

iii)
$$\mu_3 = \mu_3(a) - 3\mu_1(a)\mu_2(a) + 2\mu_1(a)^3 = 74 - 3(2)(12) + 2(2^2) = 10$$

iv)
$$\mu_4 = \mu_4(a) - 4\mu_1(a)\mu_3(a) + 6\mu_1(a)^2\mu_2(a) - 3\mu_1(a)^4$$
$$= 284 - 4(2)(74) + 6(2^2)(12) - 3(2^4)$$
$$= 384 - 592 + 288 - 48 = 128.$$

Example 7: The first four central moments for a distribution are 0,3,0 and 7. If the mean \bar{x} of the distribution is 4, find the first four raw moments.

Solution: The raw moments are the moments about origin. Given that $\mu_1 = 0, \mu_2 = 3, \mu_3 = 0$ and $\mu_4 = 7$ with $\bar{x} = 4$.

Using the relation between central moments and raw moments.

i) $\mu'_1 = \bar{x} = 4$ ii) $\mu'_2 = \mu_2 + {\mu_1}'^2 = 3 + 4^2 = 17.$ iii) $\mu'_3 = \mu_3 + 3\mu_2\mu'_1 + {\mu_1}'^3 = 0 + 3(3)(4) + 4^3 = 100.$ $\mu'_4 = \mu_4 + 4\mu_3\mu'_1 + 6\mu_2{\mu_1}'^2 + {\mu_1}'^4$
4.3.3 Computation of Moments for Grouped Data:

We have already found mean and standard deviation for the continuous data (grouped data). Now to calculate moments for the continuous data we used coding method (Short method).

When the values of x are not consecutive, but equally spaced at an interval of length'c'. We need to divide the expression by 'c'. It is called change of scale by 'c'.

Where we take x = a + cu or $u = \frac{x-a}{c}$.

We give below the effect of change of origin and scale on moments.

Let x = a + cu,

 $\therefore \bar{x} = a + c\bar{u}.$

i)The moments of x about A are given by

$$\mu_i(a) = \frac{\sum f u^i}{N} \times c^i$$

ii) The central moments of x are given by

$$\mu_i = \frac{\sum f(u - \bar{u})^i}{N} \times c^i$$

Note: When A = 0, we get the raw moment.

Example 8: Find the central moments for the following data:

Class interval	0-20	20-40	40-60	60-80
Frequency	4	7	6	3

C.I	F	Cla	u	F	<i>(u</i>	f(u	f(u	f(u	f(u
		SS	$=\frac{x-a}{a}$	u	- 0.4)	- 0.4)	$(-0.4)^2$	$(-0.4)^3$	$(-0.4)^4$
		Mar	C						
		ks							
		(x)							
0-	4	10	-1	-4	-1.4	-5.6	7.84	-10.976	15.3664
20									
20-	7	30	0	0	-0.4	-2.8	3.92	-0.448	0.1792
40									
40-	6	50	1	6	0.6	3.6	2.16	1.296	0.7776
60									
60-	3	70	2	6	1.6	4.8	7.68	12.288	19.6608
80									
Tot	2			8		0	21.6	2.16	35.984
al	0								

Solution: first find mean by coding method, taking a = 30

Here, $\bar{u} = \frac{\sum fu}{N} = \frac{8}{20} = 0.4.$

The central moments of x are given by

$$\mu_{i} = \frac{\sum f(u-\bar{u})^{i}}{N} \times c^{i}$$
i) $\mu_{1} = \frac{\sum f(u-\bar{u})^{1}}{N} \times c^{1} = \frac{0}{20} \times 20 = 0.$
ii) $\mu_{2} = \frac{\sum f(u-\bar{u})^{2}}{N} \times c^{2} = \frac{21.6}{20} \times 20^{2} = 432.$
iii) $\mu_{3} = \frac{\sum f(u-\bar{u})^{3}}{N} \times c^{3} = \frac{2.16}{20} \times 20^{3} = 864.$
iv) $\mu_{4} = \frac{\sum f(u-\bar{u})^{4}}{N} \times c^{4} = \frac{35.984}{20} \times 20^{4} = 2,87,872.$

4.4 CHARLIE'S CHECK AND SHEPPARD'S CORRECTIONS

A check which can be used to verify correct computations in a table of grouped classes. For example, consider the following table with specified class limits and frequencies f. The class marks x_i are then computed as well as the rescaled frequencies u_i , which are given by

$$u_i = \frac{f_i - x_0}{c}$$

Where the class mark is taken as $x_0 = 44.5$ and the class interval is c = 10. The remaining quantities are then computed as follows.

Class interval	x_i	f_i	u _i	$f_i u_i$	$f_i u_i^2$	$f_i(u_i + 1)^2$
0-9	4.5	2	-4	-8	32	18
10-19	14.5	3	-3	-9	27	12
20-29	24.5	11	-2	-22	44	11
30-39	34.5	20	-1	-20	20	0
40-49	44.5	32	0	0	0	32
50-59	54.5	25	1	25	25	100
60-69	64.5	7	2	14	28	63
Total		100		-20	176	236

In order to compute the variance, note that

$$V(u) = \frac{\sum f_i u_i^2}{\sum f_i} - \left(\frac{\sum f_i u_i}{\sum f_i}\right)^2$$
$$= \frac{176}{100} - \left(\frac{-20}{100}\right)^2 = 1.72$$

So the variance of the original data is

$$V(x) = c^2 V(u) = 100 \times 1.72 = 172.$$

Charlier's check makes use of the additional column $f_i(u_i + 1)^2$ added to the right side of the table. By noting that the identity

$$\sum f_i (u_i + 1)^2 = \sum f_i (u_i^2 + 2u_i + 1)$$
$$= \sum f_i u_i^2 + 2 \sum u_i + \sum f_i$$

connects columns five through seven, it can be checked that the computations have been done correctly. In the example above,

$$236 = 176 + 2(-20) + 100 \tag{8}$$

Hence, the computations pass Charlier's check.

Charlier's check in computing moments by the coding method uses the following identities:

$$\sum f(u+1) = \sum fu + \sum f$$
$$\sum f(u+1)^2 = \sum fu^2 + 2\sum fu + \sum f$$
$$\sum f(u+1)^3 = \sum fu^3 + 3\sum fu^2 + 3\sum fu + \sum f$$
$$\sum f(u+1)^4$$
$$= \sum fu^4 + 4\sum fu^3 + 6\sum fu^2 + 4\sum fu$$
$$+ \sum f$$

Sheppard's Corrections:

When the frequency distribution, consists of interval, we take x as the class mark of the interval and use this x in all the formulae.

While doing this, it is assumed that all the values in the interval concentrate at the class mark. But this assumption may not be always true and we are likely to get some errors in this calculation.

The well-known statistician Sheppard gave the corrected values of the moments as follows:

$$\mu_1(corrected) = \mu_1$$
$$\mu_2(corrected) = \mu_2 - \frac{c^2}{12}$$
$$\mu_3(corrected) = \mu_3$$
$$\mu_4 = \mu_4 - \frac{1}{2}c^2\mu_2 + \frac{7}{240}c^4$$

Where 'c' is the length of the class-interval, which is the same as the spacing between the mid values.

Note that even though this correction has great mathematical significance, we need not use these corrections in practice because the error is too small hence negligible and also in statistic, we look for estimates, which are approximate values.

Example 9:Apply Sheppard's corrections to determine the moments about the mean for the data

Class Interval	0-10	10-20	20-30	30-40	40-50
Frequency	1	2	9	2	6

		1 1		,	0				
Class	Clas	f	u	fu	<i>(u</i>	f(u	f(u	f(u	f(u
Interv	S		$-\frac{x-a}{a}$		- 0.5)	- 0.5)	$(-0.5)^2$	$(-0.5)^3$	$(-0.5)^4$
al	Mar		— с				-		-
	k								
	(X)								
0-10	5	1	-2	-2	-2.5	-2.5	12.5	-	39.062
								15.62	5
								5	
10-20	15	2	-1	-2	-1.5	-3	4.5	-6.75	10.125
20-30	25	9	0	0	-0.5	-4.5	2.25	-1.125	0.5625
30-40	35	2	1	2	0.5	1	0.5	0.25	0.125
40-50	45	6	2	1	1.5	9	13.5	20.25	30.375
				2					
Total		2		1		0	33.25	-3	80.25
		0		0					

Solution: Lets prepared table, taking A = 25.

Here, $\bar{u} = \frac{\sum fu}{N} = \frac{10}{20} = 0.5.$

The central moments of x are given by

$$\mu_i = \frac{\sum f(u - \bar{u})^i}{N} \times c^i$$

- i) $\mu_1 = \frac{\sum f(u-\bar{u})^1}{N} \times c^1 = \frac{0}{20} \times 20 = 0.$
- ii) $\mu_2 = \frac{\sum f(u-\bar{u})^2}{N} \times c^2 = \frac{33.25}{20} \times 20^2 = 665.$
- iii) $\mu_3 = \frac{\sum f(u-\overline{u})^3}{N} \times c^3 = \frac{-3}{20} \times 20^3 = -1200.$
- iv) $\mu_4 = \frac{\sum f(u-\overline{u})^4}{N} \times c^4 = \frac{80.25}{20} \times 20^4 = 6,42,000.$

Sheppard gave the corrected values of the moments as follows:

 $\mu_{1}(corrected) = \mu_{1} = 0, \qquad \mu_{2}(corrected) = \mu_{2} - \frac{c^{2}}{12} = 665 - \frac{20^{2}}{12} = 631.67$ $\mu_{3}(corrected) = \mu_{3} = -1200$ $\mu_{4} = \mu_{4} - \frac{1}{2}c^{2}\mu_{2} + \frac{7}{240}c^{4} = 6,42,000 - \frac{400}{2} \times 665 + \frac{7}{240} \times 1,60,000 = 5,13,666.67.$

4.5 MOMENTS IN DIMENSIONLESS FORM

To avoid particular units, we can define the dimensionless central moments as

$$a_r = \frac{\mu_i}{\sigma^i}$$

Where σ is the standard deviation, so, as we have $\sigma = \sqrt{\mu_2}$, We already know that for central moments, $\mu_1 = 0$, $\mu_2 = \sigma^2$. So, we get $a_0 = 0$ and $a_1 = 1$.

4.6 SKEWNESS

Skewness is one more concept which deals with the symmetry or rather asymmetry of the values of distribution around its central value. When a frequency distribution is plotted on a chart, an ideal distribution by a nice, symmetric, bell-shaped curve around the central value. Such a distribution is called symmetric distribution or a normal distribution. However in practice every distribution that we across need not be normal. Their graph will be asymmetric or skew. Such distributions are called skewed distribution.

Definition: Skewness defined by famous statistician Garrett "A distribution is said to be skewed when the mean and median fall at different points of the distribution and balance is shifted to one side or the other to left or right."

Types of Skewness: In order to understand this concept we draw the following graphs, where $\bar{x} = \text{Mean}$, $M_e = \text{Median}$ and $M_o = \text{Mode}$ of the distributions.



Figure 4.1

It is clear from the diagram that

- i) Represents a symmetric distribution, for which Mean = Median = Mode.
- ii) Represents a positive skewed distribution for which Mode < Median <Mode.
- iii) Represents a negative skewed distribution for which Mean < Median <Mode.

Measure of Skewness:

Since mean, median and mode are different for a skewed distribution, the simplest measure would be the difference between two of these in pairs. Though such measures are simple to calculate, their main drawback is the following: these measures are expressed with respect to the corresponding units of the distribution. Therefore two distributions with different units cannot be compared. In order to overcome this difficulty, relative measures are defined. These are called Coefficients of Skewness.

Karl Pearson's Coefficient of Skewness: it is defined as

$$S_k = \frac{Mean - Mode}{standard \ deviation} = \frac{\bar{x} - M_o}{\sigma}$$

Using the relation between mean, median and mode:

Mean - Mode = 3 (Mean - Median), we can write

$$S_{k} = \frac{3(Mean - Median)}{Standard \ deviation} = \frac{3(\bar{x} - M_{e})}{\sigma}$$

Interpretation on S_k

- i) If S_k is positive then the distribution is positively skewed.
- ii) If S_k is negative then the distribution is negatively skewed.
- iii) If $S_k = 0$ then the distribution is symmetric.
- iv) Theoretically the limits of S_k are from -3 to +3.

Example 10: For the following ungrouped data find the Karl Pearson's Coefficient of Skewness.

12,18,25,15, 16, 10, 8 15, 27,14

Solution: For the Karl Pearson's Coefficient of Skewness we need to find mean, mode and standard deviation of the data.

Mean =
$$\bar{x} = \frac{\sum x}{n} = \frac{12+18+25+15+16+10+8+15+27+14}{10} = \frac{160}{10} = 16.$$

Mode = 15 (number which repeated maximum time)

$$\sum x^2 = 144 + 324 + 625 + 225 + 256 + 100 + 64 + 225 + 729 + 196 = 2,888$$

$$\sigma = \sqrt{\frac{\sum x^2}{n} - (\bar{x})^2} = \sqrt{\frac{2,888}{10} - (15)^2} = \sqrt{288.8 - 225} = \sqrt{63.8} = 7.99$$

Therefore, the Karl Pearson's Coefficient of Skewness is

$$S_{k} = \frac{Mean - Mode}{standard \ deviation} = \frac{\bar{x} - M_{o}}{\sigma} = \frac{16 - 15}{7.99} = \frac{1}{7.99} = 0.125$$

Example 11: For the following grouped data find the Karl Pearson's Coefficient of Skewness. Also interpret the type of distribution.

C.I	0-4	4-8	8-12	12-16	16-20
F	1	3	10	4	2

Solution: First we find mean, mode and standard deviation.

C.I	F	Х	fx	fx^2
0-4	1	2	2	4
4-8	3	6	18	108
8-12	10	10	100	1000
12-16	4	14	56	784
16-20	2	18	36	648
Total	20		212	2,544

Mean $=\bar{x} = \frac{\sum fx}{N} = \frac{212}{20} = 10.6.$

Standard

$$\sigma = \sqrt{\frac{\sum fx^2}{N} - (\bar{x})^2} = \sqrt{\frac{2,544}{20} - (10.6)^2} = \sqrt{127.2 - 112.36} = \sqrt{14.84}$$

= 3.85.

deviation

=

$$Mode = l_1 + \frac{f_1 - f_0}{2f_1 - f_0 - f_2} \times h$$
$$Mode = 8 + \frac{10 - 3}{2(10) - 3 - 4} \times 4 = 10.15.$$

Therefore, the Karl Pearson's Coefficient of Skewness is

$$S_k = \frac{Mean - Mode}{standard \ deviation} = \frac{\bar{x} - M_o}{\sigma} = \frac{10.6 - 10.15}{3.85} = \frac{0.45}{3.85} = 0.12.$$

Bowley's Coefficient of Skewness:

This measure is based on Quartiles, hence it is also known as Quartile Coefficient of Skewness. It is given by

$$S_B = \frac{Q_3 + Q_1 - 2Q_2}{Q_3 - Q_1}$$

The limits of Bowley's Coefficient of Skewnessare between -1 to +1.

Example 12: Find the Bowley'scoefficient of Skewness for the following information are given: $Q_1 = 12.5$, $Q_2 = 17.2$, $Q_3 = 24.7$

Solution: Given that

$$Q_1 = 12.5, Q_2 = 17.2, Q_3 = 24.7$$

The Bowley's coefficient of Skewness is given by

$$S_B = \frac{Q_3 + Q_1 - 2Q_2}{Q_3 - Q_1} = \frac{24.7 + 12.5 - 2(17.2)}{24.7 - 12.5} = \frac{2.8}{12.2} = 0.23$$

Example 13:Find the Bowley's coefficient of Skewness for the following distribution:

Х	1	3	5	7	9	11
F	3	8	14	20	18	7

Solution: Let find all three quartiles for the distribution:

Х	F	cf(Cummulative frequency)
1	3	3
3	8	11
5	14	25
7	20	45
9	19	64
11	7	71
Total	N =71	

Therefore,

$$Q_{1} = value of \left[1\left(\frac{N+1}{4}\right)\right]^{th} item$$

$$= value of \left[1\left(\frac{71+1}{4}\right)\right]^{th} ite = value of 18th item = 5$$

$$Q_{2} = value of \left[2\left(\frac{N+1}{4}\right)\right]^{th} item$$

$$= value of \left[2\left(\frac{71+1}{4}\right)\right]^{th} ite = value of 36th item = 7$$

$$Q_{3} = value of \left[3\left(\frac{N+1}{4}\right)\right]^{th} item$$

$$= value of \left[3\left(\frac{71+1}{4}\right)\right]^{th} item$$

The Bowley's coefficient of Skewness is given by

$$S_B = \frac{Q_3 + Q_1 - 2Q_2}{Q_3 - Q_1} = \frac{9 + 5 - 2(7)}{9 - 5} = \frac{0}{.4} = 0.$$

Therefore, the distribution is symmetric.

4.7 KURTOSIS

Kurtosis in Greek means 'Bulginess'. In statistics, Kurtosis refers to the degree of flatness or peakedness around the mode of a frequency curve. The measure of kurtosis is with respect to a normal curve, which is accepted as a yardstick to decide the nature of other curves.

In other words measures of kurtosis tell us to what extent the given distribution is flat or peaked with respect to the standard normal curve.



- i) Normal curve is called Mesokurtic (M).
- ii) Flat one is called Platykurtic (P).
- iii) Peaked is called Leptokurtic (L).

Measures of Kurtosis:

The most prominent measure of kurtosis is the coefficient β_2 , given by

$$\beta_2 = \frac{\mu_4}{{\mu_2}^2}$$

Where μ_i 's are the moment about mean \bar{x} .

Bigger value of β_2 gives more peak to the distributions. For normal distribution $\beta_2 = 3$.

Hence the given distribution is :

- i) Leptokurtic, if $\beta_2 > 3$.
- ii) Mesokurtic, if $\beta_2 = 3$.
- iii) Platykurtic, if $\beta_2 < 3$.

Example 14: for the following distribution find β_1 and β_2 and comment on the Skewness and Kurtosis of the distribution.

Х	2	3	4	5
f	4	3	2	1

Solution: First calculate moments about mean for the given distribution.

	$\bar{x} = \frac{\sum f}{\sum f}$	$\frac{x}{x} = \frac{2(4) + 1}{2}$	(-3(3) + 4) (+3 + 2)	(2) + 5(1) + 1	$=\frac{30}{10}=3.$	
x	f	(<i>x</i> – 3)	f(x - 3)	$f(x - 3)^2$	$f(x - 3)^3$	$f(x - 3)^4$
2	4	-1	-4	4	-4	4
3	3	0	0	0	0	0
4	2	1	2	2	2	2
5	1	2	2	4	8	16
Total	10		0	10	6	22

The central moments are given by

The first moment about \bar{x} , as i)

$$\mu_1 = \frac{\sum f(x-\bar{x})}{N} = \frac{0}{10} = 0 \; .$$

ii) The second moment about \bar{x} , as

$$\mu_2 = \frac{\sum f(x - \bar{x})^2}{N} = \frac{10}{10} = 1$$

iii) The third moment about \bar{x} , as

$$\mu_3 = \frac{\sum f(x-\bar{x})^3}{N} = \frac{6}{10} = 0.6$$

iv) The fourth moments about \bar{x} , as

$$\mu_4 = \frac{\sum f(x-\bar{x})^4}{N} = \frac{22}{10} = 2.2.$$

$$\beta_1 = \frac{\mu_3^2}{\mu_2^3} = \frac{(0.6)^2}{(1)^3} = 0.36.$$

$$\beta_2 = \frac{\mu_4}{\mu_2^2} = \frac{2.2}{(1)^2} = 2.2$$

Since $\beta_1 \neq 0$, the curve is not symmetric Also $\mu_3 = 0.6 > 0$. Therefore, the curve is positively skewed.

Since $\beta_2 = 2.2 < 3$.

Therefore, the curve is flat as compared to the normal curve. Hence the distribution is platykurtic.

4.8 SOFTWARE COMPUTATION OF SKEWNESS AND **KURTOSIS**

To compute Skewness and Kurtosis by using different software are given below:

Sigma Magic: Using the Sigma Magic software, calculating the Skewness and Kurtosis is relatively straightforward. Just add a new Basic Statistics template to Excel by clicking on Stat > Basic Statistics. Copy and paste the data for which you want to Skewness and Kurtosis into the input area and then click on Compute Outputs. The analysis results will include the Skewness and Kurtosis values.

Excel: You could also calculate these values in Excel by using the formula =SKEW(...) for the Skewness value, =KURT(...) for the Kurtosis value.

Minitab: If you use the Minitab software, you can copy and paste the data into Minitab and then click on Stat > Basic Statistics > Display Descriptive Statistics. Then select the data column and then click on OK. This will print out the quartiles for the sample values. If you want the Skewness and Kurtosis values, you have to go back to the menu and click on Statistics and select the checkbox next to Skewness and Kurtosis in the statistics options. Note that the values provided by Minitab may be slightly different from Excel and Sigma Magic software.

4.10 SUMMARY

In this unit, we have discussed:

- Moments and its types for ungroup and grouped data.
- The relation between raw, arbitrary and central moments.
- The effect of change of origin and scale on moments.
- Charlie's check, and Shephard's Correction for Moments.
- Skewness and about symmetry of distribution.
- Kurtosis.

4.11 EXERCISE

1. The first four moments of a distribution are 1, 4, 10 and 46 respectively. Compute the moment coefficients of skewness and kurtosis and comment upon the nature of the distribution.

2. Compute the first four central moments from the following data. Also find the two beta coefficients.

Х	5	10	15	20	25	30	35
f	8	15	20	32	23	17	5

3. The first four central moments of a distribution are 0, 2.5, 0.7 and 18.75. Examine the skewness and kurtosis of the distribution.

		4.0			16.20
Class	0-4	4-8	8-12	12-16	16-20
interval					
Frequency	5	8	13	9	5

4. Calculate first four central moments for the following distribution:

5. Find the first four arbitrary moments about A = 7 for the following:

10, 5, 8, 7, 2, 3, 12, 14

6. Find raw moment for the following data:

C.I	5-10	10-15	15-20	20-25	25-30
f	3	4	7	4	2

7. The first four central moments of a distribution are 0, 15, 36, 78. If the mean of the distribution is 8, find the moments about A = 5.

8. The first four raw moments about origin are 4, 16, 33, 89. Find mean and the first four central moments.

9. For the following data verify Charlie's check :

C.I	2-8	8-14	14-20	20-26
f	1	4	3	2

10. Find the first four central moments using coding method, also find Sheppard's correction for moments.

C.I	0-5	5-10	10-15	15-20	20-25	25-30
f	3	8	12	13	7	2

11. For the following data, find Karl Pearson's coefficient of Skewness and also find the type of distribution:

i) 12,15,17, 12,8,25,16,6,7,41

ii) 3,7,8, 12, 15

Х	1	2	3	4	5	6	7
f	2	8	12	15	18	9	6

C.I	0-2	2-4	4-6	6-8	8-10
frequency	4	7	13	10	6

12. Find the Bowely's coefficient of Skewness for each of the following:

i) $Q_1 = 165.5$, $Q_2 = 184.3$, $Q_3 = 196.7$.

ii) 2,8,7,12,14,17,20.

Х	1	2	3	4	5	6	7
F	2	8	12	15	18	9	6

4.12 LIST OF REFERENCES

- Statistics by Murry R. Spiegel, Larry J. Stephens. Publication McGRAWHILL INTERNATIONAL.
- Fundamental Statistic by S.C. Gupta

ELEMENTARY PROBABILITY THEORY

Unit Structure

- 5.1 Objective
- 5.2 Introduction
- 5.3 Definitions of Probability
- 5.4 Conditional Probability

5.4.1 Independent and Dependent Events, Mutually Exclusive Events

- 5.5 Probability Distributions
- 5.6 Mathematical Expectation
- 5.7 Combinatorial Analysis
- 5.8 Combinations, Stirling's Approximation to n!
- 5.9 Relation of Probability to Point Set Theory, Euler or Venn Diagrams and Probability
- 5.10 Summary
- 5.11 Exercise
- 5.12 List of References

5.1 OBJECTIVE

After going through this unit, you will able to :

- Determine the probability of different experimental results.
- Explain the concept of probability.
- Calculate probability for simple, compound and complimentary events.
- Conditional probability and its examples.
- Independent events and multiplication theorem of probability.
- Probability distribution and its Expected value of probability distribution.
- Combination and Stirling's number approximation.
- Relations between probability and set theory with help of Venn diagram.

5.2 INTRODUCTION

Some times in daily life certain things come to mind like "I will be success today', I will complete this work in hour, I will be selected for job and so

on. There are many possible results for these things but we are happy when we get required result. Probability theory deals with experiments whose outcome is not predictable with certainty. Probability is very useful concept. These days many field in computer science such as machine learning, computational linguistics, cryptography, computer vision, robotics other also like science, engineering, medicine and management.

Probability is mathematical calculation to calculate the chance of occurrence of some happening, we need some basic concept on random experiment, sample space, and events.

Basic concept of probability:

Random experiment: When experiment can be repeated any number of times under the similar conditions but we get different results on same experiment, also result is not predictable such experiment is called random experiment. For e.g. A coin is tossed, A die is rolled and so on.

Outcomes: The result which we get from random experiment is called outcomes of random experiment.

Sample space: The set of all possible outcomes of random experiment is called sample space. The set of sample space is denoted by S and number of elements of sample space can be written as n(S). For e.g. A die is rolled, we get = {1,2,3,4,5,6}, n(S) = 6.

Events: Any subset of the sample space is called an event. Or a set of sample point which satisfies the required condition is called an events. Number of elements in event set is denoted by n(E). For example in the experiment of throwing of a dia. The sample space is

 $S = \{1, 2, 3, 4, 5, 6\}$ each of the following can be an event :

i) A: even number i.e. $A = \{2, 4, 6\}$ ii) B: multiple of 3 i.e. $B = \{3, 6\}$ iii) C: prime numbers i.e. $C = \{2, 3, 5\}$.

Types of events:

Impossible event: An event which does not occurred in random experiment is called impossible event. It is denoted by \emptyset set. i. e. $n(\emptyset) = 0$. For example getting number 7 when die is rolled. The probability measure assigned to impossible event is Zero.

Equally likely events: when all events get equal chance of occurrences is called equally likely events. For e.g. Events of occurrence of head or tail in tossing a coin are equally likely events.

Certain event: An event which contains all sample space elements is called certain events. i.e.n(A) = n(S).

Mutually exclusive events: Two events A and B of sample space S, it does not have any common elements are called mutually exclusive events. In the experiment of throwing of a die A: number less than 2, B: multiple of 3. There fore $n(A \cap B) = 0$

Exhaustive events: Two events A and B of sample space S, elements of event A and B occurred together are called exhaustive events. For e.g. In a

thrown of fair die occurrence of even number and occurrence of odd number are exhaustive events. There fore $n(A \cup B) = 1$.

Complement event: Let S be sample space and A be any event than complement of A is denoted by \overline{A} is set of elements from sample space S, which does not belong to A. For e.g. if a die is thrown, S = {1, 2, 3, 4, 5, 6} and A: odd numbers, A = {1, 3, 5}, then \overline{A} = {2,4,6}.

5.3 DEFINITIONS OF PROBABILITY

Probability: For any random experiment, sample space S with required chance of happing event E than the probability of event E is define as

$$P(E) = \frac{n(E)}{n(S)}$$

Basic properties of probability:

- 1) The probability of an event E lies between 0 and 1. i.e. $0 \le P(E) \le 1$.
- 2) The probability of impossible event is zero. i.e. $P(\phi) = 0$.
- 3) The probability of certain event is unity. i.e. P(E) = 1.
- 4) If A and B are exhaustive events than probability of $P(A \cup B) = 1$.
- 5) If A and B are mutually exclusive events than probability of $P(A \cap B) = 0$.
- 6) If A be any event of sample space than probability of complement of A is given by $P(A) + P(\overline{A}) = 1 \Rightarrow : P(\overline{A}) = 1 P(A)$.

Probability Axioms:

Let S be a sample space. A probability function P from the set of all event in S to the set of real numbers satisfies the following three axioms for all events A and B in S.

- i) $P(A) \ge 0$.
- ii) $P(\emptyset) = 0$ and P(S) = 1.
- iii) If A and B are two disjoint sets i.e. $A \cap B = \emptyset$ than the probability of the union of A and B is $P(A \cup B) = P(A) + P(B)$.

Theorem: Prove that for every event A of sample space S, $0 \le P(A) \le 1$.

Proof:
$$S = A \cup \overline{A}$$
, $\emptyset = A \cap \overline{A}$.
 $\therefore 1 = P(S) = P(A \cup \overline{A}) = P(A) + P(\overline{A})$
 $\therefore 1 = P(A) + P(\overline{A})$

 $\Rightarrow P(A) = 1 - P(\overline{A}) \text{ or } P(\overline{A}) = 1 - P(A).$ If $P(A) \ge 0$. than $P(\overline{A}) \le 1$.

: for every event *A*; $0 \le P(A) \le 1$.

Addition theorem of probability:

Theorem: If A and B are two events of sample space S, then probability of union of A and B is given by $P(A \cup B) = P(A) + P(B) - P(A \cap B)$.

Proof: A and B are two events of sample space S.



Now from diagram probability of union of two events A and B is given by,

 $P(A \cup B) = P(A \cap \overline{B}) + P(A \cap B) + P(B \cap \overline{A})$ But $P(A \cap \overline{B}) = P(A) - P(A \cap B)$ and $P(B \cap \overline{A}) = P(B) - P(A \cap B)$. $\therefore P(A \cup B) = P(A) - P(A \cap B) + P(A \cap B) + P(B) - P(A \cap B)$ $\therefore P(A \cup B) = P(A) + P(B) - P(A \cap B).$

Note: The above theorem can be extended to three events A, B and C as shown below:

$$P(A \cup B \cup C) = P(A) + P(B) + P(C) - P(A \cap B) - P(B \cap C)$$
$$- P(C \cap A) + P(A \cap B \cap C)$$

Example 1: A bag contains 4 black and 6 white balls; two balls are selected at random. Find the probability that balls are i) both are different colors. ii) both are of same colors.

Solution: Total number of balls in bag = 4 blacks + 6 white = 10 balls To select two balls at random, we get

$$n(S) = C(10,2) = 45.$$

i) A be the event to select both are different colors.

$$\therefore n(A) = C(4,1) \times C(6,1) = 4 \times 6 = 24.$$
$$P(A) = \frac{n(A)}{n(S)} = \frac{24}{45} = 0.53.$$

ii) To select both are same colors.

Let Abe the event to select both are black balls

$$n(A) = C(4,2) = 6$$
$$P(A) = \frac{n(A)}{n(S)} = \frac{6}{45}$$

Let B be the event to select both are white balls.

$$n(B) = C(6,2) = 15$$

$$P(B) = \frac{n(B)}{n(S)} = \frac{15}{45}$$
.

A and B are disjoint event.

∴ The required probability is

 $P(A \cup B) = P(A) + P(B) = \frac{6}{45} + \frac{15}{45} = \frac{21}{45} = 0.467.$

Example 2: From 40 tickets marked from 1 to 40, one ticket is drawn at random. Find the probability that it is marked with a multiple of 3 or 4.

Solution: From 40 tickets marked with 1 to 40, one ticket is drawn at random

$$n(S) = C(40,1) = 40$$

it is marked with a multiple of 3 or 4, we need to select in two parts.

Let A be the event to select multiple of 3,

i.e. $A = \{3, 6, 9, \dots, 39\}$

$$n(A) = C(13,1) = 13$$
$$P(A) = \frac{n(A)}{n(S)} = \frac{13}{40}$$

Let B be the event to select multiple of 4.

i.e. $B = \{4, 8, 12, \dots, 40\}$

$$n(B) = C(10,1) = 10$$
$$P(B) = \frac{n(B)}{n(s)} = \frac{10}{40}.$$

Here A and B are not disjoint.

 $A \cap B$ be the event to select multiple of 3 and 4.

i.e. $A \cap B = \{12, 24, 36\}$

$$n(A \cap B) = C(3,1) = 3$$

 $P(A \cap B) = \frac{n(A \cap B)}{n(S)} = \frac{3}{40}$

 \therefore The required probability is

$$P(A \cup B) = P(A) + P(B) - P(A \cap B) = \frac{13}{40} + \frac{10}{40} - \frac{3}{40} = \frac{20}{40} = 0.5.$$

Example 3: If the probability is 0.45 that a program development job; 0.8 that a networking job applicant has a graduate degree and 0.35 that applied for both. Find the probability that applied for atleast one of jobs. If number of graduate are 500 then how many are not applied for jobs?

Solution: Let Probability of program development job= P(A) = 0.45.

Probability of networking job= P(B) = 0.8.

Probability of both jobs = $P(A \cap B) = 0.35$.

Probability of atleast one i.e. to find $P(A \cup B)$.

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$
$$P(A \cup B) = 0.45 + 0.8 - 0.35 = 0.9$$

Now there are 500 application, first to find probability that not applied for job.

$$P(\overline{A \cup B}) = 1 - P(A \cup B) = 1 - 0.9 = 0.1$$

Number of graduate not applied for job = $0.1 \times 500 = 50$.

Check your Progress:

- 1. A card is drawn from pack of 52 cards at random. Find the probability that it is a face card or a diamond card.
- 2. If $P(A) = \frac{3}{8}$ and $(B) = \frac{5}{8}$, $P(A \cup B) = \frac{7}{8}$ than find i) $P(\overline{A \cup B})$ ii) $P(A \cap B)$.
- 3. In a class of 60 students, 50 passed in computers, 40 passed in mathematics and 35 passed in both. What is the probability that a student selected at random has i) Passed in atleast one subject, ii) failed in both the subjects, iii) passed in only one subject.

5.4 CONDITIONAL PROBABILITY

In many case we come across occurrence of an event A and for the same are required to find out the probability of occurrence an event B which depend on event A. This kind of problem is called conditional probability problems.

Definition: Let A and B be two events. The conditional probability of event B, if an event A has occurred is defined by the relation,

$$P(B|A) = \frac{P(B \cap A)}{P(A)}$$
 if and only if $P(A) > 0$.

In case when P(A) = 0, P(B|A) is not define because $P(B \cap A) = 0$ and $P(B|A) = \frac{0}{0}$ which is an indeterminate quantity.

Similarly, Let A and B be two events. The conditional probability of event A, if an event B has occurred is defined by the relation,

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$
 If and only if $P(B) > 0$.

Example 4: A pair of fair dice is rolled. What is the probability that the sum of upper most face is 6, given that both of the numbers are odd?

Solution: A pair of fair dice is rolled, therefore n(S) = 36.

A to select both are odd number, i.e. $A = \{(1,1), (1,3), (1,5), (3,1), (3,3), (3,5), (5,1), (5,3), (5,5)\}.$

$$P(A) = \frac{n(A)}{n(S)} = \frac{9}{36}$$
90

B is event that the sum is 6, i.e. $B = \{ ((1,5),(2,4), (3,3),(4,2), (5,1) \}.$

$$P(B) = \frac{n(B)}{n(S)} = \frac{5}{36}$$

 $A \cap B = \{ (1,5), (3,3), (5,1) \}$

$$P(A \cap B) = \frac{n(A \cap B)}{n(S)} = \frac{3}{36}$$

By the definition of conditional probability,

$$P(B|A) = \frac{P(A \cap B)}{P(A)} = \frac{3/_{36}}{9/_{36}} = \frac{1}{3}.$$

Example 5: If A and B are two events of sample space S, such that P(A) = 0.85, P(B) = 0.7 and $P(A \cup B) = 0.95$. Find i) $P(A \cap B)$, ii) P(A|B), iii) P(B|A).

Solution: Given that P(A) = 0.85, P(B) = 0.7 and $P(A \cup B) = 0.95$.

i) By Addition theorem,

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

0.95 = 0.85 + 0.7 - P(A \cap B)
$$P(A \cap B) = 1.55 - 0.95 = 0.6.$$

ii) By the definition of conditional probability,

$$P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{0.6}{0.7} = 0.857.$$

iii)
$$P(B|A) = \frac{P(A \cap B)}{P(A)} = \frac{0.6}{0.85} = 0.706$$

Example 6: An urn A contains 4 Red and 5 Green balls. Another urn B contains 5 Red and 6 Green balls. A ball is transferred from the urn A to the urn B, then a ball is drawn from urn B. find the probability that it is Red.

Solution: Here there are two cases of transferring a ball from urn A to B.

Case I: When Red ball is transferred from urn A to B.

There for probability of Red ball from urn A is $P(R_A) = \frac{4}{2}$

After transfer of red ball, urn B contains 6 Red and 6 Green balls.

Now probability of red ball from urn B = $P(R_B|R_A) \times P(R_A) = \frac{6}{12} \times \frac{4}{9} = \frac{24}{108}$.

Case II: When Green ball is transferred from urn A to B.

There for probability of Green ball from urn A is $P(G_A) = \frac{5}{9}$

After transfer of red ball, urn B contains 5 Red and 7 Green balls.

Now probability of red ball from urn B = $P(R_B|G_A) \times P(G_A) = \frac{5}{12} \times \frac{5}{9} = \frac{25}{12}$

108

Therefore required probability $=\frac{24}{108} + \frac{25}{108} = \frac{49}{108} = 0.4537.$

Check your progress:

- 1. A family has two children. What is the probability that both are boys, given at least one is boy?
- 2. Two dice are rolled. What is the condition probability that the sum of the numbers on the dice exceeds 8, given that the first shows 4?
- 3. Consider a medical test that screens for a COVID-19 in 10 people in 1000. Suppose that the false positive rate is 4% and the false negative rate is 1%. Then 99% of the time a person who has the condition tests positive for it, and 96% of the time a person who does not have the condition tests negative for it. a) What is the probability that a randomly chosen person who tests positive for the COVID-19 actually has the disease? b) What is the probability that a randomly chosen person who tests negative for the COVID-19 does not indeed have the disease?

5.4.1 Independent and Dependent Events, Mutually Exclusive Events: Independent events:

Two events are said to be independent if the occurrence of one of them does not affect and is not affected by the occurrence or non-occurrence of other.

i.e.
$$P(B/A) = P(B)$$
 or $P(A/B) = P(A)$.

Multiplication theorem of probability: If A and B are any two events associated with an experiment, then the probability of simultaneous occurrence of events A and B is given by

$$P(A \cap B) = P(A)P(B/A)$$

Where P(B/A) denotes the conditional probability of event B given that event A has already occurred.

OR

$$P(A \cap B) = P(B)P(A/B)$$

Where P(A/B) denotes the conditional probability of event A given that event B has already occurred.

11.5.1 For Independent events multiplication theorem:

If A and B are independent events then multiplication theorem can be written as,

$$P(A \cap B) = P(A)P(B)$$

Proof. Multiplication theorem can be given by,

If A and B are any two events associated with an experiment, then the probability of simultaneous occurrence of events A and B is given by

$$P(A \cap B) = P(A)P(B/A)$$

By definition of independent events, P(B/A) = P(B) or P(A/B) = P(A).

 $\therefore P(A \cap B) = P(A)P(B).$

Note:

1) If A and B are independent event then, \overline{A} and \overline{B} are independent event.

2) If A and B are independent event then, \overline{A} and B are independent event.

3) If A and B are independent event then, A and \overline{B} are independent event.

Example 7: Manish and Mandar are trying to make Software for company. Probability that Manish can be success is $\frac{1}{5}$ and Mandar can be success is $\frac{3}{5}$, both are doing independently. Find the probability that i) both are success. ii) Atleast one will get success. iii) None of them will success. iv) Only Mandar will success but Manish will not success.

Solution: Let probability that Manish will success is $P(A) = \frac{1}{5} = 0.2$. Therefore probability that Manish will not success is $P(\bar{A}) = 1 - P(A) = 1 - 0.2 = 0.8$.

Probability that Mandar will success is $P(B) = \frac{3}{5} = 0.6$.

Therefore probability that Mandar will not success is $P(\overline{B}) = 1 - P(B) = 1 - 0.6 = 0.4$.

- Both are success i.e. P(A ∩ B).
 P(A ∩ B) = P(A) × P(B) = 0.2 × 0.6 = 0.12 ∵A and B are independent events.
- ii) Atleast one will get success. i.e. $P(A \cup B)$ By addition theorem, $P(A \cup B) = P(A) + P(B) - P(A \cap B) = 0.2 + 0.6 - 0.12 = 0.68.$
- iii) None of them will success $P(\overline{A \cup B})$ or $P(\overline{A \cap B})$ [ByDeMorgan's law both are same] $P(\overline{A \cup B}) = 1 - P(A \cup B) = 1 - 0.68 = 0.32.$ Or

If A and B are independent than \overline{A} and \overline{B} are also independent.

 $P(\bar{A} \cap \bar{B}) = P(\bar{A}) \times P(\bar{B}) = 0.8 \times 0.4 = 0.32.$

iv) Only Mandar will success but Manish will not success. i.e. $P(\overline{A} \cap B)$.

$$P(\bar{A} \cap B) = P(\bar{A}) \times P(B) = 0.8 \times 0.6 = 0.48$$

Example 8: 50 coding done by two students A and B, both are trying independently. Number of correct coding by student A is 35 and student B is 40. Find the probability of only one of them will do correct coding.

Solution: Let probability of student A get correct coding is $P(A) = \frac{35}{50} = 0.7$

Probability of student A get wrong coding is $P(\bar{A}) = 1 - 0.7 = 0.3$

Probability of student B get correct coding is $P(B) = \frac{40}{50} = 0.8$

Probability of student B get wrong coding is $P(\overline{B}) = 1 - 0.8 = 0.2$.

The probability of only one of them will do correct coding.

i.e. A will correct than B will not or B will correct than A will not.

$$P(A \cap \bar{B}) + P(B \cap \bar{A}) = P(A) \times P(\bar{B}) + P(B) \times P(\bar{A}).$$

= 0.7 × 0.2 + 0.8 × 0.3 = 0.14 + 0.24
= 0.38

Example 9: Given that $P(A) = \frac{3}{7}$, $P(B) = \frac{2}{7}$, if A and B are independent events than find i) $P(A \cap B)$, ii) $P(\overline{B})$, iii) $P(A \cup B)$, iv) $P(\overline{A} \cap \overline{B})$. **Solution:** Given that $P(A) = \frac{3}{7}$, $P(B) = \frac{2}{7}$.

i) A and B are independent events,

$$\therefore P(A \cap B) = P(A) \times P(B) = \frac{3}{7} \times \frac{2}{7} = \frac{6}{49} = 0.122$$

ii)
$$P(\overline{B}) = 1 - P(B) = 1 - \frac{2}{7} = \frac{5}{7} = 0.714$$

iii) By addition theorem,

$$P(A \cup B) = P(A) + P(B) - P(A \cap B) = \frac{3}{7} + \frac{2}{7} - \frac{6}{49} = \frac{29}{49} = 0.592.$$

iv) $P(\bar{A} \cap \bar{B}) = P(\bar{A} \cup \bar{B}) = 1 - P(A \cup B) = 1 - 0.592 = 0.408.$

Check your progress:

- 1. If $P(A) = \frac{2}{5}$, $P(B) = \frac{1}{3}$ and if A and B are independent events, find $(i)P(A \cap B)$, $(ii)P(A \cup B)$, $(iii)P(\overline{A} \cap \overline{B})$.
- 2. The probability that A , B and C can solve the same problem independently are $\frac{1}{3}, \frac{2}{5}$ and $\frac{3}{4}$ respectively. Find the probability that i) the problem remain unsolved, ii) the problem is solved, iii) only one of them solve the problem.
- 3. The probability that Ram can shoot a target is $\frac{2}{5}$ and probability of Laxman can shoot at the same target is $\frac{4}{5}$. A and B shot independently. Find the probability that (i) the target is not shot at all, (ii) the target is shot by at least one of them. (iii) the target shot by only one of them. iv) target shot by both.

5.5 PROBABILITY DISTRIBUTIONS

In order to understand the behavior of a random variable, we may want to look at its average value. For probability we need to find Average is called expected value of random variable X. for that first we have to learn some basic concept of random variable.

Random Variable: A probability measurable real valued functions, say X, defined over the sample space of a random experiment with respective probability is called a random variable.

Types of random variables: There are two type of random variable.

Discrete Random Variable: A random variable is said to be discrete random variable if it takes finite or countably infinite number of values. Thus discrete random variable takes only isolated values.

Continuous Random variable: A random variable is continuous if its set of possible values consists of an entire interval on the number line.

Probability Distribution of a random variable: All possible values of the random variable, along with its corresponding probabilities, so that $\sum_{n=1}^{\square} \square_n = I$, is called a probability distribution of a random variable.

The probability function always follows the following properties:

i) $P(x_i) \ge 0$ for all value of *i*.

ii) $\sum_{i=1}^{n} P_i = 1$.

The set of values x_i with their probability P_i constitute a discrete probability distribution of the discrete variable X.

For e.g. Three coins are tossed, the probability distribution of the discrete variable X is getting head.

$X = x_i$	0	1	2	3
$P(x_i)$	1	3	3	1
	8	8	8	8

5.6 MATHEMATICAL EXPECTATION

All the probability information of a random variable is contained in probability mass function for random variable, it is often useful to consider various numerical characteristics of that random variable. One such number is the expectation of a random variable.

If random variable X takes values x_1, x_2, \dots, x_n with corresponding probabilities P_1, P_2, \dots, P_n respectively, then expectation of random variable X is

 $E(X) = \sum_{i=1}^{n} p_i x_i$ where $\sum_{i=1}^{n} P_i = 1$

Example 10: In Vijay sales every day sale of number of laptops with his past experience the probability per day are given below:

No. of	0	1	2	3	4	5
laptop						
Probability	0.05	0.15	0.25	0.2	0.15	0.2

Find his expected number of laptops can be sale?

Solution: Let X be the random variable that denote number of laptop sale per day.

To calculate expected value, $E(X) = \sum_{i=1}^{n} p_i x_i$

 $E(X) = (0 \times 0.05) + (1 \times 0.15) + (2 \times 0.25) + (3 \times 0.2) + (4 \times 0.15) + (5 \times 0.2)$

$$E(X) = 2.85 \sim 3$$

Therefore expected number of laptops sale per day is 3.

Example 11: A random variable X has probability mass function as follow:

$X = x_i$	-1	0	1	2	3
$P(x_i)$	K	0.2	0.3	2k	2k

Find the value of k, and expected value.

Solution: A random variable X has probability mass function,

 $\sum_{i=1}^{n} P_i = 1.$ $\Rightarrow k + 0.2 + 0.3 + 2k + 2k = 1$ $\Rightarrow 5k = 0.5$ $\Rightarrow k = 0.1$

Therefore the probability distribution of random variable X is

$X = x_i$	-1	0	1	2	3
$P(x_i)$	0.1	0.2	0.3	0.2	0.2

To calculate expected value, $E(X) = \sum_{i=1}^{n} p_i x_i$

 $E(X) = (-1 \times 0.1) + (0 \times 0.2) + (1 \times 0.3) + (2 \times 0.2) + (3 \times 0.2) = 1.2$

Example 12: A box contains 5 white and 7 black balls. A person draws 3 balls at random. He gets Rs. 50 for every white ball and losses Rs. 10 every black ball. Find the expectation of him.

Solution: Total number of balls in box = 5 white + 7 black = 12 balls.

To select 3 balls at random, $n(s) = C(12,3) = \frac{12 \times 11 \times 10}{3 \times 2 \times 1} = 220.$

Let A be the event getting white ball.

A takes value of 0, 1, 2 and 3 white ball.

Case I : no white ball. i.e. A = 0,

$$P(A=0) = \frac{C(7,3)}{220} = \frac{35}{220}$$

Case II: one white ball i.e. A = 1,

$$P(A = 1) = \frac{C(5,1) \times C(7,2)}{220} = \frac{105}{220}$$

Case III: two white balls i.e. A = 2,

$$P(A = 2) = \frac{C(5,2) \times C(7,1)}{220} = \frac{70}{220}$$

Case IV: three white balls i.e. A = 3,

$$P(A=3) = \frac{C(5,3)}{220} = \frac{10}{220}$$

Now let X be amount he get from the game.

Therefore the probability distribution of X is as follows:

$X = x_i$	-30	30	90	150
$P(x_i)$	35	105	70	10
	$\overline{220}$	220	220	220

To calculate expected value, $E(X) = \sum_{i=1}^{n} p_i x_i$ $E(X) = \left(-30 \times \frac{35}{220}\right) + \left(30 \times \frac{105}{220}\right) + \left(90 \times \frac{70}{220}\right) + \left(150 \times \frac{10}{220}\right) = \text{Rs.}$ 45.

5.7 COMBINATORIAL ANALYSIS

Multiplication Rule:

If the procedure can be broken into first and second stages, and if there are m possible outcomes for the first stage and for each of these outcomes, there are n possible outcomes for second stage, then the total procedure can be carried out in the designate order, in $m \times n$ ways.

This principle can be extended to a general form as follows:

Theorem : If a process consists of *n* steps, and

- i) The first step can be performed by n_1 ways.
- ii) The second step can be performed by n_2 ways.
- iii) The i^{th} step can be performed by n_i ways.

Then the whole process can be completed by $n_1 \times n_2 \times ... \times n_i$ different ways.

Example 13: There are 8 men and 7 women in a drama company. How many way the director has to choose a couple to play lead roles in a stage show?

Solution: The director can choose a man (task 1) in 8 ways and then a woman (task 2) in 7 ways. Then by multiplication rule he can choose a couple from $8 \times 7 = 56$ ways.

Example 14: How many four digits numbers can be formed contains each of the digits 7, 8, and 9 exactly once?

Solution: To construct four digits number we have four places.

Thousand place Hundred place Ten place Unit place

First for '7' there are 4 places, for '8' there are 3 places and for '9' there are 2 places. For last digit, we can choose any of 0,1,2,3,4,5,6 so there will be 7 digits.

Thus these can be done by $4 \times 3 \times 2 \times 7 = 168$ ways.

Example 15: To generate typical personal identification number (PIN) is a sequence of any four symbols chosen from the letters in the alphabet and the digits, How many different PIN's are generated?

- i) repetition is not allowed.
- ii) repetition is allowed.

Solution: There are 26 letters of alphabets and 10 digits. Total different symbols are 36.

i) Repetition is not allowed:

There are four place to generate PIN with four symbols,

First place can be filled by 36 ways, second place can be filled by 35 ways, third place can be filled by 34 ways and last fourth place can be filled by 33 ways.

By the multiplication rule,

Therefore these can be done by $36 \times 35 \times 34 \times 33 = 1,413,720$ ways.

ii) Repetition is allowed:

Since repetition is allowed, so each place can be filled by 36 ways, By multiplication rule,

These can be done by $36 \times 36 \times 36 \times 36 = 1,679,616$ ways.

Check your progress:

- A license plate can be made by 2 letters followed by 3 digits. How many different license plates can be made if i) repetition is not allowed.
 Repetition is allowed.
- 2. Mr. Modi buying a personal computer system is offered a choice of 4 models of basic units, 2 models keyboard, and 3 models of printer. How many distinct systems can be purchased?

Counting elements of disjoint sets with Addition Rule:

In above section we have discussed counting problem that can be solved using possibility tree. Here we discuss counting problem that can be solved using the operation sets like union , intersection and the difference between two sets.

The addition rule:

If a task can be performed in m ways and another task in n ways assuming that these two tasks cannot perform simultaneously, then the performing either task can be accomplished in any one of the m + n ways.

In general from as follows:

If there $\operatorname{are} n_1, n_2, n_3, \dots, n_m$ different objects in m different sets respectively and the sets are disjoint, then the number of ways to select an object from one of the m sets is $n_1 + n_2 + n_3 + \dots + n_m$

Example 16: How many different number of signals that can be sent by 5 flags of different colours taking one or more at a time ?

Solution: Let number of signal made by one colour flag = 5 ways.

Number of signal made by two colours flag = $5 \times 4 = 20$ ways.

Number of signal made by three flag colours = $5 \times 4 \times 3 = 60$ ways.

Number of signal made by four flag colours = $5 \times 4 \times 3 \times 2 = 120$ ways.

Number of signal made by five flag colours= $5 \times 4 \times 3 \times 2 \times 1 = 120$ ways.

Using Addition rule we get,

Therefore total number of signals = 5 + 20 + 60 + 120 + 120 = 325 ways.

Example 17: There are 4 different English books, 5 different Hindi books and 7 different Marathi books. How many ways are there to pick up an pair of two books not both with the same subjects?

Solution: One English and one Hindi book is chosen, that selection can be done by $= 4 \times 5 = 20$ ways.

One English and one Marathi book is chosen, that selection can be done by $= 4 \times 7 = 28$ ways.

One Hindi and one Marathi book is chosen, that selection can be done by $= 5 \times 7 = 35$ ways.

These three types of selection are disjoint, therefore by addition rule,

Total selection can be done by = 20 + 28 + 35 = 83 ways.

Additive Principle with Disjoint sets:

Given two sets A and B, both sets are disjoint i.e. if $A \cap B = \emptyset$, than $|A \cup B| = |A| + |B|$.

Example 18: In college 200 students visit to canteen every day of which 80 likes coffee and 70 likes tea. If no one student like both than find i) number of students like atleast one of them? ii) number of students like none of them?

Solution: Total number of students = 200

Total number of students who like coffee = |A| = 80.

Total number of students who like tea = |B| = 70.

Total number of students like atleast one $|A \cup B| = |A| + |B| = 80 + 70 = 150$.

Total number of students like none of them = 200 - 150 = 50.

Definition: An r-combination of n distinct objects is an unordered selection, or subset, of r out of the n objects. We use $C(n,r)or_r^n C$ to denote the number of r-combinations. This number is called as binomial number.

If $x_1, x_2, x_3, \dots, x_n$ are n distinct objects, and r is any integer, with $1 \le r \le n$. Therefore selecting r-objects from n objects is given by

$$C(n,r) = \frac{n!}{r! (n-r)!}$$

Example 19: How many elements of set 3-bit string with weight 2?

Solution: there are 3-bit with weight 2, i.e. n = 3, r = 2.

These can be done by = C(n, r) = C(3, 2) = 3.

Therefore the bit string is 011, 101, 110.

Example 20: A bag contains 4 red marbles and 5 green marbles. Find the number of ways that 4 marbles can be selected from the bag, if selection contain i) No restriction of colors. ii) all are of same colors.

Solution: Total number of marbles: 4 Red + 5 Green = 9 marbles

To select 4 marbles from the bag with condition,

i) No restriction of colors:

These can be done by :C(9,4) = 126 ways.

ii) All are of same colors:

First select the colors by : C(2, 1) = 2.

If all is Red in colors than these can be done by = C(4, 4) = 1.

If all is Green in colors then these can be done by = C(5, 4) = 5.

Therefore total number of ways = $2 \times 1 \times 5 = 10$ ways.

Example 21: There are 10 members in a society who are eligible to attend annual meeting. Find the number of ways a 4 members can be selected that

i) No restriction

- ii) If 2 of them will not attend meeting together.
- iii) If 2 of them will always attend meeting together.

Solution:

- i) To select 4 members from 10 members, it can be done by = C(10, 4) = 210 ways.
- ii) If 2 of them will not attend meeting together,

Let A and B denote the 2 members who will not attend meeting together.

i.e. A or B but not both are together , these can be done by $= 2 \times C(8,3) = 112$ ways.

It possible that both will not attend meeting, i.e. Neither A nor B will attend meeting, these can be done by = C(8, 2) = 28 ways.

Therefore total number of ways = 112 + 28 = 140 ways.

iii) If 2 of them will attend meeting together,

Let A and B denote the 2 members who will attend meeting together.

i.e. A or B = C(8, 2) = 28 ways.

It possible that both will not attend meeting, i.e. Neither A nor B will attend meeting, these can be done by = C(8, 4) = 70 ways.

Therefore total number of ways = 28 + 70 = 88 ways.

Example 21: How many diagonal has a regular polygon with n sides?

Solution: The regular polygon with n sides has n vertices. Any two vertices determine either a side or diagonal. Therefore these can be done by $= C(n, 2) = \frac{n(n-1)}{2}$. But there are n sides which are not diagonal.

Therefore total number of diagonals are $=\frac{n(n-1)}{2} - n = \frac{n^2 - n}{2} - \frac{2n}{2} = \frac{n^2 - 3n}{2} = \frac{n(n-3)}{2}$ diagonals.

r-combinations with Repetition Allowed:

Till now, we have seen the formula for the number of combinations when r objects are chosen from the collection of n distinct objects. The following results is very important to find the number of selection of n objects when not all n are distinct.

The number of selection with repetition of r objects chosen from n types of objects is

$$C(n+r-1,r)$$

Example 22: How many ways are there to fill a box with a dozen marbles chosen five different colors of marbles with the requirement that at least one fruit of each colors is picked?

Solution: One can pick one marble of each colors and then the remaining seven marbles in any way. There is no choice in picking one marble of

each type. The choice occurs in picking the remaining 7 marbles from 5 colors. By the result of r-combination with repetition allowed,

These can be done by = C(5 + 7 - 1, 7) = C(11,7) = 330 ways.

Example 23: How many solution does the following equation $x_1 + x_2 + x_3 + x_4 = 15$ have x_1 , x_2 , x_3 , and x_4 are non-negative integers?

Solution: Assume we have four types of unknown x_1 , x_2 , x_3 , and x_4 . There are 15 items or units (since we are looking for an integer solution). Every time an item is selected it adds one to the type it picked it up. Observe that a solution corresponds to a way of selecting 15 items from set of four elements. Therefore, it is equal to r-combinations with repetition allowed from set with four elements, we have

$$C(4 + 15 - 1, 15) = C(18, 15) = C(18, 3) = \frac{18 \times 17 \times 16}{3 \times 2 \times 1} = 816$$

Example 24: In how many ways can a teacher choose one or more students from 5 students?

Solution: Let set of student are 5, therefore total number of subsets are $2^5 = 32$.

To select one or more students, we must deleted empty set,.

Therefore total number of selection = 32 - 1 = 31 ways.

5.8 COMBINATIONS, STIRLING'S APPROXIMATION TO N!

A helpful and commonly used approximate relationship for the evaluation of the factorials of large numbers is Stirling's approximation. It is a good approximation, leading to accurate results even for small values of n. it is given by

$$n! \approx \frac{n^n}{e^n} \sqrt{2\pi n}$$

Where e = 2:71828 is the natural base of logarithms.

5.9 RELATION OF PROBABILITY TO POINT SET THEORY, EULER OR VENN DIAGRAMS AND PROBABILITY

a) Relation of Probability to Point Set Theory:

- In discrete probability we assume well defined experiment such as flipping a coin or rolling a die. Each individual result which could occur is called an outcome. The set of all outcomes is called sample space, and any subset of the sample space is called an event.
- The union of two or more sets is the set that contains all the elements of the two or more sets. Union is denoted by the symbol U.The general

probability addition rule for the union events states that $P(A \cup B) = P(A) + P(B) - P(A \cap B)$. Where $A \cap B$ is the intersection of the two sets.

Euler or Venn Diagrams and Probability:

In probability, a Venn diagram is a figure with one or more circles inside a rectangle that describes logical relation between events. The rectangle in a Venn diagram represents the sample space or the universal set, that is, the set of all possible outcomes. A circle inside the rectangle represents an event, that is, a subset of the sample space. We consider the following Venn diagram involving two events, A and B.



In the above diagram, we have two events A and B within the sample space S(or Universal set)

White and Red region represent event A, Black and Red region represent event B, only Red region represent $A \cap B$, and White, Black and Red region together represents $A \cup B$. Also Blue region represent $\overline{A \cup B}$.

• If the circles do not overlap than A and B are mutually exclusive events.

i.e $A \cap B = \emptyset$.

- If the circles are overlap than A and B are intersecting each other.
 i.eA ∩ B ≠ Ø.
- The region outside both the circles but within the rectangle represents complement of union of both the events. i.e. $\overline{A \cup B}$.

Within each divided region of a Venn diagram, we can add data in one of the following ways:

- The outcomes of the event,
- The number of outcomes in the event,
- The probability of the event.

5.10 LET US SUM UP

In this unit we have learn:

- Basic concept and Definitions of Probability.
- Conditional Probability.
- Independent and Dependent Events, Mutually Exclusive Events.
- Probability Distributions for discrete distribution.
- Mathematical Expectation for probability distribution.
- Relation between Population, Sample Mean, and Variance.
- Combinations, Stirling's Approximation to n!.
- Relations of Probability to Point Set Theory with represent Euler or Venn Diagrams.

5.11 UNIT END EXERCISES

- 1. A card is drawn at random from well shuffled pack of card find the probability that it is red or king card.
- 2. There are 30 tickets bearing numbers from 1 to 15 in a bag. One ticket is drawn from the bag at random. Find the probability that the ticket bears a number, which is even, or a multiple of 3.
- 3. In a group of 200 persons, 100 like sweet food items, 120 like salty food items and 50 like both. A person is selected at random find the probability that the person (i). Like sweet food items but not salty food items (ii). Likes neither.
- 4. A bag contains 7 white balls & 5 red balls. One ball is drawn from bag and it is replaced after noting its color. In the second draw again one ball is drawn and its color is noted. The probability of the event that both the balls drawn are of different colors.
- 5. The probability of A winning a race is $\frac{1}{3}$ & that B wins a race is $\frac{3}{5}$. Find the probability that (a). either of the two wins a race.b), no one wins the race.
- 6. Three machines A, B & C manufacture respectively 0.3, 0.5 & 0.2 of the total production. The percentage of defective items produced by A, B & C is 4, 3 & 2 percent respectively. for an item chosen at random, what is the probability it is defective.
- 7. An urn A contains 3 white &5 black balls. Another urn B contains 5 white & 7 black balls. A ball is transferred from the urn A to the urn B, then a ball is drawn from urn B. find the probability that it is white.
- 8. A husband & wife appear in an interview for two vacancies in the same post. The probability of husband selection is $\frac{1}{7}$ & that of wife's selection is $\frac{1}{5}$. What is the probability that,a). both of them will be

selected.b). only one of them will be selected.c). none of them will be selected?

- 9. A problem statistics is given to 3 students A,B & C whose chances of solving if are $\frac{1}{2}$, $\frac{3}{4}$ & $\frac{1}{4}$ respectively. What is the probability that the problem will be solved?
- 10. A bag contains 8 white & 6 red balls. Find the probability of drawing 2 balls of the same color.
- 11. Find the probability of drawing an ace or a spade or both from a deck of cards?
- 12. A can hit a target 3 times in a 5 shots, B 2 times in 5 shots & C 3 times in a 4 shots. they fire a volley. What is the probability that a).2 shots hit? b). at least 2 shots hit?
- 13. A purse contains 2 silver & 4 cooper coins & a second purse contains 4 silver & 4 cooper coins. If a coin is selected at random from one of the two purses, what is the probability that it is a silver coin?
- 14. The contain of a three urns are : 1 white, 2 red, 3 green balls; 2 white, 1 red, 1 green balls & 4 white, 5 red, 3 green balls. Two balls are drawn from an urn chosen at random. This are found to be 1 white & 1 green. Find the probability that the balls so drawn come from the second urn.
- 15. Three machines A,B& C produced identical items. Of there respective output 2%, 4% & 5% of items are faulty. On a certain day A has produced 30% of the total output, B has produced 25% & C the remainder. An item selected at random is found to be faulty. What are the chances that it was produced by the machine with the highest output?
- 16. A person speaks truth 3 times out of 7. When a die is thrown, he says that the result is a 1. What is the probability that it is actually a 1?
- 17. There are three radio stations A, B and C which can be received in a city of 1000 families. The following information is available on the basis of a survey:
- (a) 1200 families listen to radio station A
- (b) 1100 families listen to radio station B.
- (c) 800 families listen to radio station C.
- (d) 865 families listen to radio station A & B.
- (e) 450 families listen to radio station A & C.
- (f) 400 families listen to radio station B & C.
- (g) 100 families listen to radio station A,B & C.

The probability that a family selected at random listens at least to one radio station.

18. The probability distribution of a random variable x is as follows.

Х	1	3	5	7	9
P(x)	Κ	2k	3k	3k	K

Find value of (i). K (ii). E(x)

- 19. A player tossed 3 coins. He wins Rs. 200 if all 3 coins show tail, Rs. 100 if 2 coins show tail, Rs. 50 if one tail appears and loses Rs. 40 if no tail appears. Find his mathematical expectation.
- 20. The probability distribution of daily demand of cell phones in a mobile gallery is given below. Find the expected mean .

Demand	5	10	15	20
Probability	0.4	0.22	0.28	0.10

- 21. If $P(A) = \frac{4}{15}$, $P(B) = \frac{7}{15}$ and if A and B are independent events, find $(i)P(A \cap B)$, $(ii)P(A \cup B)$, $(iii)P(\overline{A} \cap \overline{B})$.
- 22. If $P(A) = \frac{5}{9}$, $P(\overline{B}) = \frac{2}{9}$ and if A and B are independent events, find $(i)P(A \cap B)$, $(ii)P(A \cup B)$, $(iii)P(\overline{A} \cap \overline{B})$.
- 23. If P(A) = 0.65, P(B) = 0.75 and $P(A \cap B) = 0.45$, where A and B are events of sample space S, find (i)P(A|B), $(ii)P(A \cup B)$, $(iii)P(\overline{A} \cap \overline{B})$.
- 24. A box containing 5 red and 3 black balls, 3 balls are drawn at random from box. Find the expected number of red balls drawn.
- 25. Two fair dice are rolled. X denotes the sum of the numbers appearing on the uppermost faces of the dice. Find the expected value.
- 26. A bag contains 5 black marbles and 6 white marbles. Find the number of ways that five marbles can be drawn from the bag such that it contains i) No restriction ii) no black marbles, iii) 3 black and 2 white, iv) at least 4 black, v) All are of same colors.
- 27. A student is to answer 8 out of 10 questions on an exam. Find the number of ways that the student can chose the 8 questions if i) No restriction, ii) student must answer the first 4 questions, iii) student must answer atleast 4 out of the five questions.
- 28. There are 12 points in a given plane, no three on the same line. i) How many triangle are determine by the points? ii) How many of these triangle contain a particular point as a vertex?
- 29. How many committees of two or more can be selected from 8 people?
- 30. Find the number of combinations if the letters of the letters of the word EXAMINATION taken out at a time.

5.12 LIST OF REFERENCES

- Statistics byMurry R. Spiegel, Larry J. Stephens. Publication McGRAWHILL INTERNATIONAL.
- Fundamental Mathematics and Statistics by S.C. Gupta and V.K kapoor Mathematical Statistics by J.N. Kapur and H.C. Saxena.

ELEMENTARY SAMPLING THEORY

6

Unit Structure

- 6.1 Objective
- 6.2 Introduction
- 6.3 Sampling Theory
 - 6.3.1 Random Samples and Random Numbers,
 - 6.3.2 Sampling With and Without Replacement,
- 6.4 Sampling Distributions,
 - 6.4.1 Sampling Distribution of Means,
 - 6.4.2 Sampling Distribution of Proportions,
 - 6.4.3 Sampling Distributions of Differences and Sums,
- 6.5 Standard Errors,
- 6.6 Summary
- 6.7 Exercise
- 6.8 List of References

6.1 **OBJECTIVE**

After going through this chapter you will able to know:

- Sampling and its requirements in statistics.
- Random sampling with and without replacement.
- Sampling distribution of Mean, Proportions, difference and sum.
- Standard errors in sampling distribution.
- Some software to use for sampling.

6.2 INTRODUCTION

In the previous chapters, we have discussed probability theory. In this chapter, we will introduce some basic concepts in statistics. The basic idea of statistical inference is to assume that the observed data is generated from some unknown probability distribution, which is often assumed to have a known functional form up to some unknown parameters. The purpose of statistical inference is to develop theory and methods to make inference on the unknown parameters based on observed data.

Sampling theory provides the tools and techniques for data collection keeping in mind the objectives to be fulfilled and nature of population. Sample surveys collect information on a fraction of total population whereas in census, the information is collected on the whole population.

The concept of sampling has a huge implementation and its application is seen in many vital fields. The importance of sampling theory is when it comes into play while making statistical analysis with different efficiency levels, there are three different methods of sampling. We have adequately thrown light on the process and methods of sampling.

6.3 SAMPLING THEORY

Often we are interested in drawing some valid inferences about a large group of individuals or objects called population in statistics. Instead of studying the entire population, which may be difficult or even impossible to study, we may study only a small portion of the population. Our objective is to draw valid inferences about certain facts for the population from results found in the sample; a process known as statistical inferences. The process of obtaining samples is called sampling and theory concerning the sampling is called sampling theory.

The sampling theory definition of the statistic is the creation of a sample set. This is recognized as one of the major processes. It retains the accuracy in bringing out the correct statistical information. The population tree is huge set and it turns out to be exhausting for the actual study and estimation process. Both money and time get exhausting in the process. The creation of the sample set saves time and effort and is a vital theory in the process of statistical data analysis.

Process of Sampling:

In this part of the chapter, we will discuss a few details regarding the process of sampling. So the steps are mentioned in the steps below:

- The first step is a wise choice of the population set.
- The second step is focusing on the sample set and the size of it.
- Then, one needs to choose an identifiable property based on which the samples will be created out of the population set.
- Then, the samples can be chosen using any of the types of sampling theory Simple random, systematic, or stratified. Each of them is thoroughly discussed in the article ahead.
- Checking the inaccuracy, if there is any.
- Hence, the set is achieved in the result.

Sampling can be done in their different method and they are given below:

- 1. Simple random type.
- 2. Systematic Sampling.
- 3. Stratified sampling.

6.3.1 Random Samples and Random Numbers:

Definition: Simple random sampling is defined as a sampling technique where every item in the population has an even chance and likelihood of
being selected in the sample. Here the selection of items entirely depends on luck or probability, and therefore this sampling technique is also sometimes known as a method of chances. For e.g. Using the lottery method is one of the oldest ways and is a mechanical example of random sampling. In this method, the researcher gives each member of the population a number. Researchers draw numbers from the box randomly to choose samples. The use of random numbers is an alternative method that also involves numbering the population. The use of a number table similar to the one below can help with this sampling technique.

Simple random sampling (SRS) is a method of selection of a sample comprising of n a number of sampling units out of the population having N number of sampling units such that every sampling unit has an equal chance of being chosen.

Simple random sampling methods:

Researchers follow these methods to select a simple random sample:

- 1. They prepare a list of all the population members initially, and then each member is marked with a specific number (for example, there are nth members, then they will be numbered from 1 to N).
- 2. From this population, researchers choose random samples using two ways: random number tables and random number generator software. Researchers prefer a random number generator software, as no human interference is necessary to generate samples.

Advantages of simple random sampling:

- 1. It is a fair method of sampling, and if applied appropriately, it helps to reduce any bias involved compared to any other sampling method involved.
- 2. Since it involves a large sample frame, it is usually easy to pick a smaller sample size from the existing larger population.
- 3. The person conducting the research doesn't need to have prior knowledge of the data he/ she is collecting. Once can ask a question to gather the researcher need not be subject expert.
- 4. This sampling method is a fundamental method of collecting the data. You don't need any technical knowledge. You only require essential listening and recording skills.
- 5. Since the population size is vast in this type of sampling method, there is no restriction on the sample size that the researcher needs to create. From a larger population, you can get a small sample quite quickly.
- 6. The data collected through this sampling method is well informed; more the sample better is the quality of the data.

Disadvantages:

1. Sampling is not feasible where knowledge about each element or unit or a statistical population is needed.

- 2. The sampling procedures must be correctly designed and followed otherwise, what we call as wild sample would crop up with misleading results.
- 3. Each type of sampling has got its own limitations.
- 4. There are numerous situations in which units, to be measured, are highly variable. Here a very large sample is required in order to yield enough cases for achieving statistically reliable information.
- 5. To know certain population characteristics like population growth rate, population density etc. census of population at regular intervals is more appropriate than studying by sampling.

6.3.2 Sampling With and Without Replacement:

Selection with Replacement (SWR): In this case, a unit is selected from a population with a known probability and the unit is returned to the population before the next selection is made (after recording its characteristic). Thus, in this method at each selection, the population size remains constant and the probability at each selection or draw remains the same. Under this sampling plan, a unit has chances of being selected more than once. For example a card is randomly drawn from a pack of cards and placed back in the pack, after noting its face value before the next card is drawn. Such a sampling method is known as sampling with replacement. There are N^n possible samples of size n from a population of N units in case of sampling with replacement.

Sampling without replacement (SWOR): In this selection procedure, if a unit from a population of size N selected, it is not returned to the population. Thus, for any subsequent selection, the population size is reduced by one. Obviously, at the time of the first selection, the population size is N and the probability of a unit being selected randomly is 1/N; for the second unit to be randomly selected, the population size is (N - 1)and the probability of selection of any one of the remaining sampling unit is 1/(N-1), similarly at the third draw, the probability of selection is 1/(N-2) and so on.

6.4 SAMPLING DISTRIBUTIONS

Sampling distribution is a statistic that determines the probability of an event based on data from a small group within a large population. Its primary purpose is to establish representative results of small samples of a comparatively larger population. Since the population is too large to analyze, the smaller group is selected and repeatedly sampled, or analyzed. The gathered data, or statistic, is used to calculate the likely occurrence, or probability, of an event. Using a sampling distribution simplifies the process of making inferences, or conclusions, about large amounts of data.

The idea behind a sampling distribution is that when you have a large amount of data (gathered from a large group, the value of a statistic from random samples of a small group will inform you of that statistic's value for the entire group. Once the data is plotted on a graph, the values of any given statistic in random samples will make a normal distribution from which you can draw inferences.

Each random sample selected will have a different value assigned to the statistic being studied. For example, if you randomly sample data three times and determine the mean, or the average, of each sample, all three means are likely to be different and fall somewhere along the graph. That's variability. You do that many times, and eventually the data you plot should look like a bell curve. That process is a sampling distribution.

Factors that influence sampling distribution:

The sampling distribution's variability can be measured either by standard deviation, also called "standard error of the mean," or population variance, depending on the context and inferences you are trying to draw. They both are mathematical formulas that measure the spread of data points in relation to the mean.

There are three primary factors that influence the variability of a sampling distribution. They are:

- The number observed in a population: This variable is represented by "*N*." It is the measure of observed activity in a given group of data.
- **The number observed in the sample:** This variable is represented by "*n*." It is the measure of observed activity in a random sample of data that is part of the larger grouping.
- The method of choosing the sample: How the samples were chosen can account for variability in some cases.

Types of distributions:

There are three standard types of sampling distributions in statistics.

- 1. Sampling Distribution of Means.
- 2. Sampling Distribution of Proportions.
- 3. Sampling Distributions of Differences and Sums.

6.4.1 Sampling Distribution of Means:

The most common type of sampling distribution is of the mean. It focuses on calculating the mean of every sample group chosen from the population and plotting the data points. The graph shows a normal distribution where the center is the mean of the sampling distribution, which represents the mean of the entire population.

The mean of the sampling distribution of the mean is the mean of the population from which the scores were sampled. Therefore, if a population has a mean μ , then the mean of the sampling distribution of the mean is also μ . The symbol $\mu_{\bar{X}}$ is used to refer to the mean of the sampling

distribution of the mean. Therefore, the formula for the mean of the sampling distribution of the mean can be written as:

 $\mu_{\bar{X}} = \mu$

The standard deviation of the sampling distribution of the mean is computed as follows:

$$\sigma_{\bar{X}} = \frac{\sigma}{\sqrt{N}}$$

That is, the standard deviation of the sampling distribution of the mean is the population Standard deviation divided by \sqrt{N} , the sample size (the number of scores used to compute a mean). Thus, the larger the sample size, the smaller the Standard deviation of the sampling distribution of the mean.

For sampling is drawn without replacement,

The mean of the sampling distribution of means given by

$$\mu_{\bar{X}} = \mu$$

The standard deviation of the sampling distribution of means is given by

$$\sigma_{\bar{X}} = \frac{\sigma}{\sqrt{N}} \sqrt{\frac{N-n}{N-1}}.$$

Example 1: A population consists of the five numbers 5, 6, 7, 12, and 15. Consider all possible samples of size 2 that can be drawn with replacement from this population. Find

- a) the mean of the population,
- b) the standard deviation of the population,
- c) the mean of the sampling distribution of means,
- d) the standard deviation of the sampling distribution of means.

Solution: Here population N = 5, sample size n = 2.

a) The mean of the population is given by

$$\mu = \frac{5+6+7+12+15}{5} = \frac{45}{5} = 9$$

b) The standard deviation of the population is given by

$$\sigma = \sqrt{\frac{(x - \bar{x})^2}{N}}$$

$$\sigma = \sqrt{\frac{(5-9)^2 + (6-9)^2 + (7-9)^2 + (12-9)^2 + (15-9)^2}{5}}$$
$$\sigma = \sqrt{\frac{16+9+4+9+36}{5}}$$

$$\sigma = \sqrt{\frac{74}{5}} = 3.85$$

c) The mean of the sampling distribution of means:

Hare 5(5) = 25 samples of size 2, that can be drawn with replacement, i.e. Samples are

$$(5,5), (5,6), (5,7), (5,12), (5,15), (6,5), (6,6), (6,7), (6,12), (6,15),$$

 $(7,5), (7,6), (7,7), (7,12), (7,15),$
 $(12,5), (12,6), (12,7), (12,12), (12,15), (15,5), (15,6), (15,7),$
 $(15,12), (15,15)$

Therefore, the corresponding sample means are

The mean of the sampling distribution of means is given by

$$\mu_{\bar{X}} = \frac{Sum \ of \ sample \ means}{25} = \frac{225}{25} = 9.$$

i.e. $\mu_{\bar{X}} = \mu$

d) The standard deviation of the sampling distribution of means.

$$\sigma_{\bar{X}} = \sqrt{\frac{(5-9)^2 + (5.5-9)^2 + (6-9)^2 + \dots + (11-9)^2 + (13.5-9)^2 + (15-9)^2}{25}}$$
$$= \sqrt{\frac{185}{25}} = \sqrt{7.4} = 2.72$$
Therefore, $\sigma_{\bar{X}} = \frac{\sigma}{\sqrt{N}} = \frac{3.85}{\sqrt{2}} = 2.72.$

Example 2: A population consists of the five numbers 7, 9, 10, 14, and 20. Consider all possible samples of size 2 that can be drawn without replacement from this population. Find

- a) the mean of the population,
- b) the standard deviation of the population,
- c) the mean of the sampling distribution of means,

d) the standard deviation of the sampling distribution of means.

Solution : Here population N = 5, sample size n = 2.

a) The mean of the population is given by

$$\mu = \frac{7+9+10+14+20}{5} = \frac{60}{5} = 12$$

b) The standard deviation of the population is given by

$$\sigma = \sqrt{\frac{(x - \bar{x})^2}{N}}$$

$$\sigma = \sqrt{\frac{(7 - 12)^2 + (9 - 12)^2 + (10 - 12)^2 + (14 - 12)^2 + (20 - 12)^2}{5}}$$

$$\sigma = \sqrt{\frac{25 + 9 + 4 + 4 + 64}{5}}$$

$$\sigma = \sqrt{\frac{106}{5}} = 4.60$$

c) The mean of the sampling distribution of means:

Here ${}^{5}C_{2} = 10$ samples of size 2, that can be drawn without replacement, i.e.(all sample are distinct selection). Samples are

Therefore, the corresponding sample means are

8, 8.5, 10.5, 13.5, 9.5, 11.5, 14.5, 12, 15, 17.

The mean of the sampling distribution of means is given by

$$\mu_{\bar{X}} = \frac{Sum \ of \ sample \ means}{10} = \frac{120}{10} = 12.$$

i.e. $\mu_{\bar{X}} = \mu$

d) The standard deviation of the sampling distribution of means.

$$\sigma_{\bar{X}} = \sqrt{\frac{(8-12)^2 + (8.5-12)^2 + (10.5-12)^2 + \dots + (12-12)^2}{+(15-12)^2 + (17-12)^2}}$$

$$= \sqrt{\frac{79.5}{10}} = \sqrt{7.95} = 2.81$$

Therefore, $\sigma_{\bar{X}} = \frac{\sigma}{\sqrt{N}} \sqrt{\frac{N-n}{N-1}} = \frac{4.6}{\sqrt{2}} \sqrt{\frac{5-2}{5-1}} = 2.81.$

6.4.2 Sampling Distribution of Proportions:

This sampling distribution focuses on proportions in a population. Samples are selected and their proportions are calculated. The mean of the sample proportions from each group represent the proportion of the entire population.

Suppose random samples of size n are drawn from a population in which the proportion with a characteristic of interest is p.

The Sampling Distribution of Proportion measures the proportion of success, i.e. a chance of occurrence of certain events, by dividing the number of successes i.e. chances by the sample size 'n'. Thus, the sample proportion is defined as

$$p = \frac{x}{n}$$

Therefore the mean μ_p and standard deviation σ_p are given by

$$\mu_p = p,$$
$$\sigma_p = \sqrt{\frac{pq}{n}}$$

Where q is probability of non-occurrence of event, which is given byq = 1 - p.

The following formula is used when population is finite, and the sampling is made without the replacement:

$$\sigma_p = \sqrt{\frac{N-n}{N-1}} \sqrt{\frac{pq}{n}}$$

If *n* is large, and *p* is not too close to 0 or 1, the binomial distribution can be approximated by the normal distribution. Practically, the Normal approximation can be used when both $np \ge 10$.

Once we have the mean and standard deviation of the survey data, we can find out the probability of a sample proportion. Here, the Z score conversion formula will be used to find out the required probability, i.e.

$$Z = \frac{X-\mu}{\sigma}.$$

Example 3:A random sample of 100 students is taken from the population of all part-time students in the Maharashtra, for which the overall proportion of females is 70%. Find sample mean and sample standard deviation.

Solution: Here $n = 100, p = 70\% = \frac{70}{100} = 0.7$ $\therefore q = 1 - p = 1 - 0.7 = 0.3$

the mean μ_p is given by

$$\mu_p = p = 0.7$$

and standard deviation σ_p is given by

$$\sigma_p = \sqrt{\frac{pq}{n}} = \sqrt{\frac{0.7 \times 0.3}{100}} = \sqrt{\frac{0.21}{100}} = 0.0458.$$

Example 4: Suppose it is known that 47% of Indian own smart phone. If a random sample of 50 Indians were surveyed, what is the probability that the proportion of the sample who owned smart phone is between 50% and 54%?

Solution: Here we have, n = 50, p = 47% = 0.47,

$$\therefore q = 1 - p = 1 - 0.47 = 0.53.$$

Now, we should check our conditions for the sampling distribution of the sample proportion.

$$np = 50 \times 0.47 = 23.5 \ge 10, nq = 50 \times 0.53 = 26.5 \ge 10.$$

Since both the condition satisfy,

 \therefore The sampling distribution that is approximately normal with mean and standard deviation,

$$\mu_P = \mu = 0.47,$$

$$\sigma = \sqrt{\frac{pq}{n}} = \sqrt{\frac{0.47 \times 0.53}{50}} = 0.07$$

Now, The probability that the proportion of the sample who owned smart phone is between 50% and 54% is given by

$$P(0.50 < Z < 0.54) = P\left(\frac{0.50 - 0.47}{0.07} < Z < \frac{0.54 - 0.47}{0.07}\right)$$
$$= P(0.429 < Z < 1)$$
$$= P(Z < 1) - P(Z < 0.429)$$
$$= 0.8413 - 0.6627$$
$$= 0.1786$$

 \therefore If the true proportion of Indians who own smart phone is 47%, then there would be a 17.86% chance that we would see a sample proportion between 50% and 54% when the sample size is 50.

6.4.3 Sampling Distributions of Differences and Sums:

Statistical analyses are very often concerned with the difference between means. A typical example is an experiment designed to compare the mean of a control group with the mean of an experimental group. Inferential statistics used in the analysis of this type of experiment depend on the sampling distribution of the difference between means.

The sampling distribution of the difference between means can be thought of as the distribution that would result if we repeated the following three steps over and over again:

- 1. Sample n_1 scores from Population 1 and n_2 scores from Population 2.
- 2. compute the means of the two samples M_1 and M_2 .
- 3. compute the difference between means, $M_1 M_2$. The distribution of the differences between means is the sampling distribution of the difference between means.

As you might expect, the mean of the sampling distribution of the difference between means is:

$$\mu_{M_1 - M_2} = \mu_{M_1} - \mu_{M_2}$$

Which says that the mean of the distribution of differences between sample means is equal to the difference between population means.

From the variance sum law, we know that:

$$\sigma^2_{M_1 - M_2} = \sigma^2_{M_1} + \sigma^2_{M_2}$$

We can write the formula for the standard deviation of the sampling distribution of the difference between means as

$$\therefore \sigma_{M_1-M_2} = \sqrt{\sigma_{M_1}^2 + \sigma_{M_2}^2}$$

Similarly we say about the sampling distribution of the sum between means is given by:

$$\mu_{M_1+M_2} = \mu_{M_1} + \mu_{M_2}$$
$$\sigma_{M_1+M_2} = \sqrt{\sigma_{M_1}^2 + \sigma_{M_2}^2}$$

Example 5: Let S_1 be a variable that stands for any of the elements of the population 4, 6, 8 and S_2 be a variable that stands for any of the elements of the population 3, 5. Compute a) μ_{S_1} , b) μ_{S_2} , c) $\mu_{S_1-S_2}$, d) σ_{S_1} , e) σ_{S_2} , and f) $\sigma_{S_1-S_2}$.

Solution:

a) Here S_1 has sample of population 4, 6, 8.

$$\mu_{S_1} = \frac{4+6+8}{3} = \frac{18}{3} = 6.$$

b) Here S_2 has sample of population 3, 5.

$$\mu_{S_2} = \frac{3+5}{2} = 4$$

c) The population consisting of the differences of any member of S_1 and any member of S_2 mean is given by

$$\mu_{S_1-S_2} = \frac{(4-3) + (6-3) + (8-3) + (4-5) + (6-5) + (8-5)}{6}$$
$$= \frac{12}{6} = 2.$$

Therefore,

we can verify that
$$\mu_{S_1-S_2} = \mu_{S_1} - \mu_{S_2} = 6 - 4 = 2.$$

d)
$$\sigma_{S_1} = \sqrt{\frac{(4-6)^2 + (6-6)^2 + (8-6)^2}{3}} = \sqrt{\frac{8}{3}} = \sqrt{2.67} = 1.63$$

e) $\sigma_{S_2} = \sqrt{\frac{(3-4)^2 + (5-4)^2}{2}} = \sqrt{\frac{2}{2}} = 1.$

f) The population consisting of the differences of any member of S_1 and any member of S_2 Standard deviation is given by

$$\sigma_{S_1 - S_2} = \sqrt{\frac{(1 - 2)^2 + (3 - 2)^2 + (5 - 2)^2 + (-1 - 2)^2 + (1 - 2)^2 + (3 - 2)^2}{6}}$$
$$= \sqrt{\frac{22}{6}} = \sqrt{\frac{11}{3}} = 1.91$$

Therefore, we can verify that,

$$\therefore \sigma_{S_1 - S_2} = \sqrt{\sigma_{S_1}^2 + \sigma_{S_2}^2} = \sqrt{2.67 + 1} = \sqrt{3.67} = 1.91$$

Example 6: The battery life of smart phone manufacturer A have a mean lifetime of 1050 days with a standard deviation of 150 days, while those of manufacturer B have a mean lifetime of 800 days with a standard deviation of 120 days. If random samples of 100 batteries of each brand are tested, what is the probability that the brand A batteries will have a mean lifetime that is at least (a) 200 days and (b) 280 days more than the brand B batteries?

Solution: Let $\overline{X_A}$ and $\overline{X_B}$ denote the mean lifetimes of samples A and B, respectively. Then

$$\mu_{\overline{X_A} - \overline{X_B}} = \mu_{\overline{X_A}} - \mu_{\overline{X_B}} = 1050 - 800 = 250 \ days.$$

$$\sigma_{\overline{X_A} - \overline{X_B}} = \sqrt{\frac{\sigma_{\overline{X_A}}^2}{N_A} + \frac{\sigma_{\overline{X_B}}^2}{N_B}} = \sqrt{\frac{(150)^2}{100} + \frac{(120)^2}{100}} = \sqrt{\frac{36900}{100}} = \sqrt{\frac{36900}{100}} = \sqrt{\frac{36900}{100}}$$

Therefore, the standardized variable for the difference in means is

$$\boldsymbol{Z} = \frac{(\overline{X_A} - \overline{X_B}) - (\mu_{\overline{X_A} - \overline{X_B}})}{\sigma_{\overline{X_A} - \overline{X_B}}}$$

and is very closely normally distributed.

a) the probability that the brand A batteries will have a mean lifetime that is at least 1200 days,

$$P(Z > 200) = 0.5 + P\left(Z = \frac{200 - 250}{19.21}\right) = 0.5 + P(Z > -2.6)$$
$$= 0.5 + 0.4953 = 0.9953.$$

b) the probability batteries will have a mean lifetime that is at least 960 days more than the brand B batteries,

$$P(Z > 280) = 0.5 - P\left(Z = \frac{280 - 250}{19.21}\right) = 0.5 - P(Z = 1.56)$$
$$= 0.5 - 0.4406 = 0.0594.$$

6.5 STANDARD ERRORS

Another measure is standard error, which is the standard deviation of the sampling distribution of an estimator. The idea is that if we draw a number of repeated samples of fixed size n from a population having a mean μ and variance σ^2 , each simple mean, say \bar{x} , will have a different value. Here \bar{x} it is a random variable and hence it has a distribution. The standard deviation of \bar{x} is called standard error. It has been proved that the standard error ' $\sigma_{\bar{x}}$ of the mean \bar{x} based on a sample of size n is,

$$\sigma_{\bar{x}} = \frac{\sigma}{\sqrt{n}}$$

From above formula, it is obvious that the larger the sample size, the smaller the standard error and vice-versa. The advantage of considering standard error instead of a standard deviation is that this measure is not influenced by the extreme values present in a population under consideration.

In reality neither we use σ to calculate the standard error of \bar{x} nor we take more than one sample. As a matter of fact, what we do is, that we select only one sample, find its standard deviation *s* and use the following formula to find out the standard error of \bar{x} .i.e.

$$S.E.(\bar{x}) = \frac{s}{\sqrt{n}}$$

Standard error is commonly used in testing of hypothesis and interval estimation. Many distributions, which are originally not normally distributed, have been taken as normal by considering the distribution of mean \bar{x} for a large n.

6.6 SUMMARY

In this unit we have learn:

- In sampling theory we have Random Samples, Sampling With and Without Replacement of sample.
- Sampling Distributions and its types.
- Standard Errors of sampling distribution.

6.7 EXERCISE

- 1. A population consists of the four numbers 8, 11, 12, and 19. Consider all possible samples of size 2 that can be drawn with replacement from this population. Find
 - a) the mean of the population,
 - b) the standard deviation of the population,
 - c) the mean of the sampling distribution of means,
 - d) the standard deviation of the sampling distribution of means.
- 2. A population consists of the seven numbers 3, 5, 7, 9, 11, 13, and 15. Consider all possible samples of size 2 that can be drawn without replacement from this population. Find
 - a) the mean of the population,
 - b) the standard deviation of the population,
 - c) the mean of the sampling distribution of means,
 - d) the standard deviation of the sampling distribution of means.
- 3. Let S_1 be a variable that stands for any of the elements of the population 5, 8, 12 and S_2 be a variable that stands for any of the elements of the population 2, 6. Compute a) μ_{S_1} , b) μ_{S_2} , c) $\mu_{S_1-S_2}$, d) σ_{S_1} , e) σ_{S_2} , and f) $\sigma_{S_1-S_2}$.
- 4. A certain type of electric light bulb has a mean lifetime of 1500 h and a standard deviation of 150 h. Three bulbs are connected so that when one burns out, another will go on. Assuming that the lifetimes are normally distributed, what is the probability that lighting will take place for (a) at least 5000 h and (b) at most 4200 h?
- 5. Two distances are measured as 27.3 centimeters (cm) and 15.6 cm with standard deviations (standard errors) of 0.16 cm and 0.08 cm,

respectively. Determine the mean and standard deviation of (a) the sum and (b) the difference of the distances.

- 6. A and B play a game of "heads and tails," each tossing 50 coins. A will win the game if she tosses 5 or more heads than B; otherwise, B wins. Determine the odds against A winning any particular game.
- 7. The average income of men in a city is Rs. 20,000 with standard deviation Rs. 10,500 and the average income of women is Rs. 16,000 and standard deviation Rs. 8,000. There are 100 sample selected from population. Find the probability of income between Rs. 15,000 to Rs. 18000.
- 8. A sample of 300 items selected at random had 32 defective items. Find mean and standard deviation of sampling distribution of proportion.
- 9. Ball bearings of a given brand weigh 0.50 g with a standard deviation of 0.02 g. What is the probability that two lots of 1000 ball bearings each will differ in weight by more than 2 g?
- 10. Find the probability that in 120 tosses of a fair coin (a) less than 40% or more than 60% will be heads and (b) 5/8 or more will be heads.

6.8 LIST OF REFERENCES

- Statistics by Murry R. Spiegel, Larry J. Stephens. Publication McGRAWHILL INTERNATIONAL.
- Fundamental Mathematics and Statistics by S.C. Gupta and V.K kapoor
- Mathematical Statistics by J.N. Kapur and H.C. Saxena.

UNIT III

STATISTICAL ESTIMATION THEORY

Unit Structure

- 7.0 Objectives
- 7.1 Basic definitions
 - 7.1.1 Population
 - 7.1.2 Sample
 - 7.1.3 Parameter
 - 7.1.4 Statistic
 - 7.1.5 Sampling distribution
 - 7.1.6 Parameter Space
 - 7.1.7 Estimator
 - 7.1.8 Estimate
- 7.2 Point estimation
 - 7.2.1 Unbiasedness
 - 7.2.2 Consistency
 - 7.2.3 Efficiency
 - 7.2.4 Minimum variance unbiased estimator
 - 7.2.5 Uniformly minimum variance unbiased estimator
 - 7.2.6 Likelihood function
 - 7.2.7 Sufficiency
- 7.3 Interval estimation
 - 7.3.1 Probable error
- 7.4 Summary
- 7.5 Exercise
- 7.6 References

7.0 OBJECTIVES

- To understand basic definitions related to point estimation.
- To find the best point estimators to represent population characteristics.
- In this chapter, students can learn the requirements of a good estimator.

• To find an appropriate confidence interval for the population parameters.

7.1 BASIC DEFINITIONS

7.1.1 Population:

- A collection of all well-defined objects under study is called population.
- <u>*Example:*</u> Suppose we want to study the economic conditions of primary teachers in Maharashtra, then the group of all primary teachers in the state of Maharashtra is a population.

7.1.2 Sample:

- A well defined finite subset of the population is called a sample.
- <u>Example</u>: Suppose we want to study the economic conditions of primary teachers in the state of Maharashtra, then the few primary teachers (set of few teachers) in the state of Maharashtra forms a sample.

7.1.3 Parameter:

• An unknown constant of a population that summarises or describes an aspect of the population (such as a mean or a standard deviation) is called parameter. Let $f(x, \theta)$ be the pdf of a random variable 'X' having an unknown constant θ .

7.1.4 Statistic:

• Any function of a sample value (observed value) is called a statistic. The sample statistic is constants but it differ from sample to sample.

7.1.5 Sampling distribution:

• The probability distribution of the sample statistic is called a sampling distribution.

7.1.6 Parameter space:

- The set of all admissible values of a parameter of the distribution is called parameter space. It is denoted by Θ.
- <u>Example</u>: X ~ Normal (μ, σ^2) $\Theta = \{(\mu, \sigma^2) / -\infty < \mu < \infty, \sigma > 0\}$

7.1.7 Estimator:

• Let $x_1, x_2, ..., x_n$ be a sample of size n taken from a distribution having pdf $f(x, \theta)$ where $\theta \in \Theta$ is an unknown parameter. A function $T = T(x_1, x_2, ..., x_n)$ which maps sample space (S) to parameter space Θ is called an estimator. In other words, If a statistic $T = T(x_1, x_2, ..., x_n)$ is used to estimate θ , and its value belongs to parameter space then it is said to be an estimator of θ .

7.1.8 Estimate:

• A particular value of an estimator corresponding to the given sample values is called an estimate of the population parameter.

In the theory of estimation, there are two parts, 1) Point Estimation, 2) Interval Estimation.

7.2 POINT ESTIMATION

- Let $x_1, x_2, ..., x_n$ be a sample of size n taken from a distribution having pdf (probability density function) $f(x, \theta)$ where $\theta \in \Theta$ is an unknown parameter. The method of using sample statistic 'T' to estimate the value of parameter θ , which is a point on real line **R**, is called "Point Estimation".
- Requirements of good and reliable estimators:
 - 1. Unbiasedness
 - 2. Consistency
 - 3. Efficiency
 - 4. Sufficiency

7.2.1 Unbiasedness:

An estimator T = T $(x_1, x_2, ..., x_n)$ is said to be an unbiased estimator of θ iff;

$$E(T) = \theta, \quad \forall \ \theta \in \Theta,$$

a parameteric function $\phi(\theta)$ is said to be estimable if there exists a statistic h (T) such that, $E(h(T)) = \phi(\theta) \quad \forall \theta \in \Theta$.

7.2.1.1 The bias of an estimator:

An estimator T = T ($x_1, x_2, ..., x_n$) is said to be a biased estimator with a bias $\frac{b(\theta)}{n}$ if;

$$E(T) = \theta + \frac{b(\theta)}{n} \quad \forall \theta \in \Theta$$

Or

$$E(T - \theta) = \frac{b(\theta)}{n} \quad \forall \ \theta \in \Theta$$

- If $b(\theta) > 0$, then estimator T is called biased estimator with an upward (positive) bias $\frac{b(\theta)}{n}$.
- If $b(\theta) < 0$, then estimator T is called biased estimator with a downward (negative) bias $\frac{b(\theta)}{n}$.
- If $b(\theta) = 0$, then estimator T is called an unbiased estimator.

• <u>Example</u>: Let x_1, x_2, x_3 be independent observation from Poisson (λ), then show that, $T = \frac{x_1 + x_2 + x_3}{3}$ is an unbiased estimator of λ .

Since x_1 , x_2 , x_3 are i.i.d. Poisson (λ).

$$E[X_1] = E[X_2] = E[X_3] = \lambda$$

Consider;

$$E[T] = E\left[\frac{x_1 + x_2 + x_3}{3}\right] = \frac{1}{3} E[x_1 + x_2 + x_3] = \lambda$$

Hence, T is an unbiased estimator of λ .

7.1.1.1 MSE of an estimator:

• Mean square error (MSE) of an estimator $T = T(x_1, x_2, ..., x_n)$ is,

$$MSE = E(T - \theta)^2$$

7.1.1 Consistency:

• A sequence of the estimator $T_n = T(x_1, x_2, ..., x_n)$ is said to be a consistent estimator for parameter θ if any given $\epsilon > 0$,

$$P\left[|T_n - \theta| < \epsilon\right] \to 1 \quad as \quad n \to \infty \quad \forall \epsilon > 0$$

- Difference between $T_n \& \theta$ becomes smaller and smaller as n goes to a large number (infinity).
- A sequence of an estimator $T_n = T(x_1, x_2, ..., x_n)$ is said to be a consistent estimator for parameter θ if;

$$E(T_n) = \theta \quad \forall \ \theta \in \Theta$$
$$V[T_n] \to 0 \quad as \ n \to \infty$$

• *Example*: Show that the sample mean is a consistent estimator of the population mean.

Let $x_1, x_2, ..., x_n$ be a random sample of size n from a population with mean μ and variance σ^2 .

First, we have to define the sample mean,

$$\overline{x} = \frac{\sum_{i=1}^{n} x_i}{n}$$

$$E[x_i] = \mu \quad \& V[x_i] = \sigma^2 \qquad \forall i = 1, 2, ..., n$$

$$E[\overline{x}] = E\left[\frac{\sum_{i=1}^{n} x_i}{n}\right]$$

$$= \frac{\sum E[x_i]}{n}$$

$$= \frac{n\mu}{n}$$

 $E[\bar{x}] = \mu$

Consider,

$$V [\bar{x}] = V \left[\frac{\sum_{i=1}^{n} x_{i}}{n}\right]$$
$$= \frac{1}{n^{2}} \sum V (x_{i})$$
$$= \frac{1}{n^{2}} n\sigma^{2}$$
$$V [\bar{x}] = \frac{\sigma^{2}}{n}$$
$$\text{Lim}_{n \to \infty} E [\bar{x}] = \mu$$
&
$$V [\bar{x}] = \frac{\sigma^{2}}{n} \to 0.$$

Therefore, the sample mean is a consistent estimator of the population mean.

• <u>**Remark**</u>: If T_n is a consistent estimator for θ , then $\phi(T_n)$ is a consistent estimator for $\phi(\theta)$, where ϕ is a continuous function.

7.1.2 Efficiency:

Lim_n

• An estimator T_1 is said to be more efficient than estimator T_2 of parameter θ if $V(T_1) < V(T_2)$. The relative efficiency of T_2 with respect to T_1 is defined as,

$$e = \frac{V[T_1]}{V[T_2]}.$$

• <u>Example</u>: Let $x_1, x_2, ..., x_n$ be a random sample of size n from a Normal (μ, σ^2) . Then select the most efficient estimator between \bar{x} (Mean) & x_m (Median).

We know that,

$$E [\bar{x}] = \mu$$

$$V [\bar{x}] = \frac{\sigma^2}{n} \to 0 \quad as \quad n \to \infty$$

$$\& \quad E [x_m] = \mu$$

$$V [x_m] = \left(\frac{\pi}{2}\right) \cdot \left(\frac{\sigma^2}{n}\right).$$

Now we have to calculate efficiency as,

$$e = \frac{V[\bar{x}]}{V[x_m]} = \frac{2}{\pi} < 1.$$

Therefore, the sample mean is a more efficient estimator than the sample median.

7.1.3 Minimum Variance Unbiased Estimator (MVUE):

• An estimator $T = T(x_1, x_2, ..., x_n)$ is said to be minimum variance unbiased estimator of parameter θ if,

1) $E[T] = \theta \quad \forall \theta \in \Theta$ and,

2) V[T] < V[T']; where, T' is any other unbiased estimator of θ .

7.1.4 Uniformly minimum variance unbiased estimator (UMVUE):

Let θ be the unknown parameter & Θ be the parameter space of θ . Let $U(\theta)$ be the set of a class of unbiased estimator of $T(\theta)$ such that,

$$E\left[T^2\right] < \infty \quad \forall \ \theta \in \Theta$$

i.e.,

$$U(\theta) = \{T: E[T] = \theta, \qquad E[T^2] < \infty \ \forall \theta \in \Theta\}$$

Then, $T_0 \in U(\theta)$ is UMVUE of θ if,

$$E [T_0 - \theta]^2 \le E [T - \theta]^2 \qquad \forall \theta \in \Theta \& T \in U(\theta).$$

7.1.5 Likelihood function:

• Let $x_1, x_2, ..., x_n$ be a sample of size n taken from a distribution having pdf $f(x, \theta)$ where $\theta \in \Theta$ is an unknown parameter. Then the likelihood function of $x_1, x_2, ..., x_n$ is defined as

$$L(X_1, X_2, \dots, X_n, \theta) = f(X_{1,\theta}) \cdot f(X_{2,\theta}) \cdot f(X_{3,\theta}) \dots f(X_{n,\theta})$$
$$L(\underline{X}, \theta) = \prod_{i=1}^n f(x_i, \theta).$$

• T is said to be the maximum likelihood estimator of θ , which maximizes the likelihood function $L(\underline{X}, \theta)$.

7.1.6 Sufficiency:

- Definition I A statistic T = T(x₁, x₂, ..., x_n) based on a sample of size n having pmf/pdf f(X, θ) θ ∈ Θ is said to sufficient statistic for θ if and only if the information contains in T about θ is same as the information contains in x₁, x₂, ..., x_n about θ.
- **Definition II** A statistic $T = T(x_1, x_2, ..., x_n)$ based on a sample of size n having pmf/pdf $f(X, \theta)$ $\theta \in \Theta$ is said sufficient statistic for θ , if and only if the conditional distribution of $x_1, x_2, ..., x_n$ given T is independent of θ .
- Definition III (Neyman factorization criterion) If $x_1, x_2, ..., x_n$ is a sample of size n having pmf/pdf $f(X, \theta) \ \theta \in \Theta$ and $T = T(x_1, x_2, ..., x_n)$ be a statistic which is said to be sufficient for θ if and only if the joint probability distribution function of $x_1, x_2, ..., x_n$ can be expressed as a product of a function of T and θ , and function of $x_1, x_2, ..., x_n$ only.

i.e.,
$$L(\underline{X}, \theta) = g(T, \theta) \cdot h(\underline{X}),$$

then *T* is said to be sufficient statistic for θ , where $g(T, \theta)$ is a function of *T* & θ only and h(X) is a function of $x_1, x_2, ..., x_n$ only.

• **<u>Example</u>**: Let $x_1, x_2, ..., x_n$ be a random sample from a population having pdf,

$$f(x) = \begin{cases} \theta x^{\theta - 1}, & 0 \le x \le 1\\ 0, & otherwise \end{cases}$$

The likelihood function of $x_1, x_2, ..., x_n$ is,

$$L\left(\underline{X},\theta\right) = \prod_{i=1}^{n} \theta x_i^{\theta-1}$$
$$= \theta^n \prod_{i=1}^{n} x_i^{\theta-1}$$
$$= \left[\theta^n \prod x_i^{\theta}\right] \left[\frac{1}{\prod x_i}\right]$$
$$L\left(\underline{X},\theta\right) = g(T,\theta) \cdot h(\underline{X})$$

Where, $(T, \theta) = \theta^n \prod x_i^{\theta}$, $h(\underline{X}) = \frac{1}{\prod x_i}$

Therefore, from the Neyman factorization criterion $\prod x_i$ is sufficient statistic for θ .

7.2 INTERVAL ESTIMATION

• A confidence interval for an unknown parameter is an interval of possible values for the parameter. It is constructed so that, with a chosen degree of confidence, the actual value of the parameter lies within the lower and upper bounds of the interval.

Let T_1 and T_2 be two statistics such that,

$$P(T_1 > \theta) = \alpha_1 \qquad \dots (1)$$

and

$$P(T_2 < \theta) = \alpha_2 \qquad \dots (2)$$

where α_1 and α_2 are constants independent of θ . Equations (1) & (2) can be combined to give

$$P(T_1 < \theta < T_2) = 1 - \alpha \qquad \dots (3)$$

where $\alpha = \alpha_1 + \alpha_2$

• <u>Example</u>: If we take a large sample from a normal population with mean μ and standard deviation σ , then $100(1 - \alpha)\%$ confidence interval for a population mean at $\alpha = 0.05$ is

$$Z = \frac{\bar{x} - \mu}{\sigma / \sqrt{n}} \sim N(0, 1)$$

In general, confidence interval for μ can be constructed by using the following normal probability approach,

$$P\left(Z_{\alpha/2} \leq Z \leq Z_{1-\alpha/2}\right) = 1 - \alpha,$$

Where $P(Z < Z_{\alpha/2}) = \alpha/2$.

In particular case, $\alpha = 0.05$ then $Z_{\alpha/2} = -Z_{1-\frac{\alpha}{2}} = -1.96$ implies, $P(-1.96 \le Z \le 1.96) = 0.95 \dots$ (From Normal Probability Tables)

$$\Rightarrow P\left(-1.96 \le \frac{\bar{x}-\mu}{\sigma/\sqrt{n}} \le 1.96\right) = 0.95$$
$$\Rightarrow P\left(\bar{x}-1.96 \ \frac{\sigma}{\sqrt{n}} \le \mu \le \bar{x} + 1.96 \ \frac{\sigma}{\sqrt{n}}\right) = 0.95$$

Thus, $\bar{x} \pm 1.96 \frac{\sigma}{\sqrt{n}}$ are 95% confidence limits for the unknown parameter μ , the population mean and interval $\left(\bar{x} - 1.96 \frac{\sigma}{\sqrt{n}}, \bar{x} + 1.96 \frac{\sigma}{\sqrt{n}}\right)$ is called the 95% confidence interval.

• **Remark** For a small sample case with unknown variance, to construct a confidence interval for a population mean, use students-t distribution.

7.2.1 Probable error:

- Probable error defines the half-range of an interval about a mean for the distribution, such that half of the values from the distribution will lie within the interval and half outside.
- Thus for a symmetric distribution it is equivalent to half the interquartile range, or the median absolute deviation.
- For a normal distribution probable error is 0.6745σ

7.3 EXERCISE

- 1. What do you understand by point estimation?
- 2. Explain the terms statistic and its sampling distribution.
- 3. Define estimator and estimate.
- 4. Write a note on unbiasedness, consistency, the efficiency of an estimator. Also, define sufficient statistics.
- 5. Define MVUE and UMVUE.
- 6. Define Neyman factorization criteria.
- 7. Construct a confidence interval for a population mean where population variance is known.

7.4 SUMMARY

- In this chapter we studied basic definitions related to point estimation of population parameters.
- The good estimator must satisfy conditions of Unbiasedness, Consistency, Efficiency and Sufficiency.
- Uniformly Minimum variance unbiased estimators popularly used techniques to estimate population parameters.
- In this chapter we studied confidence interval estimation of the population parameter.

7.5 REFERENCES

- 1. Gupta S. C. and Kapoor V. K., 2011, Fundamentals of Mathematical Statistics, 11th Ed, Sultan and Chand.
- 2. Rohatgi V. K, 1939, Introduction to Probability and Statistics, Wiley
- 3. Murray R. Spiegel, Larry J. Stephens, STATISTICS, 4th Ed, McGRAW HILL INTERNATIONAL.
- 4. J.N. KAPUR and H.C. SAXENA, 2005, MATHEMATICAL STATISTICS, 12th rev, S.Chand.
- 5. Agrawal B. L, 2003, Programmed Statistics, 2nd Ed, New Age International.

STATISTICAL DECISION THEORY

8

Unit Structure

- 8.0 Objectives
- 8.1 Basic definitions
 - 8.1.1 Statistical decision
 - 8.1.2 Hypothesis
 - 8.1.3 Null hypothesis
 - 8.1.4 Alternative hypothesis
 - 8.1.5 Simple and composite hypothesis
 - 8.1.6 Errors in the test of significance
 - 8.1.7 Critical region
 - 8.1.8 Level of significance
 - 8.1.9 Test statistic
 - 8.1.10 One-tailed and two-tailed tests
 - 8.1.11 Central limit theorem
 - 8.1.12 Critical value
 - 8.1.13 P-value
 - 8.1.14 Power of the test
 - 8.1.15 Procedure for test a hypothesis
- 8.2 Large sample test
 - 8.2.1 One sample Z-test for a mean
 - 8.2.2 Two sample Z-test for the difference of two mean
 - 8.2.3 Test for a testing population proportion
 - 8.2.4 Test for testing equality of population proportion
- 8.3 Small sample test
 - 8.3.1 A t-test for testing a population mean
 - 8.3.2 Two sample t-test for the difference of two mean
 - 8.3.3 Paired t-test
- 8.4 Control chart
 - 8.4.1 A lot acceptance sampling plan
 - 8.4.2 OC curve
 - 8.4.3 Control chart
 - 8.4.4 Variable control chart
 - 8.4.5 Attribute control chart
- 8.5 Summary

8.0 OBJECTIVES

In this chapter, students can learn,

- Basic definitions related to the testing of hypothesis/decision making
- Decision making through Testing of hypothesis
- Large and small sample tests
- Decision Making through statistical control chart

8.1 BASIC DEFINITIONS

8.1.1 Statistical decision:

• The decisions are made based on observations of a phenomenon that carry out probabilistic laws that are not completely known.

8.1.2 Hypothesis:

- A definite statement about the population parameter is called a hypothesis. A hypothesis is a claim to be tested.
- *Example*: A particular scooter gives an average of 50 km per litre.

8.1.3 Null hypothesis:

- A hypothesis having no difference is called the null hypothesis.
- **Example:** The population mean is μ_0 the hypothesis will be $H_0: \mu = \mu_0$.

8.1.4 Alternative hypothesis:

- A hypothesis that is accepted in the case H_0 is rejected is called the alternative hypothesis and usually denoted by H_1 . It is exactly opposite to H_0 .
- <u>Example</u>: If $H_0: \mu = \mu_0$ i.e., the population has a specified mean μ_0 , then the alternative hypothesis could be;

i.	$H_1: \mu \neq \mu_0 \ (\mu > \mu_0 \ or \ \mu < \mu_0)$	Two-tailed alternative
ii.	$H_1: \mu > \mu_0$	Right tailed alternative
iii.	$H_1: \mu < \mu_0$	Left tailed alternative

8.1.5 Simple and composite hypothesis:

- A statistical hypothesis that completely specifies the population parameter is called a simple hypothesis, and the hypothesis that does not specify the population parameter is called a composite hypothesis.
- <u>Example</u>: If $x_1, x_2, ..., x_n$ is a random sample from normal with mean μ and variance σ^2 then $H_0: \mu = \mu_0$ and $\sigma^2 = \sigma_1^2$ is a simple

hypothesis. The following fully not specified hypotheses is called a composite hypothesis.

1) $H_1: \mu \neq \mu_0$ 2) $H_0: \sigma^2 \neq \sigma_0^2$ 3) $H_0: \mu = \mu_0 \text{ and } \sigma^2 > \sigma_0^2$ etc.

8.1.6 Errors in the test of significance:

- The main objective in the sampling theory is to draw a valid inference about the population parameters based on sample results. In practice, we decide to accept or reject the lot after examining a sample drawn from it. In sampling theory, we are liable to commit two types of errors: a rejection of a good lot and acceptance of a bad lot.
- *i. Type-I error*: Rejecting H_0 when H_0 is true.
- *ii.* **Type-II error**: Accepting H_0 when it is false (Accepting H_0 when H_1 is true).
- iii. Size of Type-I and Type-II errors

 $P [Reject H_0 when it is true] = P [Reject H_0 | H_0] =$

- $P [Reject \ a \ lot \ when \ it \ is \ good] = \alpha$
- $P [Accept H_0 when it is wrong] = P [Accept H_0 | H_1] =$
- $P [Accept a lot when it is bad] = \beta$

In the above probabilities, $\alpha \& \beta$ are called the Type-I & Type-II errors, respectively.

Actual Situation	Decision		
Actual Situation	Reject H_0	Accept H_0	
H_0 is true	Type-I Error	Correct Decision	
H_0 is false	Correct Decision	Type-II Error	

• The four types of decisions are shown in the table as follows.

8.1.7 Critical region:

• A region in sample space S which amounts to a rejection of H_0 is called a critical region or rejection region of H_0 .

8.1.8 Level of significance:

• The probability ' α ' is that the value of the test statistic belongs to the critical region, known as 'level of significance'. That is the probability of the occurrence of the type I error is the level of significance. Usually, we use the level of significance of 5% or 1%. The level of significance is always fixed in advance before collecting the sample information.

8.1.9 Test statistic:

• A function of sample observations is used to test H_0 is called test statistic.

8.1.10 One-tailed and two-tailed tests:

- A function of any statistical hypothesis where the alternative hypothesis is one-tailed (right-tailed or left-tailed) is called a one-tailed test.
- *Example*: A test for testing the mean of a population

 $H_0: \mu = \mu_0$ Versus $H_1: \mu > \mu_0$ (Right Tailed) or $H_1: \mu < \mu_0$ (Left Tailed), is a one-tailed test. In the right-tailed test, the critical region lies entirely in the right tail of the sampling distribution of \bar{X} , while for the left tailed test, the critical region lies entirely in the left tail of the sampling distribution of \bar{X} .

• A test of any statistical hypothesis where the alternative hypothesis is two-tailed such as; $H_0: \mu = \mu_0$ Versus $H_1: \mu \neq \mu_0$ ($\mu > \mu_0$ or $\mu < \mu_0$) is known as a two-tailed test, and in such a case, the critical region is given by the portion of the area lying in both the tails (sides) of the probability curve of the test statistic \overline{X} .

8.1.11 Central limit theorem:

• In many cases, the exact probability distribution of the test statistics T cannot be obtained. The difficulty is overcome using the normal approximation. The probability distribution of standardized T is assumed to be N(0, 1) as the sample size $n \to \infty$ (i.e., n is sufficiently large). The corresponding theorem in support of the normal approximation is known as the central limit theorem.

Case-I: Parent population is Normal:

Let the random sample drawn from $N(\mu, \sigma^2)$. By the definition of a random sample, variates values $x_1, x_2, ..., x_n$ of the sample are independent and identically distributed as $N(\mu, \sigma^2)$ then the sample mean $(\overline{\mu})$

 (\bar{X}) is distributed normally with μ and variance σ^2/n i.e.,

 $\bar{X} \sim N \ (\mu, \sigma^2/n)$. The result shows how the precision of a sample mean

increases as the sample size increases and $Z = \frac{\bar{x} - \mu}{\sigma/\sqrt{n}} \sim N(0, 1)$, standard normal variate.

Case-II: Parent population is Non-Normal:

If the population from which the random sample is drawn has a nonnormal distribution with finite mean μ and finite standard deviation σ then the variate, by owing to central limit theorem,

$$Z = \frac{\bar{x} - \mu}{\sigma / \sqrt{n}} \sim N(0, 1) \quad \text{as } n \to \infty$$

8.1.12 Critical value:

• The value of the test statistic, which separates the critical (rejection) region and acceptance region, is called the *'critical value'*. It depends upon (1) The level of significance α used and (2) The alternative hypothesis, whether it is two-tailed or one-tailed.



8.1.13 P-value:

- Another approach for testing is to find out the 'p' value at which H_0 is significant. That is, to find the smallest level of significance, α at which H_0 is rejected. About the acceptance or rejection of H_0 , the experimenter can himself decide the level α by comparing it with the p-value. The criterion for this is that if the p-value is less than or equal to α , reject H_0 otherwise, accept H_0 .
- <u>Example</u>:



Fig.4: P-value for One-tailed Z-Test

8.1.14 Power of test:

• Power of Test = P [Rejecting H_0 given that H_1 is true] = P [Rejecting $H_0 | H_1$ is true] = 1 - P [Accepting $H_0 | H_1$ is true] = 1 - P [Type II Error] = $1 - \beta$

8.1.15 The procedure of testing of hypothesis:

In any testing of a hypothesis, it is a stepwise procedure that leads to rejection or acceptance of the null hypothesis based on samples drawn from the population. The various steps in testing a statistical hypothesis are as follows;

- i. Set up the null hypothesis H_0 .
- ii. Set up the alternative hypothesis H_1 , this will enable us to decide whether we have to use a one-tailed (right or left) test or a two-tailed test.
- iii. Choose an appropriate level of significance (α), i.e., α is fixed in advance.
- iv. Choose the appropriate test statistic Z or T and find its value under the null hypothesis H_0 .
- v. Determine the critical values and critical region corresponding to the level of significance and the alternative hypothesis (Z_{α} for one-tailed H_1 or $Z_{\alpha/2}$ for two-tailed H_1).
- vi. Decision rule: We compare the calculated value of Z, with the tabulated value of Z_{α} if H_1 is one-tailed & compare |Z| with $Z_{\alpha/2}$ (For symmetrical distribution). If the calculated value is greater than the tabulated value, then we reject H_0 at α % level of significance and conclude that there is a significant difference at α % level of significance otherwise accept H_0 at α % significance level and conclude that there is no significant difference at α % level of significance.

8.2 LARGE SAMPLE TESTS

8.2.1 Test for testing of a population mean:

- Let consider $x = \{x_1, x_2, ..., x_n\}$ be a random sample of size n taken from a normally distributed population having population mean μ and population variances σ^2 respectively.
- We have to test the hypothesis,

 $H_0: \mu = \mu_0$

Against

$$H_1: \mu \neq \mu_0 \text{ OR}$$
$$H_1: \mu > \mu_0 \text{ OR}$$

- $H_1: \mu < \mu_0$
- Under H_0 , the test statistic is,

$$Z = \frac{\bar{x} - \mu_0}{\sigma / \sqrt{n}} \sim N \ (0, 1),$$

where n is the sample size, \bar{x} is the sample mean, σ population standard deviation.

Let Z_{α} be the critical value at α level of significance. We compare the calculated value of Z, with the tabulated value of Z_{α} . Where $P(Z \leq Z_{\alpha}) = \alpha$.

- H_1 is two-tailed $(\mu \neq \mu_0)$ $|Z| \ge Z_{\alpha/2}$, then reject H₀.
- H_1 is one-tailed $(\mu > \mu_0)$ $Z \ge Z_{1-\alpha}$, then reject H₀.
- H_1 is one-tailed ($\mu < \mu_0$)
- $Z \leq Z_{\alpha}$, then reject H₀.

8.2.2 Test for the testing difference of the two population mean:

- Let consider $x_1 = \{x_{11}, x_{12}, ..., x_{1n_1}\}$ and $x_2 = \{x_{21}, x_{22}, ..., x_{2n_2}\}$ be two independent normally distributed samples having population means μ_1 and μ_2 and population variances σ_1^2 and σ_2^2 respectively.
- \bar{x}_1 and \bar{x}_2 are the sample means.
- We have to test the hypothesis,

 $H_0: \mu_1 - \mu_2 = \mu_0$ Against

$$H_{1}: \mu_{1} \neq \mu_{2} \text{ OR}$$
$$H_{1}: \mu_{1} - \mu_{2} > \mu_{0} \text{ OR}$$
$$H_{1}: \mu_{1} - \mu_{2} < \mu_{0}$$

Under H_0 , the test statistic is,

$$Z = \frac{(\bar{x}_1 - \bar{x}_2) - \mu_0}{\sqrt{\left[\frac{\sigma_1^2}{n_1}\right] + \left[\frac{\sigma_2^2}{n_2}\right]}} \sim N \ (0, 1).$$

Let Z_{α} be the critical value at α level of significance. We compare the calculated value of Z, with the tabulated value of Z_{α} . Where $P(Z \leq Z_{\alpha}) = \alpha$.

• H_1 is two-tailed $(\mu_1 \neq \mu_2)$ $|Z| \ge Z_{\alpha/2}$, then reject H₀.

- H_1 is one-tailed $(\mu_1 \mu_2 > \mu_0)$
- $Z \ge Z_{1-\alpha}$, then reject H₀.
- H_1 is one-tailed $(\mu_1 \mu_2 < \mu_0)$
- $Z \leq Z_{\alpha}$, then reject H₀.

8.2.4 Test for testing of the population proportion:

• We have to test the hypothesis,

$$H_0: P = P_0$$

Against

$$H_1: P \neq P_0 \text{ OR}$$
$$H_1: P > P_0 \text{ OR}$$
$$H_1: P < P_0$$

• Under H_0 , the test statistic is,

$$Z = \frac{p - P_0}{\sqrt{\frac{P_0 Q_0}{n}}} \sim N \ (0, 1),$$

where n is the sample size, p is the sample proportion.

Let Z_{α} be the critical value at α level of significance. We compare the calculated value of Z, with the tabulated value of Z_{α} . Where $P(Z \le Z_{\alpha}) = \alpha$.

- H_1 is two-tailed $(P \neq P_0)$ $|Z| \ge Z_{\alpha/2}$, then reject H₀.
- H_1 is one-tailed $(P > P_0)$ $Z \ge Z_{1-\alpha}$, then reject H_0 .
- H_1 is one-tailed ($P < P_0$) $Z \le Z_{\alpha}$, then reject H₀.

8.2.4 Test for testing equality of two population Proportion:

• We have to test the hypothesis,

$$H_0: P_1 = P_2$$

Against

$$H_1: P_1 \neq P_2 \text{ OR}$$
$$H_1: P_1 > P_2 \text{ OR}$$
$$H_1: P_1 < P_2$$

• Under H_0 , the test statistic is,

$$Z = \frac{(p_1 - p_2)}{\sqrt{\hat{P}\hat{Q}\left(\frac{1}{n_1} + \frac{1}{n_2}\right)}} \sim N(0, 1)$$
$$\hat{P} = \frac{n_1 p_1 + n_2 p_2}{n_1 + n_2} \quad \hat{Q} = 1 - \hat{P}$$

Let Z_{α} be the critical value at α level of significance. We compare the calculated value of Z, with the tabulated value of Z_{α} . Where $P(Z \leq Z_{\alpha}) = \alpha$.

• H_1 is two-tailed $(P_1 \neq P_2)$

 $|Z| \ge Z_{\alpha/2}$, then reject H₀.

- H_1 is one-tailed $(P_1 > P_2)$ $Z \ge Z_{1-\alpha}$, then reject H₀.
- H_1 is one-tailed $(P_1 < P_2)$ $Z \le Z_{\alpha}$, then reject H₀.

8.3 SMALL SAMPLE TESTS

8.3.1 Test for the population mean:

- Let consider $x = \{x_1, x_2, ..., x_n\}$ be a random sample of size n (n small) taken from a normally distributed population having population mean μ and population variances σ^2 (mean and variances are unknown) respectively.
- We have to test the hypothesis,

Against

$$H_1: \mu \neq \mu_0$$
 OR
 $H_1: \mu > \mu_0$ OR

 $H_0: \mu = \mu_0$

$$H_1: \mu < \mu_0$$

• Under H_0 , the test statistic is,

$$t = \frac{\bar{x} - \mu}{s/\sqrt{n}} \sim t_{n-1} d. f.$$

Where n is the sample size, \bar{x} is the sample mean, s sample standard deviation.

Let $t_{(n-1,\alpha)}$ be the critical value based on students t-distribution at n-1 degrees of freedom and α level of significance. We compare the

calculated value of t, with the tabulated value of $t_{(n-1,\alpha)}$. Where $P(T \le t_{(n-1,\alpha)}) = \alpha$.

- H_1 is two-tailed $(\mu \neq \mu_0)$ $|t| \ge t_{(n-1,\alpha/2)}$, then reject H₀.
- H_1 is one-tailed $(\mu > \mu_0)$
- $t \ge t_{(n-1,1-\alpha)}$, then reject H₀.
- H_1 is one-tailed ($\mu < \mu_0$)
- $t \leq t_{(n-1,\alpha)}$, then reject H₀.

8.3.2 Test for the difference of two population means (two samples):

- Let consider $x = \{x_1, x_2, ..., x_{n_1}\}$ and $y = \{y_1, y_2, ..., y_{n_2}\}$ be two independent normally distributed samples having unknown population means μ_1 and μ_2 and unknown population variances σ_1^2 and σ_2^2 respectively.
- \bar{x} and \bar{y} are the sample arithmetic means.
- We have to test the hypothesis,

 $H_0: \mu_1 - \mu_2 = \mu_0$

Against

 $H_{1}: \mu_{1} \neq \mu_{2} \text{ OR}$ $H_{1}: \mu_{1} - \mu_{2} > \mu_{0} \text{ OR}$ $H_{1}: \mu_{1} - \mu_{2} < \mu_{0}$

• Under H_0 , the test statistic is,

$$t = \frac{\bar{x} - \bar{y}}{s / \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}} \sim t_{n_1 + n_2 - 2} d. f. \quad \text{where } s^2 = \frac{(n_1 - 1)s_x^2 + (n_2 - 1)s_y^2}{n_1 + n_2 - 2}.$$

Let $t_{(n_1+n_2-2,\alpha)}$ be the critical value based on students t-distribution at $n_1 + n_2 - 2$ degrees of freedom and α level of significance. We compare the calculated value of t, with the tabulated value of $t_{(n_1+n_2-2,\alpha)}$. Where $P(T \le t_{(n_1+n_2-2,\alpha)}) = \alpha$.

- H_1 is two-tailed $(\mu \neq \mu_0)$ $|t| \ge t_{(n_1+n_2-2,\alpha/2)}$, then reject H₀.
- H_1 is one-tailed $(\mu > \mu_0)$ $t \ge t_{(n_1+n_2-2,1-\alpha)}$, then reject H₀.
- H_1 is one-tailed $(\mu < \mu_0)$
- $t \leq t_{(n_1+n_2-2,\alpha)}$, then reject H₀.

8.3.3 Paired t-test for difference of mean:

- Let consider $\{x_i, y_i\}; i = 1, 2, ..., n$ be n-pair dependent normally distributed samples having unknown population means μ_1 and μ_2 and unknown population variances σ_1^2 and σ_2^2 respectively.
- We have to test the hypothesis,

$$H_0: \mu_1 - \mu_2 = 0$$

Against

$$H_1: \mu_1 - \mu_2 \neq 0 \text{ OR}$$

 $H_1: \mu_1 - \mu_2 > 0 \text{ OR}$
 $H_1: \mu_1 - \mu_2 < 0$

• Under H_0 , the test statistic is,

 $t = \frac{\bar{X} - \bar{Y}}{s/\sqrt{n}} \sim t_{n-1} d. f.$ where $d = \bar{X} - \bar{Y} \& s^2 = \frac{1}{n-1} \sum (d_i - \bar{d})^2$

Let $t_{(n-1,\alpha)}$ be the critical value based on students t-distribution at n-1 degrees of freedom and α level of significance. We compare the calculated value of *t*, with the tabulated value of $t_{(n-1,\alpha)}$. Where $P(T \le t_{(n-1,\alpha)}) = \alpha$.

• H_1 is two-tailed $(\mu \neq \mu_0)$

 $|t| \ge t_{(n-1,\alpha/2)}$, then reject H₀.

• H_1 is one-tailed $(\mu > \mu_0)$

 $t \ge t_{(n-1,1-\alpha)}$, then reject H₀.

• H_1 is one-tailed ($\mu < \mu_0$)

 $t \leq t_{(n-1,\alpha)}$, then reject H₀.

8.4 STATISTICAL CONTROL CARTS

8.4.1 A lot acceptance sampling plan (LASP):

• A lot acceptance sampling plan (LASP) is a sampling scheme and a set of rules for making decisions. The decision, based on counting the number of defectives in a sample, can be to accept the lot, reject the lot, or even, for multiple or sequential sampling schemes, to take another sample and then repeat the decision process.

8.4.2 OC curve:

• Operating Characteristic (OC) Curve plots the probability of accepting the lot (Y-axis) versus the lot fraction or percent defectives (X-axis). The OC curve is the primary tool for displaying and investigating the properties of a LASP.

8.4.3 Control charts:

• Control charts are a statistical process control tool used to determine if a manufacturing or business process is in a state of control. It is more appropriate to say that the control charts are the graphical device for Statistical Process Monitoring (SPM). Traditional control charts are mostly designed to monitor process parameters when an underlying form of the process distributions are known.

8.4.4 Variable control chart (\overline{X} Chart):

• Dr Walter A. Shewhart proposed a general model for control charts in 1920. Let w be a sample statistic that measures some continuously varying quality characteristic of interest (e.g., thickness), and suppose that the mean of w is μ_w , with a standard deviation of σ_w . Then the centre line, the Upper Control Limit (UCL), and the Lower Control Limit (LCL) are:

$$UCL = \mu_w + k\sigma_w$$

Center Line = μ_w
 $LCL = \mu_w + k\sigma_w$

where k is the distance of the control limits from the centre line, expressed in terms of standard deviation units. When k is set to 3, we speak of 3 - sigma control charts. Historically, k = 3 has become an accepted standard in the industry. The centerline is the process mean, which in general is unknown. We replace it with a target or the average of all the data. The quantity that we plot is the sample average, \overline{X} The chart is called the \overline{X} chart.

We also have to deal with the fact that σ is, in general, unknown. Here we replace σw with a given standard value, or we estimate it by a function of the average standard deviation.

8.4.5 Attributes control charts:

The Shewhart control chart plots quality characteristics that can be measured and expressed numerically. We measure weight, height, position, thickness, etc. If we cannot represent a particular quality characteristic numerically, or if it is impractical to do so, we then often resort to using a quality characteristic to sort or classify an item that is inspected into one of two "buckets".

An example of a common quality characteristic classification would be designating units as "conforming units" or "nonconforming units". Another quality characteristic criteria would be sorting units into "non-defective" and "defective" categories. Quality characteristics of that type are called **attributes.**

• Control charts dealing with the number of defects or nonconformities are called **c charts** (for the count).

- Control charts dealing with the proportion or fraction of defective products are called **p charts** (for proportion).
- There is another chart that handles defects per unit, called the **u chart** (for the unit). This applies when we wish to work with the average number of nonconformities per unit of product.

8.6 SUMMARY

Students can get an idea about the testing of a hypothesis and make decisions about parameters of interest. In this chapter, we briefly studied various large/small One-sample and two-sample tests for mean and proportion along with variable and attribute control charts are also discussed.

8.5 EXERCISE

- 1. Explain the term hypothesis and its types.
- 2. Define Type-I and Type-II errors.
- 3. Explain the terms level of significance, p-value and power of a test.
- 4. Write down the stepwise procedure of testing of hypothesis.
- 5. Write down the procedure for testing the equality of two population proportions.
- 6. Write down the procedure to test the specified population mean, in the case of a small sample.
- 7. Explain in detail paired t-Test for difference Mean.

8.7 REFERENCES

- Gupta S. C. and Kapoor V. K., 2011, Fundamentals of Mathematical Statistics, 11th Ed, Sultan and Chand.
- Rohatgi V. K, 1939, Introduction to Probability and Statistics, Wiley
- Murray R. Spiegel, Larry J. Stephens, STATISTICS, 4th Ed, McGRAW HILL INTERNATIONAL.
- J.N. KAPUR and H.C. SAXENA, 2005, MATHEMATICAL STATISTICS, 12th rev, S.Chand.
- Agrawal B. L, 2003, Programmed Statistics, 2nd Ed, New Age International.
- Kanji G. K., 2006, 100 Statistical tests, 3rd Ed, SAGE Publication.

STATISTICS IN R

9

Unit Structure

- 9.0 Objectives
- 9.1 Descriptive statistics in R
- 9.2 Normal distribution
- 9.3 Binomial distribution
- 9.4 Frequency distribution
- 9.5 Data import and export
- 9.6 Summary
- 9.7 Exercise
- 9.8 References

9.0 OBJECTIVES

- Use of R-software to find basic statistical measures
- Use of R-software to simulate Normal Distributions
- Use of R-software to simulate Binomial Distributions
- Data import and Export in R.

9.1 DESCRIPTIVE STATISTICS IN R-SOFTWARE

G						
Sr.	Descriptiv	Syntax in R	Output			
No	e					
•	Statistics					
1.	Observatio	$x \le c(1,4,7,12,19,15,21,20)$	Х			
	n Vector		[1] 1 4 7 12 19 15			
	(c-	•	21 20			
	function)					
2.	Arithmetic	AM<-mean(x)	AM			
	mean		[1] 12.375			
3	Mode	# Create the function	Mode [1] 2			
5.	widde	# Create the function.				
		gettilode $<-$ function(x) {				
		uniqv <- unique(x)				
		uniqu[which may(tabulata(ma				
		tcn(x, uniqv)))]				
		}				
		V <-				
		c(2,1,2,3,1,2,3,4,1,5,5,3,2,3);				
		mode<-getmode(v)				
4.	median	Med=median(x)	Med [1] 13.5			
9.2 NORMAL DISTRIBUTION

R has four inbuilt functions to generate normal distribution. They are described below.

Sr.	Function	Syntax	Description
no.	Name		_
1.	Density function	dnorm(x, mean, sd)	This function gives the height of the probability distribution at each point for a given mean and standard deviation.
2.	Cumulative Probability	pnorm(x, mean, sd)	This function gives the probability of a normally distributed random number less than the value of a given number. It is also called the "Cumulative Distribution Function".
3.	Inverse function	qnorm(p, mean, sd)	This function takes the probability value and gives a number whose cumulative value matches the probability value.
4.	Random Number	rnorm(n, mean, sd)	This function is used to generate random numbers whose distribution is normal. It takes the sample size as input and generates that many random numbers.

Following is the description of the parameters used in the above functions:

x: is a vector of numbers.

- p: is a vector of probabilities.
- n: is a number of observations(sample size).

mean: is the mean value of the sample data. Its default value is zero.

sd: is the standard deviation. Its default value is 1.

9.3 BINOMIAL DISTRIBUTION

R has four in-built functions to generate binomial distribution. They are described below.

Sr. no.	Function Name	Syntax	Description			
1.	Probability	dbinom(x, size, prob)	This function gives the			

			probability density
			distribution at each point.
2.	Cumulative Probability	pbinom(x, size, prob)	This function gives the cumulative probability of an event. It is a single value representing the probability.
3.	Inverse function	qbinom(p, size, prob)	This function takes the probability value and gives a number whose cumulative value matches the probability value.
4.	Random Number	rbinom(n, size, prob)	This function generates required number of random values of given probability from a given sample.

Following is the description of the parameters used -

- x: is a vector of numbers.
- p: is a vector of probabilities.
- n: is a number of observations.

Size: is the number of trials.

prob: is the probability of success of each trial.

9.4 FREQUENCY DISTRIBUTION IN R

- Table function in R *-table()*, performs categorical tabulation of data with the variable and its frequency. *table()* function is also helpful in creating Frequency tables with the condition and cross-tabulation
- Syntax

 $x \le c(1,2,3,2,4,2,5,4,6,7,8,9)$

freq <- data.frame(table(x))</pre>

• Output

> freq	Х	Freq
1	1	1
2	2	3
3	3	1
4	4	2
5	5	1
6	6	1
7	7	1
8	8	1
9	9	1

9.5 DATA IMPORT AND EXPORT

i. Data Importing:

The sample data is frequently observed in Excel format and needs to be imported into R before use. For this, we can use the function read.xls from the gdata package. It reads from an Excel spreadsheet and returns a data frame.

library(gdata) # load gdata package > help(read.xls) # documentation > mydata = read.xls("mydata.xls") # read from the first sheet

ii. Data export:

There are numerous methods for exporting R objects into other formats.

A tab delimited tex file write.table(mydata, "c:/mydata.txt", sep="\t")

MS-Excel Spread sheet

library(xlsx) write.xlsx(mydata, "c:/mydata.xlsx")

9.7 SUMMARY

R is a programming language and free software environment for statistical computing and graphics supported by the R Core Team and the R Foundation for Statistical Computing. It is widely used for data analysis purposes in analytical industries. In this chapter, we studied various descriptive measures in R, probabilities, quantiles and random number generation syntax of the normal and binomial distribution.

9.6 EXERCISE

1. Write down the R-command for the arithmetic mean and compute for the given data.

1, 4, 7, 12, 19, 15, 21 and 20.

- 2. Generate 10 random numbers using R-command from a normal distribution with mean zero and standard deviation one.
- 3. Calculate cumulative distribution function at point zero for normal distribution with mean zero and standard deviation one.
- 4. Compute probability density function at 1, 2, 3, 4, 5 for normal distribution with mean zero and standard deviation one.
- 5. Compute quantiles at 0.1, 0.2, 0.3, 0.4, 0.5 for binomial distribution with n=5 and p=0.7.
- 6. Generate 10 random numbers using R-command from a binomial distribution with n=5 and p=0.5.

7. Explain frequency distribution in R.

9.8 REFERENCES

- Gupta S. C. and Kapoor V. K., 2011, Fundamentals of Mathematical • Statistics, 11th Ed, Sultan and Chand.
- R.B. Patil, H.J. Dand and R. Bhavsar, 2017, A Practical Approach • using R, 1st ed, SPD.
- https://www.tutorialspoint.com/r/r normal distribution.htm •
- https://cran.r-project.org/bin/windows/base/ •

UNIT IV

10

SMALL SAMPLING THEORY

Unit Structure

- 10.0 Objectives
- 10.1 Introduction
- 10.2 Student's t distribution
- 10.3 Graph of *t*-distribution
- 10.4 Critical values of t
- 10.5 Application of t-distribution
- 10.6 Test of Hypothesis and Significance
- 10.7 Confidence Interval
- 10.8 t-Test for Difference of Means
- 10.9 Degrees of Freedom
- 10.10 The F-Distribution
- 10.11 Summary
- 10.12 Reference for further reading
- 10.13 Exercises
- 10.14 Solution to Exercises
- 10.15 Tables of t-distribution and F-distribution

10.0 OBJECTIVES

In this chapter we will study about the test suitable for small samples i.e sample size less than or equal to 30 for which the tests studied in previous chapters are not applicable. We will also study the test for equality of variance.

10.1 INTRODUCTION

The entire large sample theory was based on the application of "Normal Test". However if the sample size n is small, the distribution of the various statistics, e.g. $Z = \frac{\bar{x} - \mu}{(\frac{\sigma}{\sqrt{n}})} or Z = \frac{X - nP}{\sqrt{nPQ}}$ etc., are far from normality and as such

'normal test' cannot be applied if 'n' is small. In such cases exact sample tests, pioneered by W.S. Gosset (1908) who wrote under the pen name of Student, and later on developed and extended by Prof. R. A. Fisher (1926), are used. In the following sections we shall discuss i) t - test and ii) F-test

test.

The exact sample tests can, however, be applied to large samples though the converse is not true. In all exact sample tests, the basic assumption is that "the population(s) from which the sample(s) is(are) drawn is(are) normal, i.e., the parent population(s) is(are) normally distributed."

10.2 STUDENT'S T DISTRIBUTION

Let x_i (i = 1,2,...,n) be a random sample of size n from a normal population with mean μ and variance σ^2 . Then Student's t is defined by the statistic:

$$t = t_{cal} = \frac{\bar{x} - \mu}{\frac{s}{\sqrt{n}}} , s^2 = \frac{1}{n-1} \sum_{i=1}^{l=n} (x_i - \bar{x})^2 , \bar{x} = \frac{\sum x_i}{n}$$

where \bar{x} is the sample mean and s^2 is an unbiased estimate of the population variance σ^2 , and it follows Student's distribution with v = (n-1) degree of freedom with probability density function:

$$f(t) = \frac{1}{\sqrt{\nu}B\left(\frac{1}{2}, \frac{\nu}{2}\right)} \cdot \frac{1}{\left(1 + \frac{t^2}{\nu}\right)^{\frac{\nu+1}{2}}} , -\infty < t\infty$$

10.3 GRAPH OF t-DISTRIBUTION

The probability density function of t-distribution with n degrees of freedom is:

$$f(t) = C.\left(1 + \frac{t^2}{n}\right)^{-\frac{(n+1)}{2}}, \quad -\infty < t < \infty$$

Since f(t) is an even function, the probability curve is symmetric about the line t =0. As t increases, f(t) decreases rapidly and tends to zero as $t \to \infty$, so that t-axis is an asymptote to the curve. We know that

$$\mu_2 = \frac{n}{n-2}$$
, $n > 2$; $\beta_2 = \frac{3(n-2)}{n-4}$, $n > 4$

Hence for $n \ge 2$, $\mu_2 \ge 1$ i.e., the variance of t-distribution is greater than that of standard normal distribution and for $n \ge 4$, $\beta_2 \ge 3$ and thus t-distribution is more flat on the top than the normal curve. In fact, for small n, we have

$$P(|t| \ge t_0) > -P(|Z| \ge t_0), \qquad Z \sim N(0,1)$$

i.e, the tails of the t-distribution have a greater probability (area) than the tails of standard normal distribution. Moreover we can check that for large

n, t-distribution tends to standard normal distribution. Graph of tdistribution is given by the following diagram



10.4 CRITICAL VALUES OF t

The critical values of t at level of significance α and degree of freedom v for two tailed test are given by the equation:



The values $t_v(\alpha)$ have been tabulated in table, for different values of $\alpha andv$ are given at the end of the chapter.

Since t-distribution is symmetric about t=0, we get

$$P(t < t_{\nu}(\alpha)) + P(t < -t_{\nu}(\alpha)) = \alpha \Rightarrow 2P((t > t_{\nu}(\alpha)) = \alpha$$
$$\Rightarrow P(t > t_{\nu}(\alpha)) = \frac{\alpha}{2} \therefore P(t > t_{\nu}(2\alpha)) = \alpha$$

 $t_{\nu}(2\alpha)$ (from the tables at the end of the chapter) gives the significant value of t for a single tail test(Right tail or Left tail since the distribution is symmetrical), at level of significance $\alpha \& \nu$ degree of freedom.

Hence the significant values of t at level of significance ' α ' for a single tailed test can be obtained from those of two tailed test by looking the values at level of significance 2α . For example

 $t_8(0.05)$ for single tail test = $t_8(0.1)$ for two tail test = 1.86 $t_{15}(0.01)$ for single tail test = $t_{15}(0.02)$ for two tail test = 2.602

10.5 APPLICATION OF t-DISTRIBUTION

The t-distribution has a whole number of applications in Statistics, some of which are given here below:

- i) to test if the sample mean (\bar{x}) differs significantly from the hypothetical value μ of the population mean.
- ii) to test the significance of the difference between two sample means.
- iii) to test the significance of an observed sample correlation coefficient and sample regression co efficient.
- iv) to test the significance of observed partial correlation coefficient.

10.6 TEST OF HYPOTHESIS AND SIGNIFICANCE

Suppose we want to test:

i) if a random sample $x_i(i=1,2,...,n)$ of size n has been drawn from a normal population with a specified mean say μ_0 orif the sample mean differs significantly from the hypothetical value of μ_0 of the population mean.

i.e $H_0: \mu = \mu_0$, $H_1: \mu \neq \mu_0 or\mu > \mu_0 or\mu < \mu_0$ where H_0 is the null hypothesis and H_1 is the alternative hypothesis

ii) Calculate

$$t = \frac{\bar{x} - \mu}{\frac{S}{\sqrt{n}}}, S^2 = \frac{1}{n - 1} \sum_{i=1}^{i=n} (x_i - \bar{x})^2$$

- iii) degree of freedom df = v = n-1
- iv) from the table calculate, $t_v(\alpha)$
- v) Conclusion: Reject H_0 if calculated |t| > tabulated t and Do not reject H_0 if calculated $|t| \le$ tabulated t.

Remark: We know, the sample variance:

$$s^{2} = \frac{1}{n} \sum (x_{i} - \bar{x})^{2} \Rightarrow s^{2} = \frac{1}{n} (n - 1)S^{2} \Rightarrow \frac{s^{2}}{n - 1} = \frac{S^{2}}{n}$$

Hence for numerical problems, the test statistic t stated above will become

$$t = \frac{\bar{x} - \mu}{\frac{S}{\sqrt{n}}} = \frac{\bar{x} - \mu}{\frac{S}{\sqrt{n-1}}}$$

Eg: 1) A machinist is making engine parts with axle diameter of 0.7 inch. A random sample of 10 parts show a mean diameter of 0.742 inch with a standard deviation of 0.04 inch. Compute the statistic you would use to test whether the work is meeting the specifications and state the conclusion.

Solution:

i)
$$H_0: \mu = 0.7, H_1: \mu \neq 0.7$$

- ii) $\bar{x} = 0.742, s = 0.04, n = 10,$ $t_{cal} = \frac{\bar{x} - \mu}{\frac{s}{\sqrt{n-1}}} = \frac{0.742 - 0.7}{\frac{0.04}{\sqrt{9}}} = \frac{0.042 \times \sqrt{9}}{0.04} = 3.15$
- iii) degree of freedom i.e v = n-1, v = 9
- iv) $t_v(\alpha) = t_9(0.05) = t_9(0.025) = 2.262$
- v) conclusion $t_{cal} = 3.15 > t_9(0.05) = 2.262$ Since calculated 't' is greater than the tabulated 't' we reject the null hypothesis H₀ i.e the product is not conforming to specifications.

Eg: 2) The mean weakly sales of soap bars in departmental stores was 146.3 bars per store. After an advertising campaign the mean weekly sales in 22 stores for a typical week increased to 153.7 and showed a standard deviation of 17.2. Was the advertising campaign successful?

Solution:

i) $H_0: \mu = 146.3, H_1: \mu > 146.3$

ii)
$$\bar{x} = 153.7, s = 17.2, n = 22$$

 $t_{cal} = \frac{\bar{x} - \mu}{\frac{s}{\sqrt{n-1}}} = \frac{153.7 - 146.3}{\frac{17.2}{\sqrt{21}}} = \frac{7.4 \times \sqrt{21}}{17.2} = 1.97$

- iii) degree of freedom f i.e v = n-1, v = 21
- iv) iv) $t_v(\alpha) = t_{21}(0.05) = t_{21}(0.025) = 1.72$
- v) conclusion $t_{cal} = 1.97 > t_{21}(0.05) = 1.72$ Since calculated 't' is greater than the tabulated 't' we reject the null hypothesis H₀ i.e the advertising campaign was successful in promoting the sales.

Eg: 3) A random sample of 10 boys had the following I.Q.'s is 70, 120, 110, 101, 88, 83, 95, 98, 107, 100. Do these data support the assumption of a population mean I.Q of 100?

Solution: First we calculate find the value of $\bar{x} \& S^2$.

Here $n =$	$10, \Sigma x =$	= 972	$\therefore \bar{x} =$	$=\frac{\Sigma x}{m}=$	$\frac{972}{10} =$	97.2
				n	10	

х	70	120	110	101	88	83	95	98	107	100	Total
$x - \overline{x}$	-27.2	22.8	12.8	3.8	-9.2	-14.2	-2.2	0.8	9.8	2.8	
$(x-\overline{x})^2$	739.84	519.84	163.84	14.44	84.64	201.64	4.84	0.64	96.04	7.84	1833.6

$$S^{2} = \frac{1}{n-1} \sum (x_{i} - \bar{x})^{2} = \frac{1}{9} (1833.6) = 203.73 \Rightarrow S = \sqrt{203.73}$$
$$= 14.273$$

i) $H_0: \mu = 100, H_1: \mu \neq 100$

ii)
$$\bar{x} = 97.2, S = 203.73, n = 10$$

$$t_{cal} = \frac{|\bar{x} - \mu|}{\frac{s}{\sqrt{n}}} = \frac{|97.2 - 100|}{\frac{14.273}{\sqrt{10}}} = \frac{2.8 \times \sqrt{10}}{14.273} = 0.6203$$

iii) degree of freedom d f i.e v = n-1, v = 9

iv)
$$t_v(\alpha) = t_9(0.05) = t_9(0.025) = 2.262$$

v) conclusion
$$t_{cal} = 0.6203 < t_9(0.05) = 2.262$$

Since calculated 't' is less than the tabulated 't' we accept the null hypothesis H_0 i.e the data are consistent with the assumption of mean I.Q of 100 in the population.

10.7 CONFIDENCE INTERVAL

As done with normal distribution in earlier chapter, we can define 95%, 99% or other confidence intervals by using the table given at the end of this chapter. We can estimate within specified limits of confidence the population mean μ . In general confidence limits are given by the formula

$$\bar{x} \pm \frac{t_v(\alpha)S}{\sqrt{n}}$$

In specific the 95 % confidence interval is given by

$$\left(\bar{x} - \frac{t_v(0.05)S}{\sqrt{n}}, \bar{x} + \frac{t_v(0.05)S}{\sqrt{n}}\right)$$

and 99% confidence interval is given by

$$\left(\bar{x} - \frac{t_v(0.01)S}{\sqrt{n}}, \bar{x} + \frac{t_v(0.01)S}{\sqrt{n}}\right)$$

Eg: 1) A random sample of 16 values from a normal population showed a mean of 41.5 inches and the sum of squares of deviations from this mean = 135 sq. inches. Obtain the 95 % and 99 % confidence limits for population mean.

Solution: n = 16, \bar{x} = 41.5, $\sum (x - \bar{x})^2 = 135$ $\therefore S^2 = \frac{1}{n-1} \sum (x - \bar{x})^2 = \frac{1}{15} (135) = 9 \Rightarrow S = 3$

from the table of t-distribution, we get $t_{15}(0.05) = 2.131 \& t_v(0.01) = 2.94795\%$ confidence limits for population mean are given by

$$\bar{x} \pm \frac{t_{15}(0.05)S}{\sqrt{n}} = 41.5 \pm 2.131 \times \frac{3}{\sqrt{16}} = 41.5 \pm 2.131 \times 0.75$$

 $\Rightarrow 39.902 < \mu < 43.098$

99% confidence limits for population mean are given by

$$\bar{x} \pm \frac{t_{15}(0.01)S}{\sqrt{n}} = 41.5 \pm 2.947 \times \frac{3}{\sqrt{16}} = 41.5 \pm 2.947 \times 0.75$$
$$\Rightarrow 39.29 \le \mu \le 43.71$$

10.8 t-TEST FOR DIFFERENCE OF MEANS

Suppose we want to test if two independent samples x_i ($i = 1,2,3,...,n_1$) and y_j ($j = 1,2,3,...,n_2$) of size n_1 and n_2 have been drawn from two normal populations with means $\mu_x(\mu_1) \& \mu_y(\mu_2)$ respectively.

Under the null hypothesis (H₀) that the samples have been drawn from the normal populations with mean $\mu_x(\mu_1)\&\mu_y(\mu_2)$ and the under the assumption that the population variance are equal i.e $\sigma_x^2 = \sigma_y^2 = \sigma^2$, the statistic

$$t = \frac{\left(\bar{x} - \bar{y} - (\mu_x - \mu_y)\right)}{S\left(\sqrt{\frac{1}{n_1} + \frac{1}{n_2}}\right)} = \frac{\bar{x} - \bar{y} - d_0}{S\sqrt{\frac{1}{n_1} + \frac{1}{n_2}}} , d_0 = \mu_x - \mu_y$$

Where $\bar{x} = \frac{\sum x_i}{n_1}$, $\bar{y} = \frac{\sum y_j}{n_2}$ & $S^2 = \frac{1}{n_1 + n_2 - 2} \left[\sum (x_i - \bar{x})^2 + \sum (y_j - \bar{y})^2 \right]$ is

an unbiased estimate of the common population variance σ^2 , follows

Students t-distribution with $v = (n_1 - 1) + (n_2 - 1) = (n_1 + n_2 - 2)$ degrees of freedom.

Paired t-test for Difference of means:

Let us now consider the case when (i) the sample sizes are equal i.e. $n_1 = n_2 = n$ and (ii) the two samples are not independent but the sample observations are paired together, i.e., the pair of observations $(x_i, y_i), 1 \le i \le n$ corresponds to the same ith sample unit. The problem is to test if the sample means differ significantly or not.

For example, suppose we want to test the efficacy of a particular drug, say, for inducting sleep. Let x_i and y_i (i =1,2,...,n) be the readings, in hours of sleep, on the ith individual, before and after the drug is given respectively. Here instead of applying the difference of the means test discussed above in the same section we apply the paired t-test given below:

Here we consider the increments, $d_i = x_i - y_i$, $1 \le i \le n$

Under the null Hypothesis, H_0 that increments are due to fluctuations of sampling, i.e., the drug is not responsible for these increments, the statistic is

$$t = \frac{\bar{d}}{\frac{S}{\sqrt{n}}} = \frac{\bar{d} \times \sqrt{n}}{S}$$

where

$$\bar{d} = \frac{1}{n} \sum_{i=1}^{n} d_i \,\&S^2 = \frac{1}{n-1} \sum_{i=1}^{n} (d_i - \bar{d})^2 = \frac{1}{n-1} \left\{ \sum_{i=1}^{n} d_i^2 - \frac{(\sum_{i=1}^{n} d)^2}{n} \right\}$$

follows Student's t-distribution with (n - 1) degree of freedom.

Eg: 1) For a random sample of 10 pigs fed on diet A, the increases in weight in pounds in a certain period were: 10, 6, 16, 17, 13, 12, 8, 14, 15, 9. For another sample of 12 pigs, fed on Diet B, the increase in the same period were: 7, 13, 22, 15, 12, 14, 18, 8, 21, 23, 10, 17. Test whether diets A and B differ significantly as regards to their effect on increase in weight.

Solution:

i) Null Hypothesis, $H_0: \mu_x = \mu_y$, i.e., there is no significant difference between the mean increase in weight due to diets A and B.

Alternative	hypothesis,	$H_1: \mu_x \neq$	μ_{ν} (two tailed)
-------------	-------------	-------------------	--------------------------

Xi	$x_i - \bar{x}$	$(x_i - \bar{x})^2$	yj	$y_j - \overline{y}$	$(y_j - \overline{y})^2$
10	-2	4	7	-8	64
6	-6	36	13	-2	4
16	4	16	22	7	49

ii)____

	17	5	25	15	0	0
	13	1	1	12	-3	9
	12	0	0	14	-1	1
	8	-4	16	18	3	9
	14	2	4	8	-7	49
	15	3	9	21	6	36
	9	-3	9	23	8	64
				10	-5	25
				17	2	4
Total	120	0	120	180	0	314

$$\bar{x} = \frac{\sum x}{n_1} = \frac{120}{10} = 12 , \ \bar{y} = \frac{\sum y}{n_2} = \frac{180}{12} = 15$$

$$S^2 = \frac{1}{n_1 + n_2 - 2} \left[\sum (x_i - \bar{x})^2 + (y_j - \bar{y})^2 \right] = \frac{1}{20} [120 + 314]$$

$$S^2 = 21.7 \Rightarrow S = \sqrt{21.7} = 4.6583$$

$$t = \frac{\bar{x} - \bar{y} - d_0}{S\sqrt{\frac{1}{n_1} + \frac{1}{n_2}}} = \frac{12 - 15 - 0}{4.6583\sqrt{\frac{1}{10} + \frac{1}{12}}} = \frac{-3 \times \sqrt{120}}{4.6583 \times \sqrt{22}} = -\frac{32.8634}{21.8494}$$

$$\Rightarrow t = -1.5041$$

iii)
$$df = v = 10 + 12 - 2 = 20$$

iv)
$$t_{20}(0.05) = 2.086$$

v) Conclusion: $|t_{cal}| = 1.5041 < t_{20}(0.05) = 2.086$ Since calculated value of t is less than the tabulated value, the null hypothesis H₀ is accepted at 5% level of significance and we may conclude that the two diets do not differ significantly as regards their effect on increase in weight.

Eg: 2) The yields if two types 'Type A' and 'Type B' of grains in pounds per acre in 6 replications are given below. What comments would you make on the difference in the mean yield.

Replication	1	2	3	4	5	6
Yield of Type A	20.5	24.6	2306	29.98	30.37	23.83
Yield of Type B	24.86	26.39	28.19	30.75	29.98	22.04

Solution: i) $H_0: \mu_x = \mu_y, H_1: \mu_x \neq \mu_y$

ii)
$$t = \frac{\bar{d}}{\left(\frac{S}{\sqrt{n}}\right)}$$
, $\bar{d} = \sum_{i=1}^{n} d_i$, $d_i = x_i - y_i$, $v = n - 1$
$$S^2 = \frac{1}{n-1} \left\{ \sum d^2 - \frac{(\sum d)^2}{n} \right\}$$

replication	type A	type B	d	d^2
1	20.5	24.86	-4.36	
2	24.6	26.39	-1.79	
3	23.06	28.19	-5.13	
4	29.98	30.75	-0.77	
5	30.37	29.97	0.4	
6	23.83	22.04	1.79	
total			-9.86	52.4876

$$(iii)S^{2} = \frac{1}{5} \left[52.4876 - \frac{(-9.86)^{2}}{6} \right] = \frac{1}{5} \left[52.4876 - \frac{97.2196}{6} \right]$$
$$= \frac{36.2843}{5} = 7.2569$$
$$\therefore S = \sqrt{7.2569} = 2.6939$$
$$\bar{d} = -\frac{9.86}{6} = -1.6433, t = \frac{\bar{d}}{\frac{5}{\sqrt{n}}} = \frac{-1.6433 \times \sqrt{6}}{2.6939} = -1.4942$$

iv) $st_5(0.05) = 2.571$

v) Conclusion: $|t_{cal}| = 1.4942 < t_5(0.05) = 2.571$

Since the calculated value of t is less than the tabulated value of t we accept H_0 at 5% level of significance i.e., there is no major difference between the mean yield of two types diets.

10.9 DEGREES OF FREEDOM

In order to discuss the statistic t as discussed above, it is necessary to use observations obtained from a sample as well as certain population parameters. If these parameters are unknown, they must be estimated from the sample.

The number of degrees of freedom of a statistic, generally denoted by v, is defined as the number N of independent observation in the sample (i.e. the sample size) minus the number k of population parameters, which must be estimated from sample observations. In symbols, v = N - k.

In case of the statistic t, the number of independent observations in the sample is N, from which we compute \bar{x} S. However, since we must estimate μ , k = 1 and v = N - 1.

10.10 THE *f***-DISTRIBUTION**

If X and Y are two independent chi-square variates with v_1 and v_2 degree of freedom respectively, then F-statistic is defined by

$$F = \frac{X/v_1}{Y/v_2}$$

In other words, F is defined as the ration of two independent chi-square variates divided by their respective degrees of freedom and it follows Snedecor's F-distribution with (v_1, v_2) degree of freedom with probability function given by:

$$f(F) = \frac{\left(\frac{v_1}{v_2}\right)^{\frac{v_1}{2}}}{B\left(\frac{v_1}{2}, \frac{v_2}{2}\right)} \cdot \frac{F^{\frac{v_1}{2}-1}}{\left(1 + \frac{v_1}{v_2}F\right)^{\frac{(v_1+v_2)}{2}}}$$

Remark: The sampling distribution of F-statistic does not involve any population parameters and depends only on the degrees of freedom $v_1 \& v_2$.

F-test for Equality of Two population Variances.

Suppose we want to test (i) whether two independent samples x_i , (i=1,2,...n₁) and y_j , (j =1,2,...n₂) have been drawn from the normal population with the same variance σ^2 or (ii) whether the two independent estimates of the population variance are homogenous or not.

Under the Null hypothesis $H_0: \sigma_x^2 = \sigma_y^2 = \sigma^2$ i.e., the population variances are equal, or two independent estimates of the population variance are homogenous, the statistics F is given by

$$F = \frac{S_X^2}{S_Y^2}$$

where

$$S_X^2 = \frac{1}{n_1 - 1} \sum (x_i - \bar{x})^2, \ S_Y^2 = \frac{1}{n_2 - 1} \sum (y_i - \bar{y})^2$$

are unbiased estimates of the common population variance σ^2 obtained from two independent samples and it follows Snedecor's F-distribution with $(v_1, v_2) = (n_1 - 1, n_2 - 1)$ degree of freedom.



The shaded in the diagram indicates the acceptance region $(1 - \alpha)$ and the unshaded region indicates the rejection region α .

Eg: 1) In one sample of 8 observations, the sum of the squares of deviations of the sample values from the sample mean was 84.4 and in the other sample of 10 observations it was 102.6. Test whether this difference is significant at 5% LOS given that 5% point of F for $v_1 = 7$ and $v_2 = 9$ degree of freedom is 3.29.

Solution: $H_0: \sigma_x^2 = \sigma_y^2$ i.e., the estimate of variance given by the samples are homogenous, $H_1: \sigma_x^2 \neq \sigma_y^2$ $n_1 = 8, n_2 = 10, \sum (x_i - \bar{x})^2 = 84.4, \sum (y_i - \bar{y})^2 = 102.6$ $S_x^2 = \frac{1}{n_1 - 1} \sum (x_i - \bar{x})^2 = \frac{1}{7} \times 84.4 = 12.0571$ $S_y^2 = \frac{1}{n_2 - 1} \sum (y_i - \bar{y})^2 = \frac{1}{9} \times 102.6 = 11.4$ $F = \frac{S_x^2}{S_y^2} = \frac{12.0571}{11.4} = 1.0576$ tabulated $F_{7,9}(0.05) = 3.29$

 $F_{cal} = F = 1.0576 < F_{0.05}(7,9) = 3.29$ \therefore Accept H_0 at 5% LOS

Eg: 2) Two random samples gave the following results:

	ι <i>υ</i>	
Sample no	Size	sum of squares of deviation
		from the mean
1	10	90
2	12	108

Test at 5% LOS whether there is a difference in variance. {Given $F_{0.05}(9, 11) = 2.9$, $F_{0.05}(11,9) = 3.1$ }

Solution: $H_0: \sigma_x^2 = \sigma_y^2$, $H_1: \sigma_x^2 \neq \sigma_y^2$ $n_1 = 10, n_2 = 12, \sum (x_i - \bar{x})^2 = 90, \qquad \sum (y_i - \bar{y})^2 = 108$ $S_x^2 = \frac{1}{n_1 - 1} \sum (x_i - \bar{x})^2 = \frac{1}{9} \times 90 = 10$ $S_y^2 = \frac{1}{n_2 - 1} \sum (y_i - \bar{y})^2 = \frac{1}{11} \times 108 = 9.8182$ $F_{cal} = \frac{S_x^2}{S_y^2} = \frac{10}{9.8182} = 1.0185$ $F_{cal} = 1.0185 < F_{0.05}(9, 11) = 2.9$ $\therefore accept H_0$

10.11 SUMMARY

In this chapter we had learnt about to apply t-test for the sample size less than or equal to 30. We had also learnt to test if the sample mean (\bar{x}) differs significantly from the hypothetical value μ of the population mean and to test the significance of the difference between two sample means. WE had also learnt about the confidence interval and the F-test to whether the two independent estimates of the population variance are homogenous or not.

10.13 EXERCISES

- 1) A researcher is interested in determing whether or not review sessions affect exam performance. The independent variable, a review session, is administered to a sample of students (n=9) in an attempt to determine if this has an effect on the dependent variable, exam performance. Based on the information gathered in previous semesters, the researcher knows that the population mean for a given exam is 24. The sample mean is 25. with a S.D of 4, LOS = 5%.
- 2) You conduct a survey of a sample of 25 members of this year's graduating marketing students and find that average GPA is 3.2. The standard deviation of the sample is 4. Over the last year the average GPA has been 3.0. Is the GPA of this year's students significantly different from long run average?
- 3) The heights of 10 males of given locality are found to be 70, 67, 62, 68, 61, 68, 70, 64, 64, 66 inches. Is it reasonable to believe that the average height is greater than 64 inches?
- 4) A random sample of 16 values from a normal population showed a mean of 41.5 inches and the sum of squares of deviations from this mean = 135 sq. inches. Show that the assumptions of mean 43.5 inches for the population is not reasonable. Obtain the 95 % and 99 % confidence limits.
- 5) Below are given the gain in weights (in kgs) of pigs fed on two Diets A and B

Diet A	25	32	30	34	24	14	32	24	30	31	35	25			
Diet B	44	34	22	10	47	31	40	30	32	35	18	21	35	29	22

Test if the two diets differ significantly as regards to their effect on increase in weight.

6) Samples of two types of electric light bulbs were tested for length of life and following data were obtained:



Sample Size	8	7
Sample mean	1234	1036
Sample S.D.	36	40

Is the difference in the means sufficient to warrant that Type I is superior to Type II regarding length of life?

7) Two laboratories carry out independent estimates of a particular chemicals in a medicine produced by a certain firm. A sample is taken from each batch, halved and the separate halves sent to the two laboratories. The following data is obtained.

no. of samples	10
mean value of the diff. of estimates	0.6
sum of the squares of their diff. from their mean	20

Is the difference significant at 5% LOS.

- 8) A certain stimulus administered to each of the following 12 patients resulted in the following increase of blood pressure: 5, 2, 8, -1, 0,-2, 1, 5, 0,4 and 6. Can it be obtained that the stimulus will, in general, be accompanied by an increase in blood pressure?
- 9) Two independent samples of 8 and 7 items respectively had the following values of the variables:

Sample I	9	11	13	11	15	9	12	14
Sample II	10	12	10	14	9	8	10	

Do the estimates of population variance differ significantly?

10.14 SOLUTION TO EXERCISES

		0.11	G 1 .:
Q. No.	Solution	Q. No.	Solution
1	$H_0: \mu = 24, H_1: \mu > 24, t= 0.75$	2	$H_0: \mu = 3, H_1: \mu \neq 3, t$ = 0.2445
3	t = 2, table $t = 1.833$, Reject H ₀	4	$\begin{array}{l} H_0: \mu = 43.5 \ , H_1: \mu \neq 43.5, \\ t_{cal} = 2.667, tablet \\ = 2.131 \end{array}$
5	t = -0.609, accept H ₀	6	t = 9.3924, Reject H ₀
7	t = 2.2222, Accept H ₀	8	t= 2.89, Reject H ₀
9	Cal. F = 0.8315		

10.15 TABLES OF *t*-DISTRIBUTION AND *f*-DISTRIBUTION *t* DISTRIBUTION : CRITICAL VALUES OF *t*

Significance level

Degre	Two-tailed	10%	5%	2%	1%	0.2%	0.1%
es of	test:	10/0	070	2/0	170	0.270	0.170
freedo	One-tailed	5%	2.5%	1%	0.5%	0.1%	0.05%
<i>m</i>	test:	6.01.4	10 50 6	21.021	(2) (7 7	210.200	(2)((1))
1		6.314	12.706	31.821	63.657	318.309	636.619
2		2.920	4.303	6.965	9.925	22.327	31.599
3		2.353	3.182	4.541	5.841	10.215	12.924
4		2.132	2.776	3.747	4.604	7.173	8.610
5		2.015	2.5/1	3.365	4.032	5.893	6.869
6		1.943	2.447	3.143	3.707	5.208	5.959
7		1.894	2.365	2.998	3.499	4.785	5.408
8		1.860	2.306	2.896	3.355	4.501	5.041
9		1.833	2.262	2.821	3.250	4.297	4.781
10		1.812	2.228	2.764	3.169	4.144	4.587
11		1.796	2.201	2./18	3.106	4.025	4.437
12		1.782	2.179	2.681	3.055	3.930	4.318
13		1.7/1	2.160	2.650	3.012	3.852	4.221
14		1.752	2.145	2.624	2.977	3.787	4.140
15		1.753	2.131	2.602	2.947	3.733	4.073
16		1.746	2.120	2.583	2.921	3.686	4.015
17		1.740	2.110	2.567	2.898	3.646	3.965
18		1.734	2.101	2.552	2.8/8	3.610	3.922
19		1.729	2.093	2.539	2.861	3.579	3.883
20		1.725	2.086	2.528	2.845	3.552	3.850
21		1./21	2.080	2.518	2.831	3.527	3.819
22		1.714	2.074	2.508	2.819	3.505	3.792
23		1./14	2.069	2.500	2.807	3.485	3.768
24		1./M	2.064	2.492	2.797	3.46/	3.745
25		1.706	2.060	2.485	2.787	3.450	3.725
26		1.700	2.056	2.479	2.779	3.435	3.707
27		1.703	2.052	2.4/3	2.771	3.421	3.090
28		1./01	2.048	2.467	2.763	3.408	3.6/4
29		1.699	2.045	2.462	2.750	3.396	3.639
30		1.097	2.042	2.437	2.730	2.365	3.040
$\frac{32}{24}$		1.094	2.037	2.449	2.730	2 2 4 9	3.022
34		1.091	2.032	2.441	2.720	2 2 2 2 2	2 5 8 2
30		1.000	2.028	2.434	2.719	2 2 1 0	2.566
30		1.000	2.024	2.429	2.712 2.704	2 207	2.551
40		1.004	2.021	2.425	2.704	3.307	2 5 2 9
42		1.002	2.016	2.410	2.098	2 286	2 5 2 6
 //6		1.000	2.013	2.414	2.092	3.200	3 515
40		1.077	2.013	2.410	2.007	3 260	3 505
50		1.077	2.011	2.407	2.002	3 261	3 496
60		1.671	2.007	2.705	2.070	3 232	3 460
70		1.671	1 994	2.370	2.000	3 211	3 435
80		1.664	1 990	2.301	2.040	3 105	3 416
00		1.007	1.770	4.577	4.057	5.175	5.110

90	1.662	1.987	2.368	2.632	3.183	3.402
100	1.660	1.984	2.364	2.626	3.174	3.390
120	1.658	1.980	2.358	2.617	3.160	3.373
150	1.655	1.976	2.351	2.609	3.145	3.357
200	1.653	1.972	2.345	2.601	3.131	3.340
300	1.650	1.968	2.339	2.592	3.118	3.323
400	1.649	1.966	2.336	2.588	3.111	3.315
500	1.648	1.965	2.334	2.586	3.107	3.310
600	1.647	1.964	2.333	2.584	3.104	3.307
∞	1.645	1.960	2.326	2.576	3.090	3.291

 F Distribution : Critical Values of F (5%significancelevel)

 v1
 1
 2
 3
 4
 5
 6
 7
 8

 10
 12
 14
 16
 18
 20

 9

 v_2

1	161.45	5 199.50	215.71	224.58	230.16	233.99	236.7	7 238.88	240.54	241.88	243.91	245.36	246.46	247.32	248.01
2	18.51	19.00	19.16	19.25	19.30	19.33	19.35	19.37	19.38	19.40	19.41	19.42	19.43	19.44	19.45
3	10.13	9.55	9.28	9.12	9.01	8.94	8.89	8.85	8.81	8.79	8.74	8.71	8.69	8.67	8.66
4	7.71	6.94	6.59	6.39	6.26	6.16	6.09	6.04	6.00	5.96	5.91	5.87	5.84	5.82	5.80
5	6.61	5.79	5.41	5.19	5.05	4.95	4.88	4.82	4.77	4.74	4.68	4.64	4.60	4.58	4.56
6	5.99	5.14	4.76	4.53	4.39	4.28	4.21	4.15	4.10	4.06	4.00	3.96	3.92	3.90	3.87
7	5.59	4.74	4.35	4.12	3.97	3.87	3.79	3.73	3.68	3.64	3.57	3.53	3.49	3.47	3.44
8	5.32	4.46	4.07	3.84	3.69	3.58	3.50	3.44	3.39	3.35	3.28	3.24	3.20	3.17	3.15
9	5.12	4.26	3.86	3.63	3.48	3.37	3.29	3.23	3.18	3.14	3.07	3.03	2.99	2.96	2.94
10	4.96	4.10	3.71	3.48	3.33	3.22	3.14	3.07	3.02	2.98	2.91	2.86	2.83	2.80	2.77
11	4.84	3.98	3.59	3.36	3.20	3.09	3.01	2.95	2.90	2.85	2.79	2.74	2.70	2.67	2.65
12	4.75	3.89	3.49	3.26	3.11	3.00	2.91	2.85	2.80	2.75	2.69	2.64	2.60	2.57	2.54
13	4.67	3.81	3.41	3.18	3.03	2.92	2.83	2.77	2.71	2.67	2.60	2.55	2.51	2.48	2.46
14	4.60	3.74	3.34	3.11	2.96	2.85	2.76	2.70	2.65	2.60	2.53	2.48	2.44	2.41	2.39
15	4.54	3.68	3.29	3.06	2.90	2.79	2.71	2.64	2.59	2.54	2.48	2.42	2.38	2.35	2.33
16	4.49	3.63	3.24	3.01	2.85	2.74	2.66	2.59	2.54	2.49	2.42	2.37	2.33	2.30	2.28
17	4.45	3.59	3.20	2.96	2.81	2.70	2.61	2.55	2.49	2.45	2.38	2.33	2.29	2.26	2.23
18	4.41	3.55	3.16	2.93	2.77	2.66	2.58	2.51	2.46	2.41	2.34	2.29	2.25	2.22	2.19
19	4.38	3.52	3.13	2.90	2.74	2.63	2.54	2.48	2.42	2.38	2.31	2.26	2.21	2.18	2.16
20	4.35	3.49	3.10	2.87	2.71	2.60	2.51	2.45	2.39	2.35	2.28	2.22	2.18	2.15	2.12
21	4.32	3.47	3.07	2.84	2.68	2.57	2.49	2.42	2.37	2.32	2.25	2.20	2.16	2.12	2.10
22	4.30	3.44	3.05	2.82	2.66	2.55	2.46	2.40	2.34	2.30	2.23	2.17	2.13	2.10	2.07
23	4.28	3.42	3.03	2.80	2.64	2.53	2.44	2.37	2.32	2.27	2.20	2.15	2.11	2.08	2.05
24	4.26	3.40	3.01	2.78	2.62	2.51	2.42	2.36	2.30	2.25	2.18	2.13	2.09	2.05	2.03
25	4.24	3.39	2.99	2.76	2.60	2.49	2.40	2.34	2.28	2.24	2.16	2.11	2.07	2.04	2.01
26	4.22	3.37	2.98	2.74	2.59	2.47	2.39	2.32	2.27	2.22	2.15	2.09	2.05	2.02	1.99
27	4.21	3.35	2.96	2.73	2.57	2.46	2.37	2.31	2.25	2.20	2.13	2.08	2.04	2.00	1.97
28	4.20	3.34	2.95	2.71	2.56	2.45	2.36	2.29	2.24	2.19	2.12	2.06	2.02	1.99	1.96
29	4.18	3.33	2.93	2.70	2.55	2.43	2.35	2.28	2.22	2.18	2.10	2.05	2.01	1.97	1.94
30	4.17	3.32	2.92	2.69	2.53	2.42	2.33	2.27	2.21	2.16	2.09	2.04	1.99	1.96	1.93
35	4.12	3.27	2.87	2.64	2.49	2.37	2.29	2.22	2.16	2.11	2.04	1.99	1.94	1.91	1.88
40	4.08	3.23	2.84	2.61	2.45	2.34	2.25	2.18	2.12	2.08	2.00	1.95	1.90	1.87	1.84
50	4.03	3.18	2.79	2.56	2.40	2.29	2.20	2.13	2.07	2.03	1.95	1.89	1.85	1.81	1.78
60	4.00	3.15 2	2.76	2.53	2.37 2	2.25	2.17	2.10	2.04	1.99	1.92	1.86	1.82	1.78	1.75
70	3.98	3.13	2.74	2.50	2.35	2.23	2.14	2.07	2.02	1.97	1.89	1.84	1.79	1.75	1.72
80	3.96	3.11	2.72	2.49	2.33	2.21	2.13	2.06	2.00	1.95	1.88	1.82	1.77	1.73	1.70
90	3.95	3.10	2.71	2.47	2.32	2.20	2.11	2.04	1.99	1.94	1.86	1.80	1.76	1.72	1.69
100	3.94	3.09	2.70	2.46	2.31	2.19	2.10	2.03	1.97	1.93	1.85	1.79	1.75	1.71	1.68
120	3.92	3.07	2.68	2.45	2.29	2.18	2.09	2.02	1.96	1.91	1.83	1.78	1.73	1.69	1.66
150	3.90	3.06	2.00	2.43	2.27	2.10	2.07	2.00	1.94	1.89	1.82	1.76	1.71	1.07	1.04
200	3.89	3.04	2.65	2.42	2.26	2.14	2.06	1.98	1.93	1.88	1.80	1.74	1.69	1.66	1.62

250	3.88	3.03	2.64	2.41	2.25	2.13	2.05	1 98	1.92	1.87	1 79	1.73	1.68	1.65	1.61
300	3.87	3.03	2.63	2.40	2.24	2.13	2.04	1.97	1.91	1.86	1.78	1.72	1.68	1.64	1.61
400	3.86	3.02	2.63	2.39	2.24	2.12	2.03	1.96	1.90	1.85	1.78	1.72	1.67	1.63	1.60
500	3.86	3.01	2.62	2.39	2.23	2.12	2.03	1.96	1.90	1.85	1.77	1.71	1.66	1.62	1.59
600	3.86	3.01	2.62	2.39	2.23	2.11	2.02	1.95	1.90	1.85	1.77	1.71	1.66	1.62	1.59
750	3.85	3.01	2.62	2.38	2.23	2.11	2.02	1.95	1.89	1.84	1.77	1.70	1.66	1.62	1.58
1000	3.85	3.00	2.61	2.38	2.22	2.11	2.02	1.95	1.89	1.84	1.76	1.70	1.65	1.61	1.58

(continued)

f Distribution : Critical Values of f (5% significancelevel) v_1 25 30 35 40 50 60 75 100

r V V	2 2	25 200	30	35	40	5	0 (50	75	100	150
	1	249.26	250.10	250.69	251.14	251.77	252.20	252.62	253.04	253.46	253.68
	2	19.46	19.46	19.47	19.47	19.48	19.48	19.48	19.49	19.49	19.49
1	-										

2	19.46	19.46	19.47	19.47	19.48	19.48	19.48	19.49	19.49	19.49
3	8.63	8.62	8.60	8.59	8.58	8.57	8.56	8.55	8.54	8.54
4	5.77	5.75	5.73	5.72	5.70	5.69	5.68	5.66	5.65	5.65
5	4.52	4.50	4.48	4.46	4.44	4.43	4.42	4.41	4.39	4.39
6	3.83	3.81	3.79	3.77	3.75	3.74	3.73	3.71	3.70	3.69
7	3.40	3.38	3.36	3.34	3.32	3.30	3.29	3.27	3.26	3.25
8	3.11	3.08	3.06	3.04	3.02	3.01	2.99	2.97	2.96	2.95
9	2.89	2.86	2.84	2.83	2.80	2.79	2.77	2.76	2.74	2.73
10	2.73	2.70	2.68	2.66	2.64	2.62	2.60	2.59	2.57	2.56
11	2.60	2.57	2.55	2.53	2.51	2.49	2.47	2.46	2.44	2.43
12	2.50	2.47	2.44	2.43	2.40	2.38	2.37	2.35	2.33	2.32
13	2.41	2.38	2.36	2.34	2.31	2.30	2.28	2.26	2.24	2.23
14	2.34	2.31	2.28	2.27	2.24	2.22	2.21	2.19	2.17	2.16
15	2.28	2.25	2.22	2.20	2.18	2.16	2.14	2.12	2.10	2.10
16	2.23	2.19	2.17	2.15	2.12	2.11	2.09	2.07	2.05	2.04
17	2.18	2.15	2.12	2.10	2.08	2.06	2.04	2.02	2.00	1.99
18	2.14	2.11	2.08	2.06	2.04	2.02	2.00	1.98	1.96	1.95
19	2.11	2.07	2.05	2.03	2.00	1.98	1.96	1.94	1.92	1.91
20	2.07	2.04	2.01	1.99	1.97	1.95	1.93	1.91	1.89	1.88
21	2.05	2.01	1.98	1.96	1.94	1.92	1.90	1.88	1.86	1.84
22	2.02	1.98	1.96	1.94	1.91	1.89	1.87	1.85	1.83	1.82
23	2.00	1.96	1.93	1.91	1.88	1.86	1.84	1.82	1.80	1.79
24	1.97	1.94	1.91	1.89	1.86	1.84	1.82	1.80	1.78	1.77
25	1.96	1.92	1.89	1.87	1.84	1.82	1.80	1.78	1.76	1.75
26	1.94	1.90	1.87	1.85	1.82	1.80	1.78	1.76	1.74	1.73

27	1.92	1.88	1.86	1.84	1.81	1.79	1.76	1.74	1.72	1.71
28	1.91	1.87	1.84	1.82	1.79	1.77	1.75	1.73	1.70	1.69
29	1.89	1.85	1.83	1.81	1.77	1.75	1.73	1.71	1.69	1.67
30	1.88	1.84	1.81	1.79	1.76	1.74	1.72	1.70	1.67	1.66
35	1.82	1.79	1.76	1.74	1.70	1.68	1.66	1.63	1.61	1.60
40	1.78	1.74	1.72	1.69	1.66	1.64	1.61	1.59	1.56	1.55
50	1.73	1.69	1.66	1.63	1.60	1.58	1.55	1.52	1.50	1.48
60	1.69	1.65	1.62	1.59	1.56	1.53	1.51	1.48	1.45	1.44
70	1.66	1.62	1.59	1.57	1.53	1.50	1.48	1.45	1.42	1.40
80	1.64	1.60	1.57	1.54	1.51	1.48	1.45	1.43	1.39	1.38
90	1.63	1.59	1.55	1.53	1.49	1.46	1.44	1.41	1.38	1.36
100	1.62	1.57	1.54	1.52	1.48	1.45	1.42	1.39	1.36	1.34
120	1.60	1.55	1.52	1.50	1.46	1.43	1.40	1.37	1.33	1.32
150	1.58	1.54	1.50	1.48	1.44	1.41	1.38	1.34	1.31	1.29
200	1.56	1.52	1.48	1.46	1.41	1.39	1.35	1.32	1.28	1.26
250	1.55	1.50	1.47	1.44	1.40	1.37	1.34	1.31	1.27	1.25
300	1.54	1.50	1.46	1.43	1.39	1.36	1.33	1.30	1.26	1.23
400	1.53	1.49	1.45	1.42	1.38	1.35	1.32	1.28	1.24	1.22
500	1.53	1.48	1.45	1.42	1.38	1.35	1.31	1.28	1.23	1.21
600	1.52	1.48	1.44	1.41	1.37	1.34	1.31	1.27	1.23	1.20
750	1.52	1 47	1 44	1 41	1 37	1 34	1 30	1.26	1 22	1 20
1000	1.52	1.47	1.43	1.41	1.36	1.34	1.30	1.26	1.22	1.19
	-									

f Distribution : Critical Values of f(1%significancelevel)

<i>v</i> ₁ 1	2	3	4	5	6	7	8	9	10
	12	14	16	18	20				

 v_2

1	4052.18	4999.50	5403.35	5624.58	5763.65	5858.99	5928.36	5981.07	6022.47	6055.85	6106.32	6142.67	6170.10	6191.53	6208.73
2	98.50	99.00	99.17	99.25	99.30	99.33	99.36	99.37	99.39	99.40	99.42	99.43	99.44	99.44	99.45
3	34.12	30.82	29.46	28.71	28.24	27.91	27.67	27.49	27.35	27.23	27.05	26.92	26.83	26.75	26.69
4	21.20	18.00	16.69	15.98	15.52	15.21	14.98	14.80	14.66	14.55	14.37	14.25	14.15	14.08	14.02
5	16.26	13.27	12.06	11.39	10.97	10.67	10.46	10.29	10.16	10.05	9.89	9.77	9.68	9.61	9.55
6	13.75	10.92	9.78	9.15	8.75	8.47	8.26	8.10	7.98	7.87	7.72	7.60	7.52	7.45	7.40
7	12.25	9.55	8.45	7.85	7.46	7.19	6.99	6.84	6.72	6.62	6.47	6.36	6.28	6.21	6.16
8	11.26	8.65	7.59	7.01	6.63	6.37	6.18	6.03	5.91	5.81	5.67	5.56	5.48	5.41	5.36
9	10.56	8.02	6.99	6.42	6.06	5.80	5.61	5.47	5.35	5.26	5.11	5.01	4.92	4.86	4.81
10	10.04	7.56	6.55	5.99	5.64	5.39	5.20	5.06	4.94	4.85	4.71	4.60	4.52	4.46	4.41
11	9.65	7.21	6.22	5.67	5.32	5.07	4.89	4.74	4.63	4.54	4.40	4.29	4.21	4.15	4.10
12	9.33	6.93	5.95	5.41	5.06	4.82	4.64	4.50	4.39	4.30	4.16	4.05	3.97	3.91	3.86
13	9.07	6.70	5.74	5.21	4.86	4.62	4.44	4.30	4.19	4.10	3.96	3.86	3.78	3.72	3.66
14	8.86	6.51	5.56	5.04	4.69	4.46	4.28	4.14	4.03	3.94	3.80	3.70	3.62	3.56	3.51
15	8.68	6.36	5.42	4.89	4.56	4.32	4.14	4.00	3.89	3.80	3.67	3.56	3.49	3.42	3.37
16	8.53	6.23	5.29	4.77	4.44	4.20	4.03	3.89	3.78	3.69	3.55	3.45	3.37	3.31	3.26
17	8.40	6.11	5.18	4.67	4.34	4.10	3.93	3.79	3.68	3.59	3.46	3.35	3.27	3.21	3.16
18	8.29	6.01	5.09	4.58	4.25	4.01	3.84	3.71	3.60	3.51	3.37	3.27	3.19	3.13	3.08
19	8.18	5.93	5.01	4.50	4.17	3.94	3.77	3.63	3.52	3.43	3.30	3.19	3.12	3.05	3.00
20	8.10	5.85	4.94	4.43	4.10	3.87	3.70	3.56	3.46	3.37	3.23	3.13	3.05	2.99	2.94
21	8.02	5.78	4.87	4.37	4.04	3.81	3.64	3.51	3.40	3.31	3.17	3.07	2.99	2.93	2.88
22	7.95	5.72	4.82	4.31	3.99	3.76	3.59	3.45	3.35	3.26	3.12	3.02	2.94	2.88	2.83

								<u> </u>						<u> </u>	
23	7.88	5.66	4.76	4.26	3.94	3.71	3.54	3.41	3.30	3.21	3.07	2.97	2.89	2.83	2.78
24	7.82	5.61	4.72	4.22	3.90	3.67	3.50	3.36	3.26	3.17	3.03	2.93	2.85	2.79	2.74
25	7.77	5.57	4.68	4.18	3.85	3.63	3.46	3.32	3.22	3.13	2.99	2.89	2.81	2.75	2.70
26	7.72	5.53	4.64	4.14	3.82	3.59	3.42	3.29	3.18	3.09	2.96	2.86	2.78	2.72	2.66
27	7.68	5.49	4.60	4.11	3.78	3.56	3.39	3.26	3.15	3.06	2.93	2.82	2.75	2.68	2.63
28	7.64	5.45	4.57	4.07	3.75	3.53	3.36	3.23	3.12	3.03	2.90	2.79	2.72	2.65	2.60
29	7.60	5.42	4.54	4.04	3.73	3.50	3.33	3.20	3.09	3.00	2.87	2.77	2.69	2.63	2.57
30	7.56	5.39	4.51	4.02	3.70	3.47	3.30	3.17	3.07	2.98	2.84	2.74	2.66	2.60	2.55
35	7.42	5.27	4.40	3.91	3.59	3.37	3.20	3.07	2.96	2.88	2.74	2.64	2.56	2.50	2.44
40	7.31	5.18	4.31	3.83	3.51	3.29	3.12	2.99	2.89	2.80	2.66	2.56	2.48	2.42	2.37
50	7.17	5.06	4.20	3.72	3.41	3.19	3.02	2.89	2.78	2.70	2.56	2.46	2.38	2.32	2.27
60	7.08	4.98	4.13	3.65	3.34	3.12	2.95	2.82	2.72	2.63	2.50	2.39	2.31	2.25	2.20
70	7.01	4.92	4.07	3.60	3.29	3.07	2.91	2.78	2.67	2.59	2.45	2.35	2.27	2.20	2.15
80	6.96	4.88	4.04	3.56	3.26	3.04	2.87	2.74	2.64	2.55	2.42	2.31	2.23	2.17	2.12
90	6.93	4.85	4.01	3.53	3.23	3.01	2.84	2.72	2.61	2.52	2.39	2.29	2.21	2.14	2.09
100	6.90	4.82	3.98	3.51	3.21	2.99	2.82	2.69	2.59	2.50	2.37	2.27	2.19	2.12	2.07
120	6.85	4.79	3.95	3.48	3.17	2.96	2.79	2.66	2.56	2.47	2.34	2.23	2.15	2.09	2.03
150	6.81	4.75	3.91	3.45	3.14	2.92	2.76	2.63	2.53	2.44	2.31	2.20	2.12	2.06	2.00
200	6.76	4.71	3.88	3.41	3.11	2.89	2.73	2.60	2.50	2.41	2.27	2.17	2.09	2.03	1.97
250	6.74	4.69	3.86	3.40	3.09	2.87	2.71	2.58	2.48	2.39	2.26	2.15	2.07	2.01	1.95
300	6.72	4.68	3.85	3.38	3.08	2.86	2.70	2.57	2.47	2.38	2.24	2.14	2.06	1.99	1.94
400	6.70	4.66	3.83	3.37	3.06	2.85	2.68	2.56	2.45	2.37	2.23	2.13	2.05	1.98	1.92
500	6.69	4.65	3.82	3.36	3.05	2.84	2.68	2.55	2.44	2.36	2.22	2.12	2.04	1.97	1.92
600	6.68	4.64	3.81	3.35	3.05	2.83	2.67	2.54	2.44	2.35	2.21	2.11	2.03	1.96	1.91
750	6.67	4.63	3.81	3.34	3.04	2.83	2.66	2.53	2.43	2.34	2.21	2.11	2.02	1.96	1.90
1000	6.66	4.63	3.80	3.34	3.04	2.82	2.66	2.53	2.43	2.34	2.20	2.10	2.02	1.95	1.90
															L

continued)

f Di	f Distribution : Critical Values of f (1%significancelevel)										
<i>v</i> ₁	25	30	35	40	50	60	75	100	150		
	200										

1	6239 83	6260.65	6275 57	6286 78	6302.52	6313 03	6323 56	6334 11	6344 68	6349 97
2	99.46	99.47	99.47	99.47	99.48	99.48	99.49	99.49	99.49	99.49
3	26.58	26.50	26.45	26.41	26.35	26.32	26.28	26.24	26.20	26.18
4	13.91	13.84	13.79	13.75	13.69	13.65	13.61	13.58	13.54	13.52
5	9.45	9.38	9.33	9.29	9.24	9.20	9.17	9.13	9.09	9.08
6	7.30	7.23	7.18	7.14	7.09	7.06	7.02	6.99	6.95	6.93
7	6.06	5.99	5.94	5.91	5.86	5.82	5.79	5.75	5.72	5.70
8	5.26	5.20	5.15	5.12	5.07	5.03	5.00	4.96	4.93	4.91
9	4.71	4.65	4.60	4.57	4.52	4.48	4.45	4.41	4.38	4.36
10	4.31	4.25	4.20	4.17	4.12	4.08	4.05	4.01	3.98	3.96
11	4.01	3.94	3.89	3.86	3.81	3.78	3.74	3.71	3.67	3.66
12	3.76	3.70	3.65	3.62	3.57	3.54	3.50	3.47	3.43	3.41
13	3.57	3.51	3.46	3.43	3.38	3.34	3.31	3.27	3.24	3.22
14	3.41	3.35	3.30	3.27	3.22	3.18	3.15	3.11	3.08	3.06
15	3.28	3.21	3.17	3.13	3.08	3.05	3.01	2.98	2.94	2.92
16	3.16	3.10	3.05	3.02	2.97	2.93	2.90	2.86	2.83	2.81
17	3.07	3.00	2.96	2.92	2.87	2.83	2.80	2.76	2.73	2.71
18	2.98	2.92	2.87	2.84	2.78	2.75	2.71	2.68	2.64	2.62
19	2.91	2.84	2.80	2.76	2.71	2.67	2.64	2.60	2.57	2.55
20	2.84	2.78	2.73	2.69	2.64	2.61	2.57	2.54	2.50	2.48
21	2.79	2.72	2.67	2.64	2.58	2.55	2.51	2.48	2.44	2.42

$\begin{array}{ c c c c c c c c c c c c c c c c c c c$											
23 2.69 2.62 2.57 2.54 2.48 2.45 2.41 2.37 2.34 2.32 24 2.64 2.58 2.53 2.49 2.44 2.40 2.37 2.33 2.29 2.27 25 2.60 2.54 2.49 2.45 2.40 2.36 2.33 2.29 2.25 2.23 26 2.57 2.50 2.45 2.42 2.36 2.33 2.29 2.25 2.21 2.19 27 2.54 2.47 2.42 2.38 2.33 2.29 2.26 2.22 2.18 2.16 28 2.51 2.44 2.39 2.35 2.30 2.26 2.23 2.19 2.14 2.10 30 2.45 2.39 2.34 2.30 2.25 2.21 2.17 2.13 2.09 2.07 35 2.35 2.28 2.23 2.19 2.14 2.10 2.06 2.02 1.98	22	2.73	2.67	2.62	2.58	2.53	2.50	2.46	2.42	2.38	2.36
24 2.64 2.58 2.53 2.49 2.44 2.40 2.37 2.33 2.29 2.27 25 2.60 2.54 2.49 2.45 2.40 2.36 2.33 2.29 2.25 2.23 26 2.57 2.50 2.45 2.42 2.36 2.33 2.29 2.25 2.21 2.19 27 2.54 2.47 2.42 2.38 2.33 2.29 2.26 2.22 2.18 2.16 28 2.51 2.44 2.39 2.35 2.30 2.26 2.22 2.18 2.16 30 2.45 2.39 2.34 2.30 2.25 2.21 2.17 2.13 2.09 2.07 35 2.35 2.28 2.23 2.19 2.14 2.10 2.06 2.02 1.98 1.96 40 2.27 2.20 2.15 2.11 2.06 2.02 1.98 1.94 1.90 1.87	23	2.69	2.62	2.57	2.54	2.48	2.45	2.41	2.37	2.34	2.32
25 2.60 2.54 2.49 2.45 2.40 2.36 2.33 2.29 2.25 2.23 26 2.57 2.50 2.45 2.42 2.36 2.33 2.29 2.25 2.21 2.19 27 2.54 2.47 2.42 2.38 2.33 2.29 2.26 2.22 2.18 2.16 28 2.51 2.44 2.39 2.35 2.30 2.26 2.22 2.18 2.16 30 2.45 2.39 2.34 2.30 2.25 2.21 2.17 2.13 2.09 2.07 35 2.35 2.28 2.23 2.19 2.14 2.10 2.06 2.02 1.98 1.96 40 2.27 2.20 2.15 2.11 2.06 2.02 1.98 1.96 40 2.27 2.20 2.15 2.11 2.06 2.02 1.98 1.94 1.89 50 2.17 <	24	2.64	2.58	2.53	2.49	2.44	2.40	2.37	2.33	2.29	2.27
26 2.57 2.50 2.45 2.42 2.36 2.33 2.29 2.25 2.21 2.19 27 2.54 2.47 2.42 2.38 2.33 2.29 2.26 2.22 2.18 2.16 28 2.51 2.44 2.39 2.35 2.30 2.26 2.23 2.19 2.15 2.13 29 2.48 2.41 2.36 2.33 2.27 2.23 2.00 2.16 2.12 2.10 30 2.45 2.39 2.34 2.30 2.25 2.21 2.17 2.13 2.09 2.07 35 2.35 2.28 2.23 2.19 2.14 2.10 2.06 2.02 1.98 1.96 40 2.27 2.20 2.15 2.11 2.06 2.02 1.98 1.94 1.89 1.87 1.82 1.78 1.76 60 2.10 2.03 1.98 1.84 1.88 1.84	25	2.60	2.54	2.49	2.45	2.40	2.36	2.33	2.29	2.25	2.23
27 2.54 2.47 2.42 2.38 2.33 2.29 2.26 2.22 2.18 2.16 28 2.51 2.44 2.39 2.35 2.30 2.26 2.23 2.19 2.15 2.13 29 2.48 2.41 2.36 2.33 2.27 2.23 2.20 2.16 2.12 2.10 30 2.45 2.39 2.34 2.30 2.25 2.21 2.17 2.13 2.09 2.07 35 2.35 2.28 2.23 2.19 2.14 2.10 2.06 2.02 1.98 1.96 40 2.27 2.20 2.15 2.11 2.06 2.02 1.98 1.94 1.87 50 2.17 2.10 2.05 2.01 1.95 1.91 1.87 1.82 1.78 1.76 60 2.01 1.98 1.94 1.88 1.84 1.79 1.75 1.70 1.65 1.62 80 2.01 1.94 1.89 1.85 1.79 1.75 1	26	2.57	2.50	2.45	2.42	2.36	2.33	2.29	2.25	2.21	2.19
28 2.51 2.44 2.39 2.35 2.30 2.26 2.23 2.19 2.15 2.13 29 2.48 2.41 2.36 2.33 2.27 2.23 2.20 2.16 2.12 2.10 30 2.45 2.39 2.34 2.30 2.25 2.21 2.17 2.13 2.09 2.07 35 2.35 2.28 2.23 2.19 2.14 2.10 2.06 2.02 1.98 1.99 1.97 40 2.27 2.20 2.15 2.11 2.06 2.02 1.98 1.94 1.80 50 2.17 2.10 2.05 2.01 1.95 1.91 1.87 1.82 1.78 1.76 60 2.10 2.03 1.98 1.84 1.84 1.79 1.75 1.70 1.68 70 2.05 1.98 1.89 1.83 1.78 1.74 1.70 1.65 1.61 1.58	27	2.54	2.47	2.42	2.38	2.33	2.29	2.26	2.22	2.18	2.16
29 2.48 2.41 2.36 2.33 2.27 2.23 2.20 2.16 2.12 2.10 30 2.45 2.39 2.34 2.30 2.25 2.21 2.17 2.13 2.09 2.07 35 2.35 2.28 2.23 2.19 2.14 2.10 2.06 2.02 1.98 1.96 40 2.27 2.20 2.15 2.11 2.06 2.02 1.98 1.94 1.90 1.87 50 2.17 2.10 2.05 2.01 1.95 1.91 1.87 1.82 1.78 1.76 60 2.10 2.03 1.98 1.94 1.88 1.84 1.79 1.75 1.70 1.68 70 2.05 1.98 1.93 1.89 1.83 1.78 1.74 1.70 1.65 1.62 80 2.01 1.94 1.89 1.85 1.79 1.75 1.70 1.65 1.61	28	2.51	2.44	2.39	2.35	2.30	2.26	2.23	2.19	2.15	2.13
302.452.392.342.302.252.212.172.132.092.07352.352.282.232.192.142.102.062.021.981.96402.272.202.152.112.062.021.981.941.901.87502.172.102.052.011.951.911.871.821.781.76602.102.031.981.941.881.841.791.751.701.68702.051.981.931.891.831.781.741.701.651.62802.011.941.891.851.791.751.701.651.611.58901.991.921.861.821.761.721.671.621.571.551001.971.891.841.801.741.691.651.601.551.521201.931.861.811.761.701.661.611.561.511.481501.901.831.771.731.661.621.571.521.461.432001.871.791.741.691.631.581.531.481.421.392501.851.771.721.671.611.561.511.461.401.363001.841.761.701.661.51 <t< th=""><th>29</th><th>2.48</th><th>2.41</th><th>2.36</th><th>2.33</th><th>2.27</th><th>2.23</th><th>2.20</th><th>2.16</th><th>2.12</th><th>2.10</th></t<>	29	2.48	2.41	2.36	2.33	2.27	2.23	2.20	2.16	2.12	2.10
35 2.35 2.28 2.23 2.19 2.14 2.10 2.06 2.02 1.98 1.96 40 2.27 2.20 2.15 2.11 2.06 2.02 1.98 1.94 1.90 1.87 50 2.17 2.10 2.05 2.01 1.95 1.91 1.87 1.82 1.78 1.76 60 2.10 2.03 1.98 1.94 1.88 1.84 1.79 1.75 1.70 1.68 70 2.05 1.98 1.93 1.89 1.83 1.78 1.74 1.70 1.65 1.62 80 2.01 1.94 1.89 1.85 1.79 1.75 1.70 1.65 1.61 1.58 90 1.99 1.92 1.86 1.82 1.76 1.72 1.67 1.62 1.57 1.55 100 1.97 1.89 1.84 1.80 1.74 1.69 1.65 1.60 1.55	30	2.45	2.39	2.34	2.30	2.25	2.21	2.17	2.13	2.09	2.07
40 2.27 2.20 2.15 2.11 2.06 2.02 1.98 1.94 1.90 1.87 50 2.17 2.10 2.05 2.01 1.95 1.91 1.87 1.82 1.78 1.76 60 2.10 2.03 1.98 1.94 1.88 1.84 1.79 1.75 1.70 1.68 70 2.05 1.98 1.93 1.89 1.83 1.78 1.74 1.70 1.65 1.62 80 2.01 1.94 1.89 1.85 1.79 1.75 1.70 1.65 1.62 80 2.01 1.94 1.89 1.85 1.79 1.75 1.70 1.65 1.61 1.58 90 1.99 1.92 1.86 1.82 1.76 1.72 1.67 1.62 1.57 1.52 100 1.97 1.89 1.84 1.80 1.74 1.69 1.65 1.60 1.55 1.52	35	2.35	2.28	2.23	2.19	2.14	2.10	2.06	2.02	1.98	1.96
50 2.17 2.10 2.05 2.01 1.95 1.91 1.87 1.82 1.78 1.76 60 2.10 2.03 1.98 1.94 1.88 1.84 1.79 1.75 1.70 1.68 70 2.05 1.98 1.93 1.89 1.83 1.78 1.74 1.70 1.65 1.62 80 2.01 1.94 1.89 1.85 1.79 1.75 1.70 1.65 1.61 1.58 90 1.99 1.92 1.86 1.82 1.76 1.72 1.67 1.62 1.57 1.55 100 1.97 1.89 1.84 1.80 1.74 1.69 1.65 1.60 1.55 1.52 120 1.93 1.86 1.81 1.76 1.70 1.66 1.61 1.56 1.51 1.48 150 1.90 1.83 1.77 1.73 1.66 1.62 1.57 1.52 1.46	40	2.27	2.20	2.15	2.11	2.06	2.02	1.98	1.94	1.90	1.87
60 2.10 2.03 1.98 1.94 1.88 1.84 1.79 1.75 1.70 1.68 70 2.05 1.98 1.93 1.89 1.83 1.78 1.74 1.70 1.65 1.62 80 2.01 1.94 1.89 1.85 1.79 1.75 1.70 1.65 1.61 1.58 90 1.99 1.92 1.86 1.82 1.76 1.72 1.67 1.62 1.57 1.55 100 1.97 1.89 1.84 1.80 1.74 1.69 1.65 1.60 1.55 1.52 120 1.93 1.86 1.81 1.76 1.70 1.66 1.61 1.56 1.51 1.48 150 1.90 1.83 1.77 1.73 1.66 1.62 1.57 1.52 1.46 1.43 200 1.87 1.79 1.74 1.69 1.63 1.58 1.53 1.48 1.42	50	2.17	2.10	2.05	2.01	1.95	1.91	1.87	1.82	1.78	1.76
70 2.05 1.98 1.93 1.89 1.83 1.78 1.74 1.70 1.65 1.62 80 2.01 1.94 1.89 1.85 1.79 1.75 1.70 1.65 1.61 1.58 90 1.99 1.92 1.86 1.82 1.76 1.72 1.67 1.62 1.57 1.55 100 1.97 1.89 1.84 1.80 1.74 1.69 1.65 1.60 1.55 1.52 120 1.93 1.86 1.81 1.76 1.70 1.66 1.61 1.56 1.51 1.48 150 1.90 1.83 1.77 1.73 1.66 1.62 1.57 1.52 1.46 1.43 200 1.87 1.79 1.74 1.69 1.63 1.58 1.53 1.48 1.42 1.39 250 1.85 1.77 1.72 1.67 1.61 1.56 1.51 1.46 1.40	60	2.10	2.03	1.98	1.94	1.88	1.84	1.79	1.75	1.70	1.68
80 2.01 1.94 1.89 1.85 1.79 1.75 1.70 1.65 1.61 1.58 90 1.99 1.92 1.86 1.82 1.76 1.72 1.67 1.62 1.57 1.55 100 1.97 1.89 1.84 1.80 1.74 1.69 1.65 1.60 1.55 1.52 120 1.93 1.86 1.81 1.76 1.70 1.66 1.61 1.56 1.51 1.48 150 1.90 1.83 1.77 1.73 1.66 1.62 1.57 1.52 1.46 1.43 200 1.87 1.79 1.74 1.69 1.63 1.58 1.53 1.48 1.42 1.39 250 1.85 1.77 1.72 1.67 1.61 1.56 1.51 1.46 1.40 1.36 300 1.84 1.76 1.70 1.66 1.59 1.55 1.50 1.44 1.38 <th>70</th> <th>2.05</th> <th>1.98</th> <th>1.93</th> <th>1.89</th> <th>1.83</th> <th>1.78</th> <th>1.74</th> <th>1.70</th> <th>1.65</th> <th>1.62</th>	70	2.05	1.98	1.93	1.89	1.83	1.78	1.74	1.70	1.65	1.62
90 1.99 1.92 1.86 1.82 1.76 1.72 1.67 1.62 1.57 1.55 100 1.97 1.89 1.84 1.80 1.74 1.69 1.65 1.60 1.55 1.52 120 1.93 1.86 1.81 1.76 1.70 1.66 1.61 1.56 1.51 1.48 150 1.90 1.83 1.77 1.73 1.66 1.62 1.57 1.52 1.46 1.43 200 1.87 1.79 1.74 1.69 1.63 1.58 1.53 1.48 1.42 1.39 250 1.85 1.77 1.72 1.67 1.61 1.56 1.51 1.46 1.42 1.39 250 1.85 1.77 1.72 1.67 1.61 1.56 1.51 1.46 1.40 1.36 300 1.84 1.76 1.70 1.66 1.59 1.55 1.50 1.44 1.38 <th>80</th> <th>2.01</th> <th>1.94</th> <th>1.89</th> <th>1.85</th> <th>1.79</th> <th>1.75</th> <th>1.70</th> <th>1.65</th> <th>1.61</th> <th>1.58</th>	80	2.01	1.94	1.89	1.85	1.79	1.75	1.70	1.65	1.61	1.58
100 1.97 1.89 1.84 1.80 1.74 1.69 1.65 1.60 1.55 1.52 120 1.93 1.86 1.81 1.76 1.70 1.66 1.61 1.56 1.51 1.48 150 1.90 1.83 1.77 1.73 1.66 1.62 1.57 1.52 1.46 1.43 200 1.87 1.79 1.74 1.69 1.63 1.58 1.53 1.48 1.42 1.39 250 1.85 1.77 1.72 1.67 1.61 1.56 1.51 1.46 1.40 1.36 300 1.84 1.76 1.70 1.66 1.59 1.55 1.50 1.44 1.38 1.35 400 1.82 1.75 1.69 1.64 1.58 1.53 1.48 1.42 1.36 1.32 500 1.81 1.74 1.68 1.63 1.57 1.52 1.47 1.41 1.34 </th <th>90</th> <th>1.99</th> <th>1.92</th> <th>1.86</th> <th>1.82</th> <th>1.76</th> <th>1.72</th> <th>1.67</th> <th>1.62</th> <th>1.57</th> <th>1.55</th>	90	1.99	1.92	1.86	1.82	1.76	1.72	1.67	1.62	1.57	1.55
120 1.93 1.86 1.81 1.76 1.70 1.66 1.61 1.56 1.51 1.48 150 1.90 1.83 1.77 1.73 1.66 1.62 1.57 1.52 1.46 1.43 200 1.87 1.79 1.74 1.69 1.63 1.58 1.53 1.48 1.42 1.39 250 1.85 1.77 1.72 1.67 1.61 1.56 1.51 1.46 1.40 1.36 300 1.84 1.76 1.70 1.66 1.59 1.55 1.50 1.44 1.38 1.35 400 1.82 1.75 1.69 1.64 1.58 1.53 1.48 1.42 1.36 1.32 500 1.81 1.74 1.68 1.63 1.57 1.52 1.47 1.41 1.34 1.31 600 1.80 1.73 1.67 1.63 1.56 1.51 1.46 1.40 1.34 </th <th>100</th> <th>1.97</th> <th>1.89</th> <th>1.84</th> <th>1.80</th> <th>1.74</th> <th>1.69</th> <th>1.65</th> <th>1.60</th> <th>1.55</th> <th>1.52</th>	100	1.97	1.89	1.84	1.80	1.74	1.69	1.65	1.60	1.55	1.52
150 1.90 1.83 1.77 1.73 1.66 1.62 1.57 1.52 1.46 1.43 200 1.87 1.79 1.74 1.69 1.63 1.58 1.53 1.48 1.42 1.39 250 1.85 1.77 1.72 1.67 1.61 1.56 1.51 1.46 1.40 1.36 300 1.84 1.76 1.70 1.66 1.59 1.55 1.50 1.44 1.38 1.35 400 1.82 1.75 1.69 1.64 1.58 1.53 1.48 1.42 1.36 1.32 500 1.81 1.74 1.68 1.63 1.57 1.52 1.47 1.41 1.34 1.31 600 1.80 1.73 1.67 1.63 1.56 1.51 1.46 1.40 1.34 1.30 750 1.80 1.72 1.66 1.62 1.55 1.50 1.46 1.40 1.34 </th <th>120</th> <th>1.93</th> <th>1.86</th> <th>1.81</th> <th>1.76</th> <th>1.70</th> <th>1.66</th> <th>1.61</th> <th>1.56</th> <th>1.51</th> <th>1.48</th>	120	1.93	1.86	1.81	1.76	1.70	1.66	1.61	1.56	1.51	1.48
200 1.87 1.79 1.74 1.69 1.63 1.58 1.53 1.48 1.42 1.39 250 1.85 1.77 1.72 1.67 1.61 1.56 1.51 1.46 1.40 1.36 300 1.84 1.76 1.70 1.66 1.59 1.55 1.50 1.44 1.38 1.35 400 1.82 1.75 1.69 1.64 1.58 1.53 1.48 1.42 1.36 1.32 500 1.81 1.74 1.68 1.63 1.57 1.52 1.47 1.41 1.34 1.31 600 1.80 1.73 1.67 1.63 1.56 1.51 1.46 1.40 1.34 1.30 750 1.80 1.72 1.66 1.62 1.55 1.50 1.44 1.34 1.30 750 1.80 1.72 1.66 1.62 1.55 1.50 1.46 1.40 1.34 1.30 </th <th>150</th> <th>1.90</th> <th>1.83</th> <th>1.77</th> <th>1.73</th> <th>1.66</th> <th>1.62</th> <th>1.57</th> <th>1.52</th> <th>1.46</th> <th>1.43</th>	150	1.90	1.83	1.77	1.73	1.66	1.62	1.57	1.52	1.46	1.43
250 1.85 1.77 1.72 1.67 1.61 1.56 1.51 1.46 1.40 1.36 300 1.84 1.76 1.70 1.66 1.59 1.55 1.50 1.44 1.38 1.35 400 1.82 1.75 1.69 1.64 1.58 1.53 1.48 1.42 1.36 1.32 500 1.81 1.74 1.68 1.63 1.57 1.52 1.47 1.41 1.34 1.31 600 1.80 1.73 1.67 1.63 1.56 1.51 1.46 1.40 1.34 1.31 600 1.80 1.73 1.67 1.63 1.56 1.51 1.46 1.40 1.34 1.30 750 1.80 1.72 1.66 1.62 1.55 1.50 1.44 1.33 1.29 1.80 1.72 1.66 1.62 1.55 1.50 1.45 1.39 1.33 1.29 <th>200</th> <th>1.87</th> <th>1.79</th> <th>1.74</th> <th>1.69</th> <th>1.63</th> <th>1.58</th> <th>1.53</th> <th>1.48</th> <th>1.42</th> <th>1.39</th>	200	1.87	1.79	1.74	1.69	1.63	1.58	1.53	1.48	1.42	1.39
300 1.84 1.76 1.70 1.66 1.59 1.55 1.50 1.44 1.38 1.35 400 1.82 1.75 1.69 1.64 1.58 1.53 1.48 1.42 1.36 1.32 500 1.81 1.74 1.68 1.63 1.57 1.52 1.47 1.41 1.34 1.31 600 1.80 1.73 1.67 1.63 1.56 1.51 1.46 1.40 1.34 1.30 750 1.80 1.72 1.66 1.62 1.55 1.50 1.45 1.39 1.33 1.29	250	1.85	1.77	1.72	1.67	1.61	1.56	1.51	1.46	1.40	1.36
400 1.82 1.75 1.69 1.64 1.58 1.53 1.48 1.42 1.36 1.32 500 1.81 1.74 1.68 1.63 1.57 1.52 1.47 1.41 1.34 1.31 600 1.80 1.73 1.67 1.63 1.56 1.51 1.46 1.40 1.34 1.30 750 1.80 1.72 1.66 1.62 1.55 1.50 1.45 1.39 1.33 1.29	300	1.84	1.76	1.70	1.66	1.59	1.55	1.50	1.44	1.38	1.35
500 1.81 1.74 1.68 1.63 1.57 1.52 1.47 1.41 1.34 1.31 600 1.80 1.73 1.67 1.63 1.56 1.51 1.46 1.40 1.34 1.30 750 1.80 1.72 1.66 1.62 1.55 1.50 1.45 1.39 1.33 1.29	400	1.82	1.75	1.69	1.64	1.58	1.53	1.48	1.42	1.36	1.32
600 1.80 1.73 1.67 1.63 1.56 1.51 1.46 1.40 1.34 1.30 750 1.80 1.72 1.66 1.62 1.55 1.50 1.45 1.39 1.33 1.29 1000 1.72 1.66 1.62 1.55 1.50 1.45 1.39 1.33 1.29	500	1.81	1.74	1.68	1.63	1.57	1.52	1.47	1.41	1.34	1.31
750 1.80 1.72 1.66 1.62 1.55 1.50 1.45 1.39 1.33 1.29	600	1.80	1.73	1.67	1.63	1.56	1.51	1.46	1.40	1.34	1.30
	750	1.80	1.72	1.66	1.62	1.55	1.50	1.45	1.39	1.33	1.29
1000 1.79 1.72 1.66 1.61 1.54 1.50 1.44 1.38 1.32 1.28	1000	1.79	1.72	1.66	1.61	1.54	1.50	1.44	1.38	1.32	1.28

10.12 REFERENCE FOR FURTHER READING

Following books are recommended for further reading:

- Statistics by Murray R, Spiegel, Larry J. Stephens, Mcgraw Hill International Publisher, 4th edition
- Fundamental of Mathematical Statistics by S. C. Gupta and V. K. Kapoor, Sultan Chand and Sons publisher, 11th edition

11

THE CHI-SQUARE TEST

Unit Structure

- 11.0 Objectives
- 11.1 Introduction
- 11.2 Properties of Chi Square variate
- 11.3 The Chi Square test for Goodness of fit11.3.1 Decision Criterion
- 11.4 Test for Independence of Attributes
- 11.5 Yate's Correction for continuity
- 11.6 Test in $r \times c$ Contingency Table
- 11.7 Coefficient of Contingency
- 11.8 Correlation of Attributes
- 11.9 Additive Property of Chi Square variate
- 11.10 Summary
- 11.11 Exercises
- 11.12 Solution to Exercises
- 11.13 Table of Chi-Square distribution
- 11.14 Reference for further reading

11.0 OBJECTIVES

The chi-square test is a non-parametric test that compares two or more variables from randomly selected data. It helps find the relationship between two or more variables. The chi square distribution is a theoretical or mathematical distribution which has wide applicability in statistical work. The term 'chi square' (pronounced with a hard 'ch') is used because the Greek letter χ is used to define this distribution. It will be seen that the elements on which this distribution is based are squared, so that the symbol $\chi 2$ is used to denote the distribution.

11.1 INTRODUCTION

We know that if the probability distribution of the discrete random variable X is known we can find the probability distribution of the random variable $Y = X^2$. One may be interested in knowing whether we can find the probability distribution of random variable $Y = X^2$ if the probability distribution of a continuous random variable X is known. The answer is affirmative and in particular, if X is a standard normal variate (a continuous random variable) so that its probability density function is given by

$$f(x) = \frac{1}{\sqrt{2\pi}} e^{\left(-\frac{1}{2}\right)x^2}, -\infty < x < \infty = 0, \qquad otherwise$$

then $Y = X^2$ is also a continuous random variable whose probability density function is given by

$$g(y) = \frac{1}{\sqrt{2\pi}} \frac{1}{\sqrt{y}} e^{-\frac{y}{2}}, \quad y \ge 0$$

= 0, otherwise

Here, the distribution of Y is known as Chi-square distribution with one degree of freedom.

More generally, if X_1, X_2, \dots, X_n are n independent standard normal variates, then the distribution of random variable $U = X_1^2 + X_2^2 + \dots + X_n^2$ is given by the probability density function

$$h(u) = \frac{1}{2^{\frac{n}{2}}\Gamma(\frac{n}{2})}e^{-\frac{u}{2}u^{\frac{n}{2}-1}}, \qquad u \ge 0, n \in \mathbb{N}$$

= 0, otherwise where $\Gamma\left(\frac{n}{2}\right)$ is called gamma $\frac{n}{2}$ and is given by $\Gamma\left(\frac{n}{2}\right) = \int_0^\infty e^{-t} t^{\frac{n}{2}-1} dt$

Here, the distribution of random variable U is called Chi-square distribution with n degrees of freedom n degrees of freedom and U is called a Chi-square variate. Generally, a chi square variate is denoted by the square of greek letter chi, i.e χ^2 . Thus if χ^2 denotes a Chi-square variate with n degrees of freedom, then its probability density function is given by

11.2 PROPERTIES OF CHI-SQUARE VARIATE WITH *n* **DEGREES OF FREEDOM**

Let χ^2 denote a Chi-square variate with n degrees of freedom. Then we have the following properties:

1) The probability density function of χ^2 is given by

$$f(\chi^2) = \frac{1}{2^{\frac{n}{2}}\Gamma\left(\frac{n}{2}\right)} e^{-\frac{\chi^2}{2}} (\chi^2)^{\frac{n}{2}-1}, \qquad \chi^2 \ge 0, n \in \mathbb{N}$$
$$= 0, \qquad otherwise$$

2) The mean of χ^2 is $E(\chi^2) = n$

- 3) The variance of χ^2 is $V(\chi^2) = 2n$
- 4) The mode of $\chi^2 = n 2$
- 5) The frequency curve is given by $y = f(\chi^2)$ lies in the first quadrant and it is positively skewed curve, its tail on the right extends upto infinity, as given in the below diagram



 $p = \Pr[X \ge \chi^2]$ is denoted by α and $\chi^2 = \chi^2_{n,\alpha}$

- 6) The total area under the Chi-square curve is 1.
- 7) $p(\chi^2 > c) = area under the curve <math>y = f(\chi^2)$ to the right of $\chi^2 = c$
- 8) For a chi square variate with n degrees of freedom, if $p(\chi^2 > c) = \alpha$, then c is denoted by $\chi^2_{n,\alpha}$ i.e $p(\chi^2 > \chi^2_{n,\alpha}) = \alpha$
- 9) $\chi^2_{n,\alpha}$ is called α probability point of chi square distribution with n degrees of freedom.

Eg: 1) If a random variable X follows chi square distribution with 10 degrees of freedom find i) x_0 ii) x_1 & iii) α such that $p(X > x_0) = 0.95$, $p(X \le x_1) = 0.01 \& p(X > 18.3) = \alpha$

Solution: n = 10 – degree of freedom i) to find x_0 such that $p(X > x_0) = 0.95 \Rightarrow x_0 = \chi^2_{10,0.95} = 3.9403$

ii) to find x₁ such that $p(X \le x_1) = 0.01$ $p(X \le x_1) = 1 - p(X > x_1) \Rightarrow p(X > x_1) = 1 - 0.01 = 0.99$ $\Rightarrow p(X > x_1) = 0.99 \Rightarrow x_1 = \chi^2_{10,0.99} = 2.5582$ iii) $p(X > 18.3) = \alpha \Rightarrow \chi^2_{10,\alpha} = 18.3 \Rightarrow \alpha = 0.05$

Eg: 2) A random variable Y follows chi square distribution with S.D 4, Find y_0 if $p(Y \le y_0) = 0.05$.

Solution: S. $D = 4 \Rightarrow var = 16 \Rightarrow 2n = 16 \Rightarrow n = 8$ $p(Y \le y_0) = 0.05 \Rightarrow p(Y \le y_0) = 1 - p(Y > y_0) \Rightarrow p(Y > y_0)$ = 1 - 0.05 = 0.95 $p(Y > y_0) = 0.95 \Rightarrow y_0 = \chi^2_{8,0.95} \Rightarrow y_0 = 2.7326$

Eg: 3) If a random variable X follows chi square distribution with S.D 4. Find mean and mode.

Solution: *S*. *D* = 4 \Rightarrow *Var* = 16, *but variance of* $\chi^2 = 2n \Rightarrow n = 8$ *mean of* $\chi^2 = n = 8$ & *mode of* $\chi^2 = n - 2 = 6$

11.3 THE CHI SQUARE TEST FOR GOODNESS OF FIT

When we come across some observations on a random variable, our curiosity may tempt us to investigate whether it can be considered to be a random variable following a certain specified probability law. A technique to test whether a given frequency distribution of a random variable follows a certain specified distribution (known as theoretical distribution) was proposed by Karl Pearson. It involves a test statistic that can be shown to follow Chi-square distribution under certain assumptions and the test is known as *Chi-square test for goodness of fit.*

Suppose we have an observed frequency distribution with n classes having observed frequencies

 $O_1, O_2, O_3, \dots, O_n$ with $\sum_{i=1}^{i=n} O_i = N$. (Here the classes may correspond to the discrete values of a variable or groups of values of a variable or even to the groups corresponding to an attribute).

Further, suppose that according to our assumption to be called the null hypothesis H_0 , the expected frequencies are $E_1, E_2, E_3, \dots, E_n$ such that $\sum_{i=1}^{i=n} E_i = N$.

Then the *test statistic* proposed by Karl Pearson is given by $\chi^2 = \sum_{i=1}^{i=n} \frac{(O_i - E_i)^2}{E_i}$

It can be shown that under the assumptions

- 1) observations are drawn independently and at random.
- 2) null hypothesis H_0 is true i.e the expected frequencies are $E_1, E_2, E_3, \dots, E_n$ respectively.

- 3) total number of observations made = N is large
- 4) observed frequencies O_i^{s} are large
- 5) expected frequencies E_i^{s} are large so that the terms $\frac{O_i E_i}{E_i}$ can be considered to be negligible.

The test statistic $\chi^2 = \sum_{i=1}^{i=n} \frac{(O_i - E_i)^2}{E_i}$ follow Chi-square distribution with (n-1) degrees of freedom.

For all practical purposes, the assumptions regarding the distribution of test statistic χ^2 can be considered to be reasonable if

a) the number of observations = $N \ge 50$

b) the expected frequencies $E_i^{'s} \ge 5$.

In case when the expected frequencies is less than 5 we require to combine more than one neighboring classes so that the expected frequency for such class is not less than 5.

11.3.1 DECISION CRITERION

While performing a test for goodness of fit, we shall use the following decision criteria:

Reject
$$H_0$$
 if $\chi^2 = \sum_{i=1}^{i=n} \frac{(O_i - E_i)^2}{E_i} > \chi^2_{(n-1),\alpha}$

Do not reject
$$H_0$$
 i.e accept H_0 if $\chi^2 \leq \chi^2_{(n-1),\alpha}$

Eg: 1)The following data represents the last digit of the cars passing at a certain traffic signal observed during last 30 minutes for 180 cars.

last digit	0	1	2	3	4	5	6	7	8	9
frequency	12	20	14	12	21	18	17	26	19	21

Can we retain at 5% level of significance that all the digits are equally likely to occur?

Solution: we want to test whether all digits are equally likely to occur. If

all digits are equally likely,
$$p = \frac{1}{10}$$
. $E_i = Np = 180 \times \frac{1}{10} = 18$

*H*₀: All digits are equally likely to occur

 H_1 : not H_0 i.e logical negation to H_0 (i.e all digits are not equally likely)

 $LOS = 5\% i.e \ \alpha = 0.05, N = 180 \ge 50 \ and \ E_i = 18 \le 5, n = 10\chi^2$

$$=\sum_{i=0}^{n} \frac{(O_i - E_i)^2}{E_i}$$
 follows chiq square distribution with 9 degrees of freedom

Decision criteria is given by reject H_0 if $\chi^2 > \chi^2_{9,0.05}$ where $\chi^2_{9,0.05} = 16.9190$ do not reject H_0 if $\chi^2 \le \chi^2_{9,0.05}, \chi^2 = \sum_{i=0}^n \frac{(O_i - E_i)^2}{E_i}$

Digit	observed freq. (O _i)	Exp. freq. (E _i)	O _i - E _i	$\frac{(\boldsymbol{O}_i - \boldsymbol{E}_i)^2}{\boldsymbol{E}_i}$
0	12	18	-6	36/18
1	20	18	2	4/18
2	14	18	-4	16/18
3	12	18	-6	36/18
4	21	18	3	9/18
5	18	18	0	0
6	17	18	-1	1/18
7	26	18	8	64/18
8	19	18	1	1/18
9	21	18	3	9/18
			total	176/18 = 9.7778

$$\chi^2 = \sum_{i=0}^n \frac{(O_i - E_i)^2}{E_i} = 9.7778 < 16.9190 = \chi^2_{9,0.05}$$

do not reject H_0 i.e accept H_0 .

Eg: 2) As per Mendel's theory according to the shape and color, certain variety of pea that can be classified into four categories Round and yellow, Round and green, Angular and yellow, Angular and green occur in the proportion of 9:3:3:1. To test this a sample of N = 128 peas was taken and the following were the observed frequencies

RY - 66	RG – 28
AY – 29	AG – 5

Perform the chi square test for goodness of fit.

Solution: N = 128, n =4

The probability of occurrence and Expected frequencies are given by

category	\mathbf{p}_{i}	$E_i = Np_i$
RY	9/16	128 X 9/16 = 72
RG	3/16	128 X 3/16 = 24
AY	3/16	128 X 3/16 = 24
AG	1/16	128 X1/16 = 8

 \therefore H₀: The four categories of peas i.e RY, RG, AY, AG have expected frequencies 72, 24, 24, 8 resp.

H₁: not H₀ LOS = 5%, $\alpha = 0.05$ Decision criteria is given by *Reject H*₀ if $\chi^2 > \chi^2_{3,0.05} = 7.8147$ Do not reject H₀ if $\chi^2 \le 7.8147$, where $\chi^2 = \sum_{i=1}^{n} \frac{(o_i - E_i)^2}{E_i}$

category of peas	obs. freq.(O _i)	Exp. freq. (E _i)	$O_i - E_i$	$\frac{(O_i - E_i)^2}{E_i}$
RY	66	72	-6	0.5
RG	28	24	4	0.6667
AY	29	24	5	1.0417
AG	5	8	-3	1.125
			total	3.3334

$$\chi^2 = 3.3334 < \chi^2_{3,0.05} = 7.8147$$

 \Rightarrow Accept H_0 at 5% LOS.

Eg: 3) Four identical coins are tossed 100 times and the following results are obtained.

no. of heads (x)	0	1	2	3	4
frequency	8	29	40	19	4

Are there sufficient evidences to conclude that the coins are biased at 5% LOS.

Solution: let p denote the probability of getting a head with each of the four coins,

X: no. of heads follows binomial distribution with n= 4, p

 $X \sim B(n, p)$ i.e $X \sim B(4,p)$

H₀: the coins are unbiased i.e p = 1/2

 H_1 : the coins are baised i.e p $\neq \frac{1}{2}$

LOS = 5 %,
$$\alpha = 0.05$$

$$X \sim B(4,p) \Rightarrow p(x) = {4 \choose x} p^{x} q^{4-x}$$
$$p(0) = p^{0} q^{4} = \frac{1}{16}, p(1) = 4pq^{3} = \frac{4}{16}, p(2) = 6p^{2}q^{2} = \frac{6}{16}$$
$$175$$

$$p(3) = 4p^3q = \frac{4}{16}, \qquad p(4) = p^4q^0 = \frac{1}{16}$$

Decision criteria Reject H_0 if $\chi^2 > \chi^2_{4,0.05} = 9.4877$

Expected frequencies $E_i = N \cdot p(x)$

X	obs. freq. O _i	Exp. freq. E _i	O _i - E _i	$(O_i - E_i)^2$
				$\overline{E_i}$
0	8	100p(0) = 6.25	1.75	0.49
1	29	100p(1) = 25	4	0.64
2	40	100p(2) = 37.5	2.5	0.1667
3	19	100p(3) = 25	-6	1.44
4	4	100p(4) = 6.25	-2.25	0.81
			total	3.5467

 $\chi^{2} = 3.5467 < \chi^{2}_{4,0.05} = 9.4877$ Accept $H_{0}at 5\% LOS \Rightarrow coins are unbiased$

Eg: 4) The random variable X denotes the number of street accidents per week.

Х	0	1	2	3	4	5	6	7
obs. freq.	15	30	28	14	8	4	0	1
exp. freq.	14	27	27	18	9	4	1	0

Test whether the random variable X follows Poisson distribution with parameter m = 2 at 1% level of significance.

Solution: As the expected frequency is less than 5 for X = 5, 6 & 7 we combine them into one class

Х	0	1	2	3	4	5-7
obs. freq.	15	30	28	14	8	5
exp. freq.	14	27	27	18	9	5

 H_0 : X follows Poisson distributions with parameter m =2

 H_1 : not H_0

LOS = 1%, $\alpha = 0.01$

Decision criteria is given by Reject H_0 if $\chi^2 > \chi^2_{5,0.01} = 15.086$

Do not reject H₀ if $\chi^2 \le \chi^2_{5,0.01}$, where $\chi^2 = \sum_{i=1}^n \frac{(o_i - E_i)^2}{E_i}$

i	Х	obs. freq. O _i	exp. freq. E _i	$(O_i - E_i)$	$\frac{(O_i - E_i)^2}{E_i}$
1	0	15	14	1	0.0714
2	1	30	27	3	0.3333

3	2	28	27	1	0.0370
4	3	14	18	-4	0.8889
5	4	8	9	-1	0.1111
6	5-7	5	5	0	0
				total	1.4417

$$\chi^2 = 1.4417 < \chi^2_{5,0.01} = 15.086$$

Accept H_0 i.e at 1% level of significance the hypothesis that the variable X follows Poisson distribution with parameter m =2 is retainable.

11.4 TEST FOR INDEPENDENCE OF ATTRIBUTES

In this section, we consider the presence of two attributes among units from single population and the interest is centered around the possible dependence or independence of the attributes. In other words, on the basis of data regarding two attributes for some units from the population, we shall investigate whether the observed data provide sufficient reasons to reject the claim that the two attributes are independent of each other for the population under consideration. Such a test is called **test for independence of attributes**.

To understand the mechanism of the test, we consider the following table representing data for two attributes known as 2×2 contingency table. It is called a contingency table as it represents the information which can be attributed to chance as the information is regarding randomly selected persons.

		Attribute B		total
		B1	B2	
Attribute A	A1	a	b	a+b
	A2	с	d	c+d
		a+c	b+d	a+b+c+d=N

2×2 contingency table is given as below:

With the help of the test statistic $\chi^2 = \frac{N(ad-bc)^2}{(a+b)(a+c)(b+d)(c+d)}$ we can perform a test for testing the hypothesis H₀: the attributes A and B under the consideration are independent against the logical alternative H₁: not H₀i.e the attributes A and B are dependent subject to the conditions N \geq 50 and each of the observed frequencies a, b, c, d \geq 5.

The decision criterion at level of significance = α is given by $Reject H_0: if \chi^2 > \chi^2_{1,\alpha}, do not reject H_0: if \chi^2 \le \chi^2_{1,\alpha}$ where $\chi^2 = \frac{N(ad-bc)^2}{(a+b)(a+c)(b+d)(c+d)}$ where a,b,c,d are the observed frequencies with a+b+a+d = N

frequencies with a+b+c+d = N.

Eg: 1) The following results are obtained at the end of six months of a kind of psychotherapy given to a group of 120 patients and also for another group of 120 patients who were not given the psychotherapy.

	psychotherapy				
	given	not given			
condition improved	71	42			
condition did not improved	49	78			

Can we conclude at 5% LOS that the psychotherapy is effective?

Solution: H₀: Psychotherapy is not effective

H₁: psychotherapy is effective

 $LOS = 5\%, \alpha = 0.05$

Decision criteria Reject H_0 if $\chi^2 > \chi^2_{1,0.05} = 3.8415$

N = 240, a =71, b = 42, c = 49, d = 78

$$\chi^{2} = \frac{N(ad - bc)^{2}}{(a+b)(a+c)(b+d)(c+d)} = \frac{240(71 \times 78 - 49 \times 42)^{2}}{113 \cdot 120 \cdot 120 \cdot 127}$$

$$\chi^2 = 240 \times \frac{12110400}{206654400} = 14.0645$$

 $\Rightarrow \chi^2 = 14.0645 > \chi^2_{1,0.05} = 3.8415$

 \Rightarrow Reject H₀ at 5% level of significance, we may say that the Psychotherapy is effective at 5% LOS.

11.5 YATE'S CORRECTION FOR CONTINUITY

When the cell frequencies a, b, c, d as observed in case of the four classes corresponding to two attributes are small, we cannot use the test statistic χ^2 as defined in previous section i.e for when the assumption a, b, c, d are greater than or equal to 5 does not hold, the distribution of

 $\chi^2 = \frac{N(ad-bc)^2}{(a+b)(a+c)(b+d)(c+d)}$ cannot be considered to be Chi-square with one degree of freedom.

In such a case i.e if the cell frequencies a, b, c, d are not all greater than or equal to 5, we make the following adjustment called **Yate's correction**.

- if ad < bc add $\frac{1}{2}$ to a and b and subtract $\frac{1}{2}$ from both b and c both.
- if ad > bc add $\frac{1}{2}$ to b and c and subtract $\frac{1}{2}$ from both a and b both.

With this adjustments, we get the test statistic as $\chi^2 = \frac{N(|ad-bc|-\frac{N}{2})^2}{(a+b)(a+c)(b+d)(c+d)}$

The decision criterion at level of significance = α is given by $Reject H_0: if \chi^2 > \chi^2_{1,\alpha}, do \ not \ reject H_0: if \chi^2 \le \chi^2_{1,\alpha}$

where $\chi^2 = \frac{N(|ad-bc|-\frac{N}{2})^2}{(a+b)(a+c)(b+d)(b+c)}$

Eg: 1) In an experiment on immunization of cattle from tuberculosis the following results were obtained

	affected	unaffected
Inoculated	11	31
Not inoculated	14	4

Examine the effect of vaccine in controlling the incidence of the disease at 1% LOS.

Solution: H₀: the attributes are independent

H1: the attributes are not independent LOS = 1%, $\alpha = 0.01$ Decision criteria Reject H_0 iff $\chi^2 > \chi^2_{1,0.01} = 6.6349$

N = 60, a = 11, b = 31, c = 14, d = 4

$$\chi^{2} = \frac{N\left(|ad - bc| - \frac{N}{2}\right)^{2}}{(a+b)(a+c)(b+d)(c+d)} = \frac{60(|11 \times 4 - 31 \times 14| - 30)^{2}}{42 \cdot 18 \cdot 25 \cdot 35}$$
$$\chi^{2} = \frac{60(390 - 30)^{2}}{661500} = \frac{60 \times 360 \times 360}{661500} = 11.7551$$
$$\chi^{2} = 11.7551 > \chi^{2}_{1,0.01} = 6.6349$$
$$\Rightarrow reject H_{0} at 1\% LOs$$

We can say at 1% level of significance that the Inoculation and affection due to disease are dependent.

11.6 TEST IN r × c CONTINGENCY TABLE

If we have two attributes A and B classified into r and c classes respectively denoted by $A_1, A_2, \dots, A_r \& B_1, B_2, \dots, B_c$ then the observed frequency can be put in a tabular form with r rows and c columns called r × c contingency table.

If we use O_{ij} (i = 1,2,3,...,r, j = 1,2,3,...,c)to denote the observed frequency for attribute class A_iB_j , then r × c contingency table can be represented as shown below:

Attribute	B ₁	B ₂	B ₃		B _C	TOTAL
A_1	O ₁₁	O ₁₂	O ₁₃		O_{1C}	a_1
A_2	O ₂₁	O ₂₂	O ₂₃		O _{2C}	a ₂
A ₃	O ₃₁	O ₃₂	O ₃₃		O _{3C}	a 3
A _R	O _{R1}	O _{R2}	O _{R3}		O _{RC}	ar
TOTAL	b ₁	b ₂	b ₃		b _c	N

Here a_i 's represent total observed frequencies for attribute classes A_i 's and b_j 's represent the same for classes B_j 's, N being the overall total frequency.

Then with these notations, we can write down the test statistic as $\chi^2 = \sum_i \sum_j \frac{(o_{ij} - E_{ij})^2}{E_{ij}}$, where $E_{ij} = \frac{a_i b_j}{N}$ are the expected frequencies.

With the help of the test statistic $\chi^2 = \sum_i \sum_j \frac{(o_{ij} - E_{ij})^2}{E_{ij}}$ we can perform a test for testing the hypothesis H₀: the attributes A and B under the consideration are independent against the logical alternative H₁: not H₀i.e the attributes A and B are dependent subject to the conditions N \geq 50 and each of the observed frequencies \geq 5(O_{ij} \geq 5).

The decision criterion at level of significance = α is given by

Reject H_0 : if $\chi^2 > \chi^2_{(r-1)(c-1),\alpha}$ do not reject H_0 : if $\chi^2 \le \chi^2_{(r-1)(c-1),\alpha}$

Eg: 1) Using the data given in the following table decide whether we can conclude that standard of clothing of a salesman has significant effect on his performance in field selling at 5% LOS.

	Performa	Performance in Field Selling						
	Disappointing	Disappointing Satisfactory Excellent						
Poorly dressed	21	15	6	42				
Well dressed	24	35	26	85				
Very well dressed	35	80	58	173				
Total	80	130	90	300				

Solution: H₀: attributes are independent LOS = 5%, $\alpha = 0.05$

Reject H₀ iff $\chi^2 > \chi^2_{4,0.05} = 9.4877$, where $\chi^2 = \sum_i \sum_j \frac{(o_{Ij} - E_{ij})^2}{E_{ij}}$

$$E_{11} = \frac{a_1 b_1}{N} = \frac{42 \cdot 80}{300} = 11.2, E_{12} = \frac{a_1 b_2}{N} = \frac{42 \cdot 130}{300} = 18.2$$

$$E_{13} = \frac{a_1 b_3}{N} = \frac{42 \cdot 90}{300} = 12.6, E_{21} = \frac{a_2 b_1}{N} = \frac{85 \cdot 80}{300} = 22.67$$
$E_{31} = \frac{1}{N} = \frac{1}{300} = 46.13, E_{32} = \frac{1}{N} = \frac{1}{300} = 74.97$				
$E_{33} = \frac{a_3 b_3}{N} = \frac{173 \cdot 90}{300} = 51.9$				
O _{ij}	E _{ij}	O _{ij} - E _{ij}	$\frac{\left(O_{ij}-E_{ij}\right)^2}{E_{ij}}$	
21	11.2	9.8	8.575	
15	18.2	-3.2	0.5626	
6	12.6	-6.6	3.4571	
24	22.67	1.33	0.078	
35	36.83	-1.83	0.0909	
26	25.5	0.5	0.0098	
35	46.13	-11.13	2.6854	
80	74.97	5.03	0.3375	
58	51.9	6.1	0.7170	
		total	16.5133	

 $173 \cdot 130$

 a_3b_2

 $173 \cdot 80$

 a_3b_1

 $\chi^2 = 16.5133 > \chi^2_{4,0.05} = 9.488$

 \Rightarrow reject H_0 at 5% LOS

We decide to reject H_0 at 5% level of significance and conclude that the standard of clothing of a salesman has significant effect on his performance in field selling.

11.7 COEFFICIENT OF CONTINGENCY

A measure of the degree of relationship, association, or dependence of the classifications in a contingency table is given by

$$C = \sqrt{\frac{\chi^2}{\chi^2 + N}}$$

which is called the *coefficient of contingency*. The larger the C, the greater is the degree of association. The number of rows and columns in the contingency table determines the maximum value of C, which is never greater than 1. If the number of rows and columns of a contingency table

is equal to k, the maximum value of C is given by $\sqrt{\frac{k-1}{k}}$.

Eg: 1) Using the data given in the following table decide whether we can conclude that standard of clothing of a salesman has significant effect on his performance in field selling at 5% LOS.

	Performance in Field Selling			
	Disappointing	Satisfactory	Excellent	Total
Poorly dressed	21	15	6	42
Well dressed	24	35	26	85
Very well dressed	35	80	58	173
Total	80	130	90	300

Also find the coefficient of contingency.

Solution: H₀: attributes are independent

LOS = 5%,
$$\alpha = 0.05$$

Reject H₀ iff $\chi^2 > \chi^2_{4,0.05} = 9.4877$, where $\chi^2 = \sum_i \sum_j \frac{(o_{Ij} - E_{ij})^2}{E_{ij}}$
 $E_{11} = \frac{a_1 b_1}{N} = \frac{42 \cdot 80}{300} = 11.2, E_{12} = \frac{a_1 b_2}{N} = \frac{42 \cdot 130}{300} = 18.2$
 $E_{13} = \frac{a_1 b_3}{N} = \frac{42 \cdot 90}{300} = 12.6, E_{21} = \frac{a_2 b_1}{N} = \frac{85 \cdot 80}{300} = 22.67$
 $E_{31} = \frac{a_3 b_1}{N} = \frac{173 \cdot 80}{300} = 46.13, E_{32} = \frac{a_3 b_2}{N} = \frac{173 \cdot 130}{300} = 74.97$
 $E_{33} = \frac{a_3 b_3}{N} = \frac{173 \cdot 90}{300} = 51.9$

O _{ij}	E _{ij}	O _{ij} - E _{ij}	$\frac{\left(O_{ij}-E_{ij}\right)^2}{E_{ij}}$
21	11.2	9.8	8.575
15	18.2	-3.2	0.5626
6	12.6	-6.6	3.4571
24	22.67	1.33	0.078
35	36.83	-1.83	0.0909
26	25.5	0.5	0.0098
35	46.13	-11.13	2.6854
80	74.97	5.03	0.3375
58	51.9	6.1	0.7170
		total	16.5133

 $\chi^2 = 16.5133 > \chi^2_{4,0.05} = 9.488 \Rightarrow reject H_0 \ at \ 5\% \ LOS$

We decide to reject H_0 at 5% level of significance and conclude that the standard of clothing of a salesman has significant effect on his performance in field selling.

$$C = \sqrt{\frac{\chi^2}{\chi^2 + N}} = \sqrt{\frac{16.5133}{16.5133 + 300}} = 0.2284$$

11.8 CORRELATION OF ATTRIBUTES

Because classifications in a contingency table often describe characteristics of individuals or objects, they are often referred to as *attributes*, and the degree of dependence, association, or relationship is called the *correlation of attributes*. For k X k tables, we define

$$r = \sqrt{\frac{\chi^2}{N(k-1)}}$$

as the correlation coefficient between attributes(or classification). This coefficient lies between 0 and 1. For 2×2 tables in which k = 2, the correlation is often called *tetrachoric* correlation.

11.9 ADDITIVE PROPERTY OF CHI SQUARE VARIATE

Suppose that the results of repeated experiments yield sample values of χ^2 given by

 $\chi_1^2, \chi_2^2, \chi_3^2, \dots, \chi_n^2$ with $v_1, v_2, v_3, \dots, v_n$ degrees of freedom, respectively. Then the result of all these experiment can be considered equivalent to χ^2 value given by $\chi_1^2 + \chi_2^2 + \chi_3^2 + \dots + \chi_n^2$ with $v_1 + v_2 + v_3 + \dots + v_n$ degrees of freedom.

11.10 SUMMARY

In this chapter we discussed about the chi-square test which is a nonparametric test that compares two or more variables from randomly selected data. It helps to find the relationship between two or more variables. The chi square distribution is a theoretical or mathematical distribution which has wide applicability in statistical work. We had also seen the various properties of Chi-square variate along with different tests to check whether the variable follows Chi-square distribution, along with Yate's correction.

11.11 EXERCISES

1) If X is a chi square variate with 17 degreed of freedom find x_0 , x_1 and α such that

 $p(X > x_0) = 0.01, p(X \le x_1) = 0.95 \& p(X \le 8.67) = \alpha.$

2) The following table gives the result of investigation in the association of eye color and hair color. Can we deduce that the two attributes are independent?

	Brown	Black
Eye color Blue	75	25
Brown	65	35

3) The following data show the classification of individuals with respect to gender and literacy in a random sample of 200. Test the data for independence of attributes using chi square at 1% LOS.

	literate	illiterate
male	95	5
female	75	25

4) The eyesight of 100 randomly selected people from a town were tested with the following results:

	poor	good
	eyesight	eyesight
male	200	350
female	200	250

Can we conclude at 5% level of significance that gender has no bearing on the quality of eyesight?

5) The following table shows results of inoculation against cholera

	not attacked	attacked
Inoculated	446	4
Not inoculated	291	9

Can we say at 5% LOS that inoculation is effective in controlling susceptibility of cholera?

6) The following are the results of the tests performed on two brands of tyres manufactured by a manufacturer.

	Brand A	Brand B
Lasted more than 30000km	27	38
Failed to last 30000 km	18	27

Use chi square at 5% LOS to test whether we can say that the two brands of tyres differ significantly or not as regards their lifespan.

- 7) Determine at 1% LOS whether vaccination can be regarded as a preventive measure for small pox on the basis of following report: Out of 1482 persons in a locality exposed to small pox, 368 in all were attacked. Out of 1482 persons, 343 had been vaccinated and of these only 35 were attacked.
- 8) During a market research survey organized by ABC Ltd. the households were asked whether they used "Beauty Soap"(the price of which is Rs. 7.25 a piece) and whether their per capita monthly expenditure exceeded Rs. 800 with the following results.

		Monthly per capita expenditure		
		Exceeded Rs 800 Did not exceed Rs. 800		
Whether they used	yes	6	20	
"Beauty Soap"	no	4	30	

Can we say the use of "Beauty Soap" depends on monthly per capita expenditure at 5% LOS.

9) In a household survey conducted in certain locality, the following information is collected.

	Whether exclusive Indoor toilet facility is available			
	yes no			
owned house	9	4		
rented house	21	16		

Use chi square test for independence at 1% LOS and give your conclusions.

10) The following data refer to an investigation carried out to examine the effect of T.V as a medium of advertisement on the turnover of the company's manufacturing and selling consumer products. A random sample of 40 companies was selected. Analyze the data and comment on your findings.

	Annual turnover rupees	exceeding one crore
	Yes	No
T.V advertisement	7	3
No T.V advertisement	10	20

11) A random sample of students of Mumbai University was selected and asked their opinion about autonomous college. The results are given below. The same number of each gender was included within each class group. Test the hypothesis at 5% LOS that opinions are independent of the class groupings.

	favoring autonomous colleges	opposed to autonomous college
First Year	120	80
Second Year	130	70
Third year	70	30
Post Graduation	80	20

12) Test for independence between health and working capacity from one following data:

		Health				
		very good good fair				
working	good	20	25	15		
Capacity	bad	10	15	15		

13) ABC Ltd. employ a large number of handicapped persons. The following is an account of the performance of 200 randomly chosen employees of the company:

	performance		
	above average	average	below average
handicapped	29	31	20
non handicapped	55	30	35

Can we retain at 5% LOS that handicapped employees are equally efficient as the non-handicapped employees of the company.

14) In a survey 100 couples are interviewed and they were asked to give an opinion on the importance of amiable nature of partner in selection of a bride or a groom. The ranks were given independently by them as I or II or III.

		Rankings by wives				
I I II III						
Ranking	Ι	25	12	8		
by	II	10	23	8		
Husbands	III	5	5	4		

Use chi square test for independence and comment at 5% LOS.

15) A socio economic survey conducted in 1981 in Mumbai revealed the following results:

	Monthly family income						
	Below 1200 1200 to 1800 1800 and above						
No child	18	15	12				
one child	31	25					
Two or more children	81	51	63				

Can we regard at 1% LOS that the number of children in the family has no association with monthly income?

16) The data in the table were collected on how individuals prepared their taxes and their education level. The null hypothesis is that the way people prepare their taxes (computer software or pen and paper) is independent of their education level. The following table is the contingency table.

Education		
High School	Bachelors	Masters
23	35	42
45	30	25
	Education High School 23 45	EducationHigh SchoolBachelors23354530

Find the coefficient of contingency.

11.12 SOLUTION TO EXERCISE

Q.	Solution	Q. No.	Solution
No.			

1	$ \begin{array}{l} x_0 = 33.4087, x_1 \\ = 27.5871, \\ \alpha = 0.05 \end{array} $	2	2.38, accept H ₀
3	15.686, reject H ₀	4	3.84, donot reject H ₀
5	3.552, accept H ₀	6	0.026, accept
7	51.157 reject	8	0.6652 accept
9	0.2122, accept	10	2.762, accept
11	12.046, reject	12	1.9097, accept
13	4.904, accept	14	10.8125, accept
15	4.305, accept	16	0.236

11.13 TABLE OF CHI-SQUARE DISTRIBUTION



df\area	0.995	0.99	0.975	0.95	0.9	0.75	0.5	0.25	0.1	0.05	0.025	0.01	0.005
1	0.00004	0.00016	0.00098	0.00393	0.01579	0.10153	0.45494	1.3233	2.70554	3.84146	5.02389	6.6349	7.87944
2	0.01003	0.0201	0.05064	0.10259	0.21072	0.57536	1.38629	2.77259	4.60517	5.99146	7.37776	9.21034	10.59663
3	0.07172	0.11483	0.2158	0.35185	0.58437	1.21253	2.36597	4.10834	6.25139	7.81473	9.3484	11.34487	12.83816
4	0.20699	0.29711	0.48442	0.71072	1.06362	1.92256	3.35669	5.38527	7.77944	9.48773	11.14329	13.2767	14.86026
5	0.41174	0.5543	0.83121	1.14548	1.61031	2.6746	4.35146	6.62568	9.23636	11.0705	12.8325	15.08627	16.7496
6	0.67573	0.87209	1.23734	1.63538	2.20413	3.4546	5.34812	7.8408	10.64464	12.59159	14.44938	16.81189	18.54758
7	0.98926	1.23904	1.68987	2.16735	2.83311	4.25485	6.34581	9.03715	12.01704	14.06714	16.01276	18.47531	20.27774
8	1.34441	1.6465	2.17973	2.73264	3.48954	5.07064	7.34412	10.21885	13.36157	15.50731	17.53455	20.09024	21.95495
9	1.73493	2.0879	2.70039	3.32511	4.16816	5.89883	8.34283	11.38875	14.68366	16.91898	19.02277	21.66599	23.58935
10	2.15586	2.55821	3.24697	3.9403	4.86518	6.7372	9.34182	12.54886	15.98718	18.30704	20.48318	23.20925	25.18818
11	2.60322	3.05348	3.81575	4.57481	5.57778	7.58414	10.341	13.70069	17.27501	19.67514	21.92005	24.72497	26.75685
12	3.07382	3.57057	4.40379	5.22603	6.3038	8.43842	11.34032	14.8454	18.54935	21.02607	23.33666	26.21697	28.29952
13	3.56503	4.10692	5.00875	5.89186	7.0415	9.29907	12.33976	15.98391	19.81193	22.36203	24.7356	27.68825	29.81947
14	4.07467	4.66043	5.62873	6.57063	7.78953	10.16531	13.33 927	17.11693	21.06414	23.68479	26.11895	29.14124	31.31935
15	4.60092	5.22935	6.26214	7.26094	8.54676	11.03654	14.33886	18.24509	22.30713	24.99579	27.48839	30.57791	32.80132
16	5.14221	5.81221	6.90766	7.96165	9.31224	11.91222	15.3385	19.36886	23.54183	26.29623	28.84535	31.99993	34.26719
17	5.69722	6.40776	7.56419	8.67176	10.08519	12.79193	16.33818	20.48868	24.76904	27.58711	30.19101	33.40866	35.71847
18	6.2648	7.01491	8.23075	9.39046	10.86494	13.67529	17.3379	21.60489	25.98942	28.8693	31.52638	34.80531	37.15645
19	6.84397	7.63273	8.90652	10.11701	11.65091	14.562	18.33765	22.71781	27.20357	30.14353	32.85233	36.19087	38.58226
20	7.43384	8.2604	9.59078	10.85081	12.44261	15.45177	19.33743	23.82769	28.41198	31.41043	34.16961	37.56623	39.99685
21	8.03365	8.8972	10.2829	11.59131	13.2396	16.34438	20.33723	24.93478	29.61509	32.67057	35.47888	38.93217	41.40106
22	8.64272	9.54249	10.98232	12.33801	14.04149	17.23962	21.33704	26.03927	30.81328	33.92444	36.78071	40.28936	42.79565
23	9.26042	10.19572	11.68855	13.09051	14.84796	18.1373	22.33688	27.14134	32.0069	35.17246	38.07563	41.6384	44.18128
24	9.88623	10.85636	12.40115	13.84843	15.65868	19.03725	23.33673	28.24115	33.19624	36.41503	39.36408	42.97982	45.55851
25	10.51965	11.52398	13.11972	14.61141	16.47341	19.93934	24.33659	29.33885	34.38159	37.65248	40.64647	44.3141	46.92789
26	11.16024	12.19815	13.8439	15.37916	17.29188	20.84343	25.33646	30.43457	35.56317	38.88514	41.92317	45.64168	48.28988
27	11.80759	12.8785	14.57338	16.1514	18.1139	21.7494	26.33634	31.52841	36.74122	40.11327	43.19451	46.96294	49.64492
28	12.46134	13.56471	15.30786	16.92788	18.93924	22.65716	27.33623	32.62049	37.91592	41.33714	44.46079	48.27824	50.99338
29	13.12115	14.25645	16.04707	17.70837	19.76774	23.56659	28.33613	33.71091	39.08747	42.55697	45.72229	49.58788	52.33562
30	13.78672	14.95346	16.79077	18.49266	20.59923	24.47761	29.33603	34.79974	40.25602	43.77297	46.97924	50.89218	53.67196

11.14 REFERENCE FOR FURTHER READING

Following books are recommended for further reading:

- Statistics by Murray R, Spiegel, Larry J. Stephens, Mcgraw Hill International Publisher, 4th edition
- Fundamental of Mathematical Statistics by S. C. Gupta and V. K. Kapoor, Sultan Chand and Sons publisher, 11th edition

UNIT V

12

CURVE FITTING AND THE METHOD OF LEAST SQUARES

Unit Structure

- 12.0 Objectives
- 12.1 Introduction
- 12.2 Relationship between variables
- 12.3 Curve fitting
- 12.4 Equations of Approximating Curves
- 12.5 Freehand Method of Curve Fitting
- 12.6 The Straight line Method
- 12.7 Least Square Curve fitting
 - 12.7.1 Straight Line
 - 12.7.2 Parabola
 - 12.7.3 Non-Linear relationship
- 12.8 Regression
- 12.9 Applications to Time Series
- 12.10 Problems involving more than two variables
- 12.11 Summary
- 12.12 Exercises
- 12.13 Solution to Exercises
- 12.14 Logarithm tables
- 12.15 Reference for further reading

12.0 OBJECTIVES

The main objective is to study the fitting of curves using the method of least square which minimizes the sum of the square of the errors.

12.1 INTRODUCTION

When we come across data for two variables and think about the relation between each other, two objects come to our mind:

- 1) To translate the possible relationship into a mathematical equation(called equation of curve)
- 2) To exploit the relationship between the variables for estimating the value of one variable corresponding to a given value of the other variable.

Curve fitting helps us in serving the first object directly, i.e., it leads to a mathematical equation describing the relationship between the variables. Further, using this equation one can estimate the value of the other variable.

12.2 RELATIONSHIP BETWEEN VARIABLES

Very often in practice a relationship is found to exist between two (or more) variables. For example, weights of adult males depend to some degree on their heights, the circumferences of circles depend on their radii, and the pressure of a given mass of gas depends on its temperature and volume.

It is frequently desirable to express this relationship in mathematical form by determining an equation that connects the variables.

12.3 CURVE FITTING

To determine an equation that connects variables, a first step is to collect data that show corresponding values of the variables under consideration. For example, suppose X and Y denote respectively, the height and weight of adult males; then a sample of N individuals would reveal the heights $X_1, X_2, X_3, \dots, X_N$ and the corresponding weights $Y_1, Y_2, Y_3, \dots, Y_N$.

A next step is plot the points $(X_1, Y_1), (X_2, Y_2), (X_3, Y_3), \dots, (X_N, Y_N)$ on a rectangular coordinate system. The resulting set of points is some times called a **scatter diagram**. From the scatter diagram it is often possible to visualize a smooth curve that approximates the data. Such a curve is called an **approximating curve**.

For example, in the following diagram the data appear to be approximated by a straight line, and so we call that a linear relationship exists between the variables.



In the following diagram, however the relationship exists between the variables, it is not a linear relationship, and so we call it a non-linear relationship.



The general problem of finding equations of approximating curves that fit the given sets of data is called **curve fitting**.

12.4 EQUATIONS OF APPROXIMATING CURVES

Various common types of approximating curves and their equations are given below for reference. All letters other than x and y shall be treated as constants. The variables x and y are termed as independent variable and dependent variable respectively, roles of the variables can be interchanged as per the requirement.

Straight Line	y = a + bx
Parabola or quadratic curve	$y = a + bx + cx^2$
Cubic curve	$y = a + bx + cx^2 + dx^3$
Quadratic curve	$y = a + bx + cx^2 + dx^3 + ex^4$
nth degree curve	$y = a_0 + a_1 x + a_2 x^2 + \dots + a_n x^n$

The above equations refer to the polynomial equations of the degree one, two, three, four and n respectively. The following are some examples of other equations most commonly used.

Hyperbola	$y = \frac{1}{a_0 + a_1 x}$
Exponential curve	$y = a \cdot b^x$
Geometric curve	$y = a \cdot x^b$
Gompertz Curve	$y = pq^{b^x}$
Logistic Curve	$y = \frac{1}{ab^x + c}$

To decide which curve to be used, it is helpful to obtain scatter diagrams of the transformed variables. For example, if a scatter diagram of log yversus x shows a linear relationship, then the equation is of the $y = a \cdot b^x$, while if log y versus log x shows a linear relationship, the equation is of the form $y = a \cdot x^b$.

12.5 FREEHAND METHOD OF CURVE FITTING

Individual judgment can often be used to draw an approximating curve to fit a set of data. This is called a freehand method of curve fitting. If the type of equation of this curve is known, it is possible to obtain the constants in the equation by choosing as many points on the curve as there are constants in the equation. For example, if the curve is a straight line, two points are necessary; if it is a parabola, three points are necessary. The method has the disadvantage that different observers will obtain different curves and equations.

12.6 THE STRAIGHT LINE METHOD

The simplest type of the approximating curve is a straight line, whose equation can be written

$$y = a + bx$$

Given any two points $(x_1, y_1) \& (x_2, y_2)$ on the line, the constants a and b can be determined. The resulting equation of the straight line can be written as

$$y - y_1 = m(x - x_1), m = \frac{y_2 - y_1}{x_2 - x_1}$$

where m is known as the slope of the line.

When the equation is written in the form of y = a + bx, the constant 'b' denotes the slope m. The constant 'a' is the value of y when x = 0 is called the y-intercept.

12.7 LEAST SQUARE CURVE FITTING

We can find the trend curve by fitting a mathematical equation. The method is more precise and can be used even for forecasting. We can fit either a straight line or a curve to the given data. We fit a straight line $y_c = a + bx$, where a and b are constants. We determine the constants a and b so that the following conditions are fulfilled:

i) The sum of the deviations of all the values of y from their trend values is zero, when, deviations above the line are given positive sign and deviations below, negative i.e if y_c is the trend value obtained from the trend line, and y is the actual value in the data.

$$\sum (y - y_c) = 0$$

ii) The sum of the squares of the deviations is the least, i.e.,

 $\Sigma(y - y_c)^2$ is minimum.

The method gets the name 'least square method' because of this second property. This is also called *line of best fit*. In a sense this line is like arithmetic mean since arithmetic mean is a single value possessing the above two properties.

Remark: When the value of x is given in terms of years or value of x is big then we apply the following technique to get the values of x:

- 1) If the value of n is odd then we take the value of central most observation i.e value of $\frac{n+1}{2}$ th observation as 0 and we add 1 as we move downward and we subtract 1 as we move upwards.
- 2) If the value if n is even then we take the value of $\frac{n}{2}$ th observation as -1 and we add 2 as we move downward and subtract 2 as we move upwards.

12.7.1 Straight Line:

Suppose o two variables x and y, n pairs of observations, say $(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$ are available and we want to fit a linear curve of the form y = a + bx to the data. Then according to the least squares method we have to find a and b such that $\sum (y - y_i)^2$ is minimum where $y_i = a + bx_i$.

Thus, we have to minimize $D = \sum (y - a - bx)$. The first order conditions are $\frac{\delta D}{\delta a} = 0$ and $\frac{\delta D}{\delta b} = 0$

which are known as *normal equations* and they are given by

$$\sum y = an + b\sum x$$
 and $\sum xy = a\sum x + b\sum x^2$

It can be easily verified that the second order condition for minima are satisfied and therefore the best choice of a and b can be made by finding their values satisfying the normal equations. This equations will be used to fit a linear curve to the given data.

Eg: 1) Fit a straight line of the form y = a + bx using least square method

Х	0	1	2	3	4
у	1	2.9	4.8	6.7	8.6

Solution: n = 5

	Х	У	xy	\mathbf{x}^2
	0	1	0	0
	1	2.9	2.9	1
	2	4.8	9.6	4
	3	6.7	20.1	9
	4	8.6	34.4	16
total	10	24.0	67.0	30

The straight line equation is given by y = a + bx. To find values of a and b we solve the following equations

 $\sum y = an + b\sum x, \sum xy = a\sum x + b\sum x^2 \quad \dots \quad (1)$ by substituting the values in eq (1), we get

$$24 = 5a + 10b - - - -(2)$$

67 = 10a + 30b - - - -(3)

multiplying eq (2) by 2 we get 48 = 10a + 20b - - - -(4)subtracting eq (4) by eq (3), we get $19 = 10b \Rightarrow b = 1.9$ substituting b = 1.9 in eq (2) gives a = 1 $\therefore y = 1 + 1.9x$

is the required straight line.

Eg: 2) Fit a straight line of the form $y = a_0 + a_1 x$

ſ	X	1	2	3	4	6	8
	у	2.4	3.1	3.5	4.2	5	6

Solution: n = 6

	Х	у	xy	\mathbf{x}^2
	1	2.4	2.4	1
	2	3.1	6.2	4
	3	3.5	10.5	9
	4	4.2	16.8	16
	6	5	30	36
	8	6	48	64
total	24	24.2	113.9	130

The straight line equation is given by $y = a_0 + a_1 x$. To find values of a_0 and a_1 we solve the following equations

 $\sum y = a_0 n + a_1 \sum x, \sum xy = a_0 \sum x + a_1 \sum x^2 \quad \dots \quad (1)$ by substituting the values in eq (1), we get

$$24.2 = 6a_0 + 24a_1 - - - (2)$$

$$113.9 = 24a_0 + 130a_1 - - - (3)$$

multiplying eq (2) by 4, we get $96.8 = 24a_0 + 96a_1 - - (4)$ subtracting eq (4) from eq (3), we get $17.1 = 34a_1 \Rightarrow a_1 = 0.5029$

subs $a_1 = 0.5029$ in (2) gives $a_0 = 2.0217$

$$\therefore y = 2.0217 + 0.5029x$$

is the required straight line.

Eg: 3) Fit a straight line using least square method and estimate the exchange rate for the year 1993-94 and 1984-85.

year	1985-	1986-	1987-	1988-	1989-	1990-	1991-
-	86	87	88	89	90	91	92
Exchange rate	12.24	12.78	12.97	14.48	16.65	17.94	24.47

Solution: n = 7

The straight line equation we take is y = a + bx. To find values of a and b we solve the following equations

	year	ex. rate(y)	Х	xy	\mathbf{x}^2
	1985-86	12.24	-3	-36.72	9
	1986-87	12.78	-2	-25.56	4
	1987-88	12.97	-1	-12.97	1
	1988-89	14.48	0	0	0
	1989-90	16.65	1	16.65	1
	1990-91	17.94	2	35.88	4
	1991-92	24.47	3	73.41	9
total		111.53	0	50.69	28

 $\sum y = an + b\sum x, \sum xy = a\sum x + b\sum x^2 \qquad \dots \qquad (1)$

by substituting the values in eq (1), we get

$$111.53 = 7a - - - (2)$$

$$50.69 = 28b - - - (3)$$

from eq (2), we get $a = \frac{111.53}{7} = 15.9328$

from eq (3), we get $b = \frac{50.69}{28} = 1.8104$

$$\therefore y = 15.9328 + 1.8104x - - - -(4)$$

is the required straight line.

To the get exchange rate for the year 1993-94, we subs x = 5 in eq (4)

 $\therefore y = 15.9328 + 1.8104(5) = 24.9848$

To the get the exchange rate for the year 1984-85, we subs x = -4 in eq (4) $\therefore y = 15.9328 + 1.8104(-4) = 8.6912$

12.7.2 Parabola:

Suppose we want to fit a second degree curve called parabola of the form $y = a + bx + cx^2$ to *n* pairs of observations $(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$. Then according to the least squares method we have to minimize $D = \sum (y - a - bx - cx^2)^2$ with respect to a, b and c. The first order conditions are given by

$$\frac{\delta D}{\delta a} = 0, \frac{\delta D}{\delta b} = 0 \text{ and } \frac{\delta D}{\delta c} = 0$$

Applying these we get,

$$\sum y = na + b\sum x + c\sum x^{2}$$
$$\sum xy = a\sum x + b\sum x^{2} + c\sum x^{3}$$
$$\sum x^{2}y = a\sum x^{2} + b\sum x^{3} + c\sum x^{4}$$

Eg: 1) Fit a parabola of the form $y = a + bx + cx^2$ using least square method

Х	0	1	2
у	1	6	17

Solution: n=3

	Х	у	\mathbf{x}^2	x^3	x ⁴	xy	x^2y
	0	1	0	0	0	0	0
	1	6	1	1	1	6	6
	2	17	4	8	16	34	68
total	3	24	5	9	17	40	74

The equation of the parabola is given by $y = a + bx + cx^2$. To find values of *a*, *b* and *c* we solve the following equations

$$\begin{split} \sum y &= na + b \sum x + c \sum x^2, \qquad \sum xy = a \sum x + b \sum x^2 + c \sum x^3, \\ \sum x^2 y &= a \sum x^2 + b \sum x^3 + c \sum x^4 \quad - - - - (*) \end{split}$$

subs values in eq (*), we get

*), we get

$$24 = 3a + 3b + 5c - - - -(1)$$

 $40 = 3a + 5b + 9c - - - -(2)$
 $74 = 5a + 9b + 17c - - - -(3)$

Subtracting eq (1) by (2), we get $16 = 2b + 4c \Rightarrow 8 = b + 2c - - - (4)$ mul eq (2) by 5, we get 200 = 15a + 25b + 45c - - - (5)mul eq (3) by 3, we get 222 = 15a + 27b + 51c - - - (6)subtractting eq (5) by (6), we get 22 = 2b + 6c $\Rightarrow 11 = b + 3c - - - (7)$ subtracting eq (4) by (7), we get 3 = csub c = 3 in eq (4) we get b = 2sub b = 2, c = 3 in (1), we get a = 1 $\therefore y = 1 + 2x + 3x^{2}$ is the required equation of the parabola.

Eg: 2) Fit a para	abola of the f	form $y = a +$	$bx + cx^2$

0 / 1								
year	1989	1990	1991	1992	1993	1994	1995	1996
no.of students	15	17	20	25	30	31	30	32
A 1 C	1.1	<u> </u>	1 ()	.1	2000	0 100	-	

Also find the no. of students for the year 2000 & 1987.

Solution: n = 8

year	no. of students(y)	Х	x ²	x ³	x ⁴	xy	x ² y
1989	15	-7	49	-343	2401	-105	735
1990	17	-5	25	-125	625	-85	425
1991	20	-3	9	-27	81	-60	180
1992	25	-1	1	-1	1	-25	25

	1993	30	1	1	1	1	30	30
	1994	31	3	9	27	81	93	279
	1995	30	5	25	125	625	150	750
	1996	32	7	49	343	2401	224	1568
total		200	0	168	0	6216	222	3992

The equation of the parabola is given by $y = a + bx + cx^2$. To find values of *a*, *b* and *c* we solve the following equations

$$\begin{split} \sum y &= na + b \sum x + c \sum x^2, \qquad \sum xy = a \sum x + b \sum x^2 + c \sum x^3, \\ \sum x^2 y &= a \sum x^2 + b \sum x^3 + c \sum x^4 \quad ----(*) \end{split}$$

subs values in eq (*), we get

$$200 = 8a + 168c - - - -(1)$$

$$222 = 168b - - - -(2)$$

$$3992 = 168a + 6216c - - - -(3)$$
from (2), we get $b = \frac{222}{168} = 1.3214$
mul (1)by 21, we get 4200 = $168a + 3528c - - -(4)$
sub (4)by (3), we get $208 = -2688c \Rightarrow c = -\frac{208}{2688} = -0.0774$
sub $c = -0.0774$ in (1), we get $a = \frac{200 - 168(-0.0774)}{8} = 26.6254$
 $\therefore y = 26.6254 + 1.3214x - 0.0774x^2 - - - -(4)$
is the required equation of the parabola

is the required equation of the parabola.

To get the no. of students for the year 2000 we substitute x = 15 in (4)

$$\therefore y = 26.6254 + 1.3214(15) - 0.0774(15)^2 = 29.0314$$

To get the no. of students for the year 1987 we substitute x = -11 in (4)

$$\therefore y = 26.6254 + 1.3214(-11) - 0.0774(-11)^2 = 2.7246$$

Eg: 3) Fit a parabola of the form $y = a + bx + cx^2$

year	1985	1986	1987	1988	1989			
milk (in 100 litres)	20	25	27	35	38			
Find the graduation of will for the year 1082 and 1005								

Find the production of milk for the year 1982 and 1995.

Solution: n =5

year	milk	Х	\mathbf{x}^2	x^{3}	x^4	xy	x^2y
	(y)						
1985	20	-2	4	-8	16	-40	80
1986	25	-1	1	-1	1	-25	25
1987	27	0	0	0	0	0	0

	1988	35	1	1	1	1	35	35
	1989	38	2	4	8	16	76	152
total		145	0	10	0	34	46	292

The equation of the parabola is given by $y = a + bx + cx^2$. To find values of *a*, *b* and *c* we solve the following equations

$$\sum y = na + b\sum x + c\sum x^2, \qquad \sum xy = a\sum x + b\sum x^2 + c\sum x^3$$
$$\sum x^2 y = a\sum x^2 + b\sum x^3 + c\sum x^4 \quad ----(*)$$

subs values in eq (*), we get

145 = 5a + 10c - -(1) $46 = 10b - -(2) \implies b = 4.6$ 292 = 10a + 34c - -(3)multiply eq (1) by 2, we get 290 = 10 a + 20c - -(4)subtracting eq (4) from eq (3), we get $2 = 14c \Rightarrow c = \frac{2}{14} = 0.1428$ substituting c = 0.1428 in eq (1), we get a = 28.7144 $\therefore y = 28.7144 + 4.6x + 0.1428x^2 - - - (5)$ is the required equation of the parabola. To get the production of milk for the year 1995, we put x = 8 in (5) $\therefore y = 28.7144 + 4.6(8) + 0.1428(8)^2 = 74.6536$ To get the production of milk for the year 1982, we put x = -5 in (5) $\therefore y = 28.7144 + 4.6(-5) + 0.1428(-5)^2 = 9.2844$

12.7.3 Non-Linear Relationship: Fitting of a power Curve:

Suppose we want to fit a curve whose equation is of the form $y = a \cdot x^b$ to variables x and y on which n paired observations are available. Then we first rewrite the equation of the curve as

$$\log y = \log a + b \log x$$

which can be written in the linear form Y = A + bX where $Y = \log y$, A =log a and $X = \log x$. Therefore this curve can be fitted by the method of least squares solving the following normal equations.

$$\Sigma Y = nA + b\Sigma X$$
 , $\Sigma XY = A\Sigma X + b\Sigma X^2$

Once we get the value of A, we can find a = antilog (A). The method of fitting is explained in the following example.

Eg: 1) Fit a curve of the form $y = a \cdot x^b$ for the following data

Х	1	2	4	5	7	10
у	2.1	4.9	20.8	32.7	60	131

Solution: n = 6

We can write the given curve in the form of Y = A + bX where $Y = \log y$, $A = \log a$, $X = \log x$

	Х	у	$X = \log x$	$Y = \log y$	XY	X^2
	1	2.1	0	0.3222	0	0
	2	4.9	0.3010	0.6902	0.2078	0.0906
	4	20.8	0.6021	1.3181	0.7936	0.3625
	5	32.7	0.6990	1.5142	1.0586	0.4886
	7	60	0.8451	1.7782	1.5028	0.7142
	10	131	1	2.1173	2.1173	1
Total			3.4472	7.7405	5.6801	2.6559

The given curve is of the form $y = ax^b$, therefore the normal equations are given by

 $\sum Y = nA + b\sum X$, $\sum XY = A\sum X + b\sum X^2 - - - (1)$

substituting the values in eq (1), we get

$$7.7405 = 6A + 3.4472b - - - (2)$$

$$5.6801 = 3.4472A + 2.6559b - - - - (3)$$

multiplying eq (1) by 3.4472 and eq (2) by 6, then subtracting the two eq., we get

$$4.0522b = 7.3975 \Rightarrow b = \frac{7.3975}{4.0522} = 1.8256$$

substituting b = 1.8256 in eq (1), we get a = 0.02412

Now a = antilog(A) = antilog(0.2412) = 1.7426

 $\therefore y = 1.7426x^{1.8256}$

is the required equation.

Fitting of the curve $y = ae^{bx}$

We first try to convert the equation $y = ae^{bx}$ to a linear equation by applying logarithms on both sides with respect to base 10. By applying logarithms on both sides, we get

$$\log y = \log a + bx \log_{10} e$$

Let Y = log y, A = log a, $log_{10} e = 0.4343b$. Then the equation of the curve reduces to a linear equation of the form

$$Y = A + Bx$$

By using the least square method the constants A and b can be found by solving the normal equations

$$\sum Y = nA + B\sum x$$
, $\sum xY = A\sum x + B\sum x^2$

Once A and B are found, we can find a = antilog (A) and b = $\frac{B}{0.4343}$

Eg: 1) Fit a curve of the form $y = ae^{bx}$ for the following data

х	0	2	4	6	8
у	3	55	1095	22000	442000

Solution : n =5

	Х	У	$Y = \log y$	xY	x^2
	0	3	0.4771	0	0
	2	55	1.7404	3.4808	4
	4	1095	3.0394	12.1576	16
	6	22000	4.3424	26.0544	36
	8	442000	5.6454	45.1632	64
Total	20		15.2447	86.8560	120

The given equation is of the form $y = ae^{bx}$, therefore the normal equations are given by

$$\sum Y = nA + B\sum x$$
, $\sum xY = A\sum x + B\sum x^2$ $---(1)$

substituting the values in eq (1), we get

15.2447 = 5A + 20B - - - -(2)

$$86.8560 = 20A + 120B - - - - (3)$$

multiplying eq (1) by 4 and subtracting it from eq (2), we get

$$25.8772 = 40B \Rightarrow B = \frac{25.8772}{40} = 0.6469$$

substituting B = 0.6469 in eq (1), we get

$$15.2447 = 5A + 20(0.6469) \Rightarrow A = \frac{2.3067}{5} = 0.4613$$

Now a = antilog (A) = antilog(0.4613) = 2.892

$$b = \frac{B}{0.4343} = \frac{0.6469}{0.4343} = 1.4895$$

 $\therefore y = 2.892e^{1.4895x}$

is the required equation.

12.8 REGRESSION

Often, on the basis of sample data, we wish to estimate the value of a variable Y corresponding to a given value of a variable x. This can be accomplished by estimating the value of y from a least-squares curve that fits the sample data. The resulting curve is called a regression curve of y on x, since y is estimated from x.

If we want to estimate the value of x from a given value of y, we would use a regression curve of x on y, which amounts to interchanging the variables in the scatter diagram so that x is the dependent variable and y is the independent variable. This is equivalent to replacing the vertical deviations in the definition of the least-squares curve with horizontal deviations.

In general, the regression line or curve of y on x is not the same as the regression line or curve of x on y.

12.9 APPLICATIONS TO TIME SERIES

If the independent variable x is time, the data show the values of y at various times. Data arranged according to time are called time series. The regression line or curve of y on x in this case is often called atrend line or trend curve and is often used for purposes of estimation, prediction, or forecasting.

12.10 PROBLEMS INVOLVING MORE THAN TWO VARIABLES

Problems involving more than two variables can be treated in a manner analogous to that for two variables. For example, there may be a relationship between the three variables X, Y, and Z that can be described by the equation

$$z = a + bx + cy$$

which is called a linear equation in variables x, y and z.

In a three dimensional rectangular coordinate system this equation represents a plane, and the actual sample points $(x_1, y_1, z_1), (x_2, y_2, z_2), \dots, (x_N, y_N, z_N)$ may scatter not too far from this plane which we call an *approximating plane*.

By extension of the least square method, we can talk about a least square plane approximating the data. If we are estimating z from the given values of x and y, this would be called a regression plane of z on x and y. The normal equations corresponding to the least square plane are given by

$$\sum z = aN + b\sum x + c\sum y$$

$$\sum xz = a\sum x + b\sum x^{2} + c\sum xy$$

$$\sum yz = a\sum y + b\sum xy + c\sum y^{2}$$

Problems involving the estimation of a variable from two or more variables are called problems of multiple regression which is not there in the syllabus.

12.11 SUMMARY

In this chapter we learnt about curve fitting, various equations of approximating curves, the straight line method. We also learnt about the

fitting of linear and non-linear curves using the technique of least square which minimizes the sum of the square of the error.

12.12 EXERCISES

1)	Fit a stra	ight line of th	ght line of the form $y = ax + b$ using least square method								
	Х	0	1	2	3						
	у	2	5	8	11						

2) Fit a straight line of the form y = a + bx

Х	10	12	13	16	17	20	25
у	19	22	24	27	29	33	37

3) Fit a straight line of the form y = a + bx

	•			•					
Х	1	3	5	7	9	11	13	15	17
у	10	15	20	27	31	35	30	35	40

Also estimate y when x = 21.

4) Fit a parabola of the form $y = a + bx + cx^2$ using least square method

Х	0	1	2	3	4
у	1	0	3	10	21

5) Fit a parabola of the form $y = a + bx + cx^2$

Х	0	1	2	3	4
у	1	1.8	1.3	2.5	6.3

6) Fit a parabola using least square method $y = ax + bx^2$

X	1	2	3	4	5
у	1.8	5.1	8.9	14.1	19.8

7) Fit a curve of the form $y = ax^b$ for the following data

Х	1	2	3	4
у	0.7	0.86	0.97	1.06

8) Fit a curve of the form $y = ae^{bx}$ for the following

Х	0	2	4
У	5.012	10	31.62

9) Fit a curve of the form $y = ae^{bx}$ for the following

Х	1	2	3	4	5
у	1.230	2.042	3.162	3.981	5.624

10) The number y of bacteria per unit volume present in a culture after x hours is given in the following table. Fit a of the form $y = ab^x$ using least square method for the following data.

Х	0	1	2	3	4	5	6
У	30	45	60	90	130	190	275

11) Fit a curve of the form $y = ab^x$ for the following

Х	1	2	3
У	8.3	15.4	33.1

12) The population of a state at ten yearly intervals is given below. Fit a curve of the form $y = ab^x$ using least square method and also estimate the population for the year 1961.

year	1881	1891	1901	1911	1921	1931	1941	1951
population in millions	3.9	5.3	7.3	9.6	12.9	17.1	23.2	30.5

13) Fit a straight line of the form y = a + bx for the following data

x	5	4	3	2	1
у	1	2	3	4	5

14) Fit a straight line of the form y = a + bx for the following data

х	3	5	7	9	11
у	2.3	2.6	2.8	3.2	3.5

15) Fit a straight line to the following data on production

Year	1996	1997	1998	1999	2000
Production	40	50	62	58	60

16) Fit a straight line to the following data on profit

	-							
Year	1992	1993	1994	1995	1996	1997	1998	1999
Profit	38	40	65	72	69	60	87	95

17) Fit a second degree parabolic equation of the form $y = a + bx + cx^2$

x	12	10	8	6	4	2
у	6	5	4	3	2	1

12.14 SOLUTION TO EXERCISES

Q. No.	Solution	Q. No.	Solution
1	2, 3	2	a = 7.7044, b = 1.213
3	a=11.43, b=1.73	4	1, -3, 2
5	1.42, -1.07, 0.55	6	a= 1.44 , b = 0.51
7	$y = 4.642e^{0.46x}$	8	$y = 4.642e^{0.46x}$
9	$y = 0.7762e^{0.472x}$	10	$y = 30(1.306)^x$
11	$y = 8.099(1.997)^x$	12	$y = 11.074(1.0298)^{x-1916}$
13	6, -1	14	1.83, 0.15
15	54, 4.8	16	62.0833, 7.3333
17	0, 0.5, 0		

12.15 LOGARITHM TABLES

LOGARITHM TABLES

LOGARITHMS

185

				_		_	_	_	_				M	əan	Dif	fere	nce		
	0	1	2	3	4	5	6	7	8	9	1	2	3	4	5	6	7	8	9
10	0000	0043	0086	0128	0170	0212	0253	0294	0334	0374	4	8	12	17	21	25	29	33	37
11 12 13 14 15	0414 0792 1139 1461 1761	0453 0828 1173 1492 1790	0492 0864 1206 1523 1818	0531 0899 1239 1553 1847	0569 0934 1271 1584 1875	0607 0969 1303 1614 1903	0645 1004 1335 1644 1931	0682 1038 1367 1673 1959	0719 1072 1399 1703 1987	0755 1106 1430 1732 2014	433333	8 7 6 6 6	11 10 10 9 8	15 14 13 12 11	19 17 16 15 14	23 21 19 18 17	26 24 23 21 20	30 28 26 24 22	34 31 29 27 25
16 17 18 19 20	2041 2304 2553 2788 3010	2068 2330 2577 2810 3032	2095 2355 2601 2833 3054	2122 2380 2625 2856 3075	2148 2405 2648 2878 3096	2175 2430 2672 2900 3118	2201 2455 2695 2923 3139	2227 2480 2718 2945 3160	2253 2504 2742 2967 3181	2279 2529 2765 2989 3201	3 2 2 2 2 2 2	5 5 4 4	8 7 7 6	11 10 9 9 8	13 12 12 11 11	16 15 14 13 13	18 17 16 16 15	21 20 19 18 17	24 22 21 20 19
21 22 23 24 25	3222 3424 3617 3802 3979	3243 3444 3636 3820 3997	3263 3464 3655 3838 4014	3284 3483 3674 3856 4031	3304 3502 3692 3874 4048	3324 3522 3711 3892 4065	3345 3541 3729 3909 4082	3365 3560 3747 3927 4099	3385 3579 3766 3945 4116	3404 3598 3784 3962 4133	2 2 2 2 2 2 2 2	4 4 4 3	6 6 6 5 5	8 8 7 7 7	10 10 9 9 9	12 12 11 11 10	14 14 13 12 12	16 15 15 14 14	18 17 17 16 15
26 27 28 29 30	4150 4314 4472 4624 4771	4166 4330 4487 4639 4786	4183 4346 4502 4654 4800	4200 4362 4518 4669 4814	4216 4378 4533 4683 4829	4232 4393 4548 4698 4843	4249 4409 4564 4713 4857	4265 4425 4579 4728 4871	4281 4440 4594 4742 4886	4298 4456 4609 4757 4900	2 2 1 1	3 3 3 3 3	5 5 5 4 4	7 6 6 6	8 8 7 7	10 9 9 9	11 11 11 10 10	13 13 12 12 11	15 14 14 13 13
31 32 33 34 35	4914 5051 5185 5315 5441	4928 5065 5198 5328 5453	4942 5079 5211 5340 5465	4955 5092 5224 5353 5478	4969 5105 5237 5366 5490	4983 5119 5250 5378 5502	4997 5132 5263 5391 5514	5011 5145 5276 5403 5527	5024 5159 5289 5416 5539	5038 5172 5302 5428 5551	1 1 1 1	3 3 3 2	4 4 4 4 4	6 5 5 5 5	7 7 6 6	8 8 8 7	10 9 9 9	11 11 10 10 10	12 12 12 11 11
36 37 38 39 40	5563 5682 5798 5911 6021	5575 5694 5809 5922 6031	5587 5705 5821 5933 6042	5566 5717 5832 5944 6053	5611 5729 5843 5955 6064	5623 5740 5855 5966 6075	5635 5752 5866 5977 6085	5647 5763 5877 5988 6096	5658 5775 5888 5999 6107	5670 5786 5899 6010 6117	1 1 1 1	2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	4 3 3 3 3 3	5 5 4 4	6 6 5 5	7 7 7 6	8 8 8 8	10 9 9 9	11 10 10 10 10
41 42 43 44 45	6128 6232 6335 6435 6532	6138 6243 6345 6345 6542	6149 6253 6355 6454 6551	6160 6263 6365 6464 6561	6170 6274 6375 6474 6571	6180 6284 6385 6484 6580	6191 6294 6395 6493 6590	6201 6304 6405 6503 6599	6212 6314 6415 6513 6609	6222 6325 6425 6522 6618	1 1 1 1	2 2 2 2 2 2 2 2 2	0 0 0 0 0 0 0 0 0	4 4 4 4	5 5 5 5 5 5	6 6 6 6	7 7 7 7 7	8 8 8 8	9 9 9 9
46 47 48 49 50	6628 6721 6812 6902 6990	6637 6730 6821 6911 6998	6646 6739 6830 6920 7007	6656 6749 6839 6928 7016	6665 6758 6848 6937 7024	6675 6767 6857 6946 7033	6684 6776 6866 6955 7042	6693 6785 6875 6964 7050	6702 6794 6884 6972 7059	6712 6803 6893 6981 7067	1 1 1 1	2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3	4 4 4 3	5 5 4 4 4	6 5 5 5 5 5 5	7 6 6 6	7 7 7 7 7	8 8 8 8 8
51 52 53 54	7076 7160 7243 7324	7084 7168 7251 7332	7093 7177 7259 7340	7101 7185 7267 7348	7110 7193 7275 7356	7118 7202 7284 7364	7126 7210 7292 7372	7135 7218 7300 7380	7143 7226 7308 7388	7152 7235 7316 7396	1 1 1	2 2 2 2 C	3222	333	4 4 4 4	5 5 5 5	6 6 6 7	7 7 6 6	8 7 7 7
	0	1	2	3	4	Э	6	1	8	9	11	2	3	4	5	Б.	1	2	- 9

	0	4	•	•	4	e	6	-		•			Me	an	Diffe	ərən	сө		
	U	_ '	2	3	4	5	0		•	3	1	2	3	4	5	6	7	8	9
.00	1000	1002	1005	1007	1009	1012	1014	1016	1019	1021	0	0	1	1	1	1	2	2	2
.01 .02 .03 .04 .05	1023 1047 1072 1096 1122	1026 1050 1074 1099 1125	1028 1052 1076 1102 1127	1030 1054 1079 1104 1130	1033 1057 1081 1107 1132	1035 1059 1084 1109 1135	1038 1062 1086 1112 1138	1040 1064 1089 1114 1140	1042 1067 1091 1117 1143	1045 1069 1094 1119 1146	0 0 0 0	0 0 1 1	1 1 1 1	1 1 1 1	1 1 1 1	1 1 2 2	2 2 2 2 2 2	2 2 2 2 2 2 2	2 2 2 2 2 2 2 2
.06 .07 .08 .09 .10	1148 1175 1202 1230 1259	1151 1178 1205 1233 1262	1153 1180 1208 1236 1265	1156 1183 1211 1239 1268	1159 1186 1213 1242 1271	1161 1189 1216 1245 1274	1164 1191 1219 1247 1276	1167 1194 1222 1250 1279	1169 1197 1225 1253 1282	1172 1199 1227 1256 1285	0 0 0 0	1 1 1 1	1 1 1 1	1 1 1 1	1 1 1 1	2 2 2 2 2	2 2 2 2 2 2 2 2	2 2 2 2 2 2 2	2 2 3 3 3
.11 .12 .13 .14 .15	1288 1318 1349 1380 1413	1291 1321 1352 1384 1416	1294 1324 1355 1387 1419	1297 1327 1358 1390 1422	1300 1330 1361 1393 1426	1303 1334 1365 1396 1429	1306 1337 1368 1400 1432	1309 1340 1371 1403 1435	1312 1343 1374 1406 1439	1315 1346 1377 1409 1442	0 0 0 0	1 1 1 1	1 1 1 1	1 1 1 1	2 2 2 2 2	2 2 2 2 2	2 2 2 2 2 2 2 2	2 2 3 3 3	3 3 3 3
.16 .17 .18 .19 .20	1445 1479 1514 1549 1585	1449 1483 1517 1552 1289	1452 1486 1521 1556 1592	1455 1489 1524 1560 1596	1459 1493 1528 1563 1600	1462 1496 1531 1567 1603	1466 1500 1535 1570 1607	1469 1503 1538 1574 1611	1472 1507 1542 1578 1614	1476 1510 1545 1581 1618	0 0 0 0 0	1 1 1 1	1 1 1 1	1 1 1 1	2 2 2 2 2 2 2 2	2 2 2 2 2	2 2 2 3 3	3 3 3 3	3 3 3 3 3
.21 .22 .23 .24 .25	1622 1660 1698 1738 1778	1626 1663 1702 1742 1782	1629 1667 1706 1746 1786	1633 1671 1710 1750 1791	1637 1675 1714 1754 1795	1641 1679 1718 1758 1799	1644 1683 1722 1762 1803	1648 1687 1726 1766 1807	1652 1690 1730 1770 1811	1656 1694 1734 1774 1816	0 0 0 0 0	1 1 1 1	1 1 1 1	2 2 2 2 2 2 2	2 2 2 2 2 2 2 2 2	22222	3 3 3 3 3	3 3 3 3	3 3 4 4 4
.26 .27 .28 .29 .30	1820 1862 1905 1950 1995	1824 1866 1910 1954 2000	1828 1871 1914 1959 2004	1832 1875 1919 1963 2009	1837 1879 1923 1968 2014	1841 1884 1928 1972 2018	1845 1888 1932 1977 2023	1849 1892 1936 1982 2028	1854 1897 1941 1986 2032	1858 1901 1945 1991 2037	0 0 0 0	1 1 1 1	1 1 1 1	2 2 2 2 2 2 2	2 2 2 2 2 2 2	3 3 3 3 3	3 3 3 3 3	3 3 4 4 4	4 4 4 4
.31 .32 .33 .34 .35	2042 2089 2138 2188 2239	2046 2094 2143 2193 2244	2051 2099 2148 2198 2249	2056 2104 2153 2203 2254	2061 2109 2158 2208 2259	2065 2113 2163 2213 2265	2070 2118 2168 2218 2270	2075 2123 2173 2223 2275	2080 2128 2178 2228 2280	2084 2133 2183 2234 2286	0 0 1 1	1 1 1 1	1 1 2 2	2 2 2 2 2 2 2 2	2 2 2 3 3	3 3 3 3 3	3 3 4 4	4 4 4 4	4 4 5 5
.36 .37 .38 .39 .40	2291 2344 2399 2455 2512	2296 2350 2404 2460 2518	2301 2355 2410 2466 2523	2307 2360 2415 2472 2529	2312 2366 2421 2477 2535	2317 2371 2427 2483 2541	2323 2377 2432 2489 2547	2328 2382 2438 2495 2553	2333 2388 2443 2500 2559	2339 2393 2449 2506 2564	1 1 1 1	1 1 1 1	2 2 2 2 2 2 2	2 2 2 2 2 2 2	3 3 3 3 3	3 3 3 4	4 4 4 4	4 4 5 5	5 5 5 5 5
.41 .42 .43 .44 .45	2570 2630 2692 2754 2818	2576 2636 2698 2761 2825	2582 2642 2704 2767 2831	2588 2649 2710 2773 2838	2594 2655 2716 2780 2844	2600 2661 2723 2786 2851	2606 2667 2729 2793 2858	2612 2673 2735 2799 2864	2618 2679 2742 2805 2871	2624 2685 2748 2812 2877	1 1 1 1	1 1 1 1	2 2 2 2 2 2	2 2 3 3	3 3 3 3 3	4 4 4 4	4 4 4 5	5 5 5 5 5	5 6 6 6
.46 .47 .48 .49	2884 2951 3020 3090	2891 2958 3027 3097	2897 2965 3034 3105	2904 2972 3041 3112	2911 2979 3048 3119	2917 2985 3055 3126	2924 2992 3062 3133	2931 2999 3069 3141	2938 3006 3076 3148	2944 3013 3083 3155	1 1 1	1 1 1	2 2 2 2	3 3 3 3	3 3 4 4	4 4 4 4	5 5 5 5	5 5 6 6	6 6 6
	U	1	2	3	4	5	5	7	8	9	1	2	3	4	5	6	1	8	9

ANTILOGARITHMS

			_	_					000				Мө	an	Diff	ere	nce		
	0	1	2	3	4	5	6	7	8	9	1	2	3	4	5	6	7	8	9
.50	3162	3170	3177	3184	3192	3199	3206	3214	3221	3228	1	1	2	з	4	4	5	6	7
.51 .52 .53 .54 .55	3236 3311 3388 3467 3548	3243 3319 3396 3475 3556	3251 3327 3404 3483 3565	3258 3334 3412 3491 3573	3266 3342 3420 3499 3581	3273 3350 3428 3508 3589	3281 3357 3436 3516 3597	3289 3365 3443 3524 3606	3296 3373 3451 3532 3614	3304 3381 3459 3540 3622	1 1 1 1	2 2 2 2 2 2	2 2 2 2 2	3 3 3 3 3	4 4 4 4	5 5 5 5 5 5 5	5 5 6 6	6 6 6 7	7 7 7 7 7
.56 .57 .58 .59 .60	3631 3715 3802 3890 3981	3639 3724 3811 3899 3990	3648 3733 3819 3908 3999	3656 3741 3828 3917 4009	3664 3750 3837 3926 4018	3673 3758 3846 3936 4027	3681 3767 3855 3945 4036	3690 3776 3864 3954 4046	3698 3784 3873 3963 4055	3707 3793 3882 3972 4064	1 1 1 1	2 2 2 2 2 2 2	3 3 3 3	3 3 4 4 4	4 4 4 5 5	ខេលល	6 6 6 6	7 7 7 7 7	8 8 8 8 8
.61 .62 .63 .64 .65	4074 4169 4266 4365 4467	4083 4178 4276 4375 4477	4093 4188 4285 4385 4487	4102 4198 4295 4395 4498	4111 4207 4305 4406 4508	4121 4217 4315 4416 4519	4130 4227 4325 4426 4529	4140 4236 4335 4436 4539	4150 4246 4345 4446 4550	4159 4256 4355 4457 4560	1 1 1 1	2 2 2 2 2 2 2	3 3 3 3	4 4 4 4	5 5 5 5 5 5 5	6 6 6 6 6	7 7 7 7 7	8 8 8 8	<i>0</i> 0 0 0 0 0
.66 .67 .68 .69 .70	4571 4677 4786 4898 5012	4581 4688 4797 4909 5023	4592 4699 4808 4920 5035	4603 4710 4819 4932 5047	4613 4721 4831 4943 5058	4624 4732 4842 4955 5070	4634 4742 4853 4966 5082	4645 4753 4864 4977 5093	4656 4764 4875 4989 5105	4667 4775 4887 5000 5117	1 1 1 1	2 2 2 2 2 2 2 2	3 3 3 3 4	4 4 4 5 5	5 5 6 6 6	6 7 7 7 7	7 8 8 8	9 9 9 9	10 10 10 10 11
.71 .72 .73 .74 .75	5129 5248 5370 5495 5623	5140 5260 5383 5508 5636	5152 5272 5395 5521 5649	5164 5284 5408 5534 5662	5176 5297 5420 5546 5675	5188 5309 5433 5559 5689	5200 5321 5445 5572 5702	5212 5333 5458 5585 5715	5224 5346 5470 5598 5728	5236 5358 5483 5610 5741	1 1 1 1	22333	4 4 4 4 4	5 5 5 5 5 5	6 6 6 7	7 7 8 8 8	8 9 9 9	10 10 10 10 10	11 11 11 12 12
.76 .77 .78 .79 .80	5754 5888 6026 6166 6310	5768 5902 6039 6180 6324	5781 5916 6053 6194 6339	5794 5929 6067 6209 6353	5808 5943 6081 6223 6368	5821 5957 6095 6237 6383	5834 5970 6109 6252 6397	5848 5984 6124 6266 6412	5861 5998 6138 6281 6427	5875 6012 6152 6295 6442	1 1 1 1	33333	4 4 4 4	5 5 6 6	7 7 7 7 7	8 8 9 9	9 10 10 10 10	11 11 11 11 12	12 12 13 13 13
.81 .82 .83 .84 .85	6457 6607 6761 6918 7079	6471 6622 6776 6934 7096	6486 6637 6792 6950 7112	6501 6653 6808 6966 7129	6516 6668 6823 6982 7145	6531 6683 6839 6998 7161	6546 6699 6855 7015 7178	6561 6715 6871 7031 7194	6577 6730 6887 7047 7211	6592 6745 6902 7063 7228	2 2 2 2 2 2 2	33333	55555	6 6 6 7	8 8 8 8	9 9 10 10	11 11 11 11 12	12 12 13 13 13	14 14 14 15 15
.86 .87 .88 .89 .90	7244 7413 7586 7762 7943	7261 7430 7603 7780 7962	7278 7447 7621 7798 7980	7295 7464 7638 7816 7998	7311 7482 7656 7834 8017	7328 7499 7674 7852 8035	7345 7516 7691 7870 8054	7362 7534 7709 7889 8072	7379 7551 7727 7907 8091	7396 7568 7745 7925 8110	2 2 2 2 2 2 2	3 3 4 4 4	55556	7 7 7 7 7	8 9 9 9	10 10 11 11 11	12 12 12 12 13	13 14 14 14 15	15 16 16 16 17
.91 .92 .93 .94 .95	8128 8318 8511 8710 8913	8147 8337 8531 8730 8933	8166 8356 8551 8750 8954	8185 8375 8570 8770 8974	8204 8395 8590 8790 8995	8222 8414 8610 8810 9016	8241 8433 8630 8831 9036	8260 8453 8650 8851 9057	8279 8472 8670 8872 9078	8299 8492 8690 8892 9099	2 2 2 2 2 2 2	4 4 4 4	6 6 6 6	8 8 8 8	9 10 10 10 10	11 12 12 12 12	13 14 14 14 15	15 15 16 16 17	17 17 18 18 19
.96 .97 .98 .99	9120 9333 9550 9772	9141 9354 9572 9795	9162 9376 9594 9817	9183 9397 9616 9840	9204 9419 9638 9863	9226 9441 9661 9886	9247 9462 9683 9908	9268 9484 9705 9931	9290 9506 9727 9954	9311 9528 9750 9977	2222	4 4 5	6 7 7 7	8 9 9 9	11 11 11 11	13 13 13 14	15 15 16 16	17 17 18 18	19 20 20 20
	0	1	2	3	4	5	6	7	8	9	1	2	3	4	5	6	7	8	9

ANTILOGARITHMS

Note: The logarithmic tables are directly taken from internet and inserted in picture format.

12.13 REFERENCE FOR FURTHER READING

Following books are recommended for further reading:

- Statistics by Murray R, Spiegel, Larry J. Stephens, Mcgraw Hill International Publisher, 4th edition
- Fundamental of Mathematical Statistics by S. C. Gupta and V. K. Kapoor, Sultan Chand and Sons publisher, 11th edition
- Descriptive Statistics by R. J. Shah

13

CORRELATION THEORY

Unit Structure

- 13.0 Objectives
- 13.1 Introduction
- 13.2 Correlation
 - 13.2.1 Scatter Diagram and Linear Correlation
 - 13.2.2 Coefficient of Correlation
 - 13.2.3 Coefficient of Rank Correlation
 - 13.2.4 The Least Square Regression Lines
- 13.3 The Least Square Regression Lines
 - 13.3.1 Regression
 - 13.3.2Least Square Method
 - 13.3.3 Regression Lines and Regression Coefficients
- 13.4 Standard Error of Estimate
- 13.5 Explained and Unexplained Variation
- 13.6 Coefficient of determination
- 13.7 Summary
- 13.8 Exercises
- 13.9 Solution to exercises
- 13.10 Reference for further reading

13.0 OBJECTIVES

This Chapter would make you understand about the following concepts about correlation and regression:

- Scatter Diagram
- Linear Correlation
- Least square regression
- Explained and unexplained variation
- Regression Lines
- Correlation of Time series and Attributes

13.1 INTRODUCTION

By now we know how to find averages and dispersion of a distribution. Involving one variable. These measures give a complete idea about the structure of a distribution.

Sometimes it is necessary to know the relationship between two variables.For instance, a businessman would like to know the effect of production on price. If the two are related, he would like to know the nature of the relationship and to use that knowledge to his benefit. If we get heights and weights of a group of students we can see that in general taller students will be heavier, though there will be some exceptions. As a man's income increases he spends more. That is, there will be some relation between income and expenditure of any person. For such data we would like to find answer for the following questions:

- i) Are the two variables related?
- ii) If they are related, how?
- iii) To what extent they are related?

Correlation helps to find out answers to these questions.

If two variables vary together in the same direction or in opposite directions, they are said to be correlated. That is if as X increases, Y increases consistently, we say that X and Y are positively relatedile the variables are directly related with each other. In this case the values of X and Y for a particular individual have roughly the same relative position among their respective distributions, i.e if X is far above mean of X, then corresponding Y will be above mean of Y. If X is near to mean of X then Y will be near to mean of Y. We know that if the weight of the parents is above average then the son/daughter also will be having more weight.

There are some variables which are negatively correlated where, as X increases, Y decreases and as X decreases, Y increases i.e the variables are inversely proportional to each other, e.g Price increases as the supply decreases, that is, as the commodity becomes scare, the price increases.

If the change in one variable is proportional to the change in the other, the two variables are said to perfectly correlated.

If the two variables are not related to each other, we say the two variables have zero correlation, e.g length of the hair of an individual and the I.Q level of the same individual.

13.2 CORRELATION

The various method of finding whether two given variables are realted or not are:

- Scatter diagram
- Coefficient of Correlation

13.2.1 Scatter Diagram:

Scatter Diagram can be obtained by simply plotting the points on a graph where the two variables, say x and y are taken along x - axis and y - axis respectively.

Eg: A manager of a firm may want to appoint salesman for promoting his sales. When he gets a number of applications, he conducts an aptitude test and selects on the basis of their results. But after employing them,

he wants to know whether there is any relation between the actual sales and the marks in the aptitude test.

The data	for	ceven	calecman	are	26	follows
The uata	101	SUVUI	Salesinan	arc	as	IUIIUWS.

Salesman	1	2	3	4	5	6	7
Aptitude test score	47	49	60	55	59	70	83
Actual sales in '000 in Rs.	70	69	80	75	75	87	90

Solution:

Here the test scores are plotted on the X –axis and actual sales on the Y –axis.



Here we can see that as X increases, Y also increases therefore the two variables are positively related. The businessman knows by this that it is useful to select salesman on the basis of the aptitude test conducted in this case.

Various possible patterns are shown here below



Types of Correlation:

Perfect Positive Correlation: When all plotted values lie exactly on a straight line and the line runs from lower left to upper right corner it is called perfect positive correlation. Here r = 1

Perfect Negative Correlation: When all plotted values lie exactly on a straight line and the line runs from upper left to lower right corner it is called perfect negative correlation. Here r = -1

Positive Correlation: If the value of two variables deviate in the same direction. It is known as Positive correlation or direct correlation.

Negative Correlation: If the value of two variables deviate in the opposite direction. It is also known as the Negative correlation or inverse correlation.

No Correlation: The variables are independent i.e there is no relation between the two variables i.e r = 0.

13.2.2 Coefficient of Correlation:

The things discussed in the previous section gives us the direction of existence of correlation. But we also require to find exact numerical measurement for the degree or extent of correlation. It is useful to have a numerical measure, which is independent of units of the original data, so that the two variables can be compared. For this we calculate the coefficient of correlation. It is denoted by r.

Definition: The coefficient of Correlation denoted by r and name after **Karl Pearson** is defined as

$$r = r_{x,y} = \frac{\sum ((x - \bar{x})(y - \bar{y}))}{N\sigma_x \sigma_y}$$
$$\sigma_x = \sqrt{\left(\frac{\sum (x - \bar{x})^2}{N}\right)}, \sigma_y = \sqrt{\left(\frac{\sum y - \bar{y}}{N}\right)}$$

where there are N pairs.

This is also called **Product Moment Coefficient of Correlation.** Covariance of x and y is defined as $cov(x, y) = \frac{\sum (x - \bar{x})(y - \bar{y})}{N}$ $\therefore r = \frac{cov(x, y)}{N}$

The formula of r can be simplified as
$$r = \frac{\sum xy - N\bar{x}\bar{y}}{\sqrt{\sum x^2 - N\bar{x}^2}\sqrt{\sum y^2 - N\bar{y}^2}}$$

$$=\frac{\sum xy - \frac{\sum x \sum y}{N}}{\sqrt{\sum x^2 - \frac{(\sum x)^2}{N}}\sqrt{\sum y^2 - \frac{(\sum y)^2}{N}}}$$

Properties of Correlation:

- $1) \quad -1 \le r \le 1$
- 2) r = 1, perfect + vecorrelation
- 3) r = -1, perfect vecorrelation
- 4) If 0 < r < 1, the correlation is positive
- 5) If -1 < r < 0, the correlation is negative
- 6) r = 0, no correlation
- r is a pure number and is not affected by change of origin and scale in magnitude.

$$i.eifu = \frac{x-a}{b}$$
, $v = \frac{y-c}{d}thenr_{xy} = \frac{bd}{|b||d|}r_{ux}$

- a) If b and d are of same sign then $r_{xy} = r_{uv}$
- b) If b and d are of opposite signs then $r_{xy} = -r_{uv}$
- 8) If y = ax + b, $r_{xy} = 1$ if a > 0 and $r_{xy} = -1$, if a < 0

Eg 1: $r_{xy} = 0.6$, find r_{uv} if 2u - 3x + 4 = 0, 4v - 16y + 11 = 0

solution: $2u - 3x + 4 = 0 \Rightarrow u = \frac{3x - 4}{2} = \frac{x - \frac{4}{3}}{\frac{2}{3}} \Rightarrow b = \frac{2}{3}$ $4v - 16y + 11 = 0 \Rightarrow v = \frac{16y - 11}{4} = \frac{y - \frac{11}{16}}{\frac{4}{16}} \Rightarrow d = \frac{4}{16}$

b and d are of same signs $\therefore r_{uv} = r_{xy} = 0.6$

Eg 2: $r_{xy} = 0.6$, find r_{uv} if 2u + 3x + 4 = 0, 4v - 16y + 11 = 0

solution: $2u + 3x + 4 = 0 \Rightarrow u = -\frac{x + \frac{4}{3}}{\frac{2}{3}} \Rightarrow b = -\frac{2}{3}$,

$$4v - 16y + 11 = 0 \Rightarrow v = \frac{y - \frac{11}{16}}{\frac{4}{16}} \Rightarrow d = \frac{4}{16}$$

b and d are of opposite signs $\therefore r_{uv} = -r_{xy} = -0.6$

Eg 3: If 2x + y = 3 what is the value of r_{xy} .

solution: $2x + y = 3 \Rightarrow y = -2x + 3 \Rightarrow a = -2 < 0 \Rightarrow r_{xy} = -1$

Eg4: Calculate the karlpearson's coefficient of correlation given $cov(x, y) = -15, \sigma_x = 5, \sigma_y = 4.$

solution:
$$r = \frac{cov(x,y)}{\sigma_x \sigma_y} = -\frac{15}{5 \times 4} = -\frac{15}{20} = -\frac{3}{4} = -0.75$$

Eg 5: Calculate the coefficient of correlation for the following:

Х	-2	-1	0	1	2
У	4	1	0	1	4

solution: $N = 5, \Sigma x = 0, \Sigma y = 10, \Sigma (x - \bar{x})(y - \bar{y}) = 0, \sigma_x = \sqrt{\frac{10}{5}} =$

$$\sqrt{2}, \sigma_y = \sqrt{\frac{14}{5}} = \sqrt{2.8}$$

$$r = \frac{\sum (x - \bar{x})(y - \bar{y})}{N\sigma_x \sigma_y} = \frac{0}{5 \cdot \sqrt{2} \cdot \sqrt{2.8}} = 0$$

Remark: If there is no correlation between the two variables, r = 0 but the converse is not true. In this example the values are related by the equation $y = x^2$ but the observed value of coefficient of correlation is zero.

Eg 6: Calculate the coefficient of correlation for the following:

Х	1	2	3	4	-5	6	7	8	9	10
у	2	4	9	7	10	5	14	16	2	20

solution:N = 10

Х

y

XУ

Total

$$r = \frac{\sum xy - \frac{\sum x \sum y}{n}}{\sqrt{\sum x^2 - \frac{(\sum x)^2}{n}} \sqrt{\sum y^2 - \frac{(\sum y)^2}{n}}} = \frac{\left(589 - \frac{55 \times 89}{10}\right)}{\sqrt{385 - \frac{(55)^2}{10}} \sqrt{1131 - \frac{(89)^2}{10}}}$$

$$(589 - 489.5) \qquad 99.5$$

$$r = \frac{(303^{\circ} - 403.3)}{\sqrt{82.5} \times \sqrt{338.9}} = \frac{333.3}{(9.08)(18.41)} = 0.6$$

Eg 7: Find the coefficient of correlation for the following:

Х	60	50	45	47	53	70	75	57	73	48
у	30	29	29	28	29	35	40	32	35	28

solution: N =	10, '	we take a	c = 60, c =	= 35	(assumed	l mean)	
---------------	-------	-----------	-------------	------	----------	---------	--

solution: N	= 10	, we ta	ke a =	= 60, c =	= 35(as	ssume	d mea	n)			Total
Х	60	50	45	47	53	70	75	57	73	48	
у	30	29	29	28	29	35	40	32	35	28	
u = x - 60	0	-10	-15	-13	-7	10	15	-3	13	-12	-22
v = y - 35	-5	-6	-6	-7	-6	0	5	-3	0	-7	-35
uv	0	60	90	91	42	0	75	9	0	84	451
u^2	0	100	125	169	49	100	225	9	169	144	1190
v^2	25	36	36	49	36	0	25	9	0	49	265

here b and d are same i.e 1:
$$r_{xy} = r_{uv} = \frac{\sum uv - \frac{\sum u \sum v}{N}}{\sqrt{\sum u^2 - \frac{(\sum u)^2}{N}} \sqrt{\sum v^2 - \frac{\sum v^2}{N}}}$$

$$=\frac{451 - \frac{(-22)(-35)}{10}}{\sqrt{1190 - \frac{(-22)^2}{10}}\sqrt{265 - \frac{(-35)^2}{10}}} = \frac{374}{\sqrt{1141.6}\sqrt{142.5}} = 0.927$$

Eg 8:Calculate the coefficient of correlation for the following:

Х	5	10	10	15	15	20	25	30
у	15	17	17	19	21	21	19	17

soluti	on: $N = 8$, we t	ake a =	= 20, c	= 19, b	0 = 5,	d = 2				Total
	Х	5	10	10	15	15	20	25	30	
	у	15	17	17	19	21	21	19	17	
	x - 20	-3	-2	-2	-1	-1	0	1	2	-6
	$u = \frac{1}{5}$									
	y - 19	-2	-1	-1	0	1	1	0	-1	-3
	$v = \frac{1}{2}$									
	uv	6	2	2	0	-1	0	0	-2	7
	u^2	9	4	4	1	1	0	1	4	24
	v^2	4	1	1	0	1	1	0	1	9

Since b and d are of same signs,
$$r_{xy} = r_{uv} = \frac{\sum uv - \frac{\sum u \sum v}{N}}{\sqrt{\sum u^2 - \frac{(\sum u)^2}{N}} \sqrt{\sum v^2 - \frac{\sum v^2}{N}}}$$

$$r_{xy} = r_{uv} = \frac{7 - \frac{(-6)(-3)}{8}}{\sqrt{24 - \frac{(-6)^2}{8}}\sqrt{9 - \frac{(-3)^2}{8}}} = \frac{7 - 2.25}{\sqrt{19.5}\sqrt{7.87}} = 0.38$$

Eg 9: From the data given below find the number of items n, r = $0.5, \sum (x - \bar{x})(y - \bar{y}) = 120, \sigma_y = 8, \sum (x - \bar{x})^2 = 90.$

solution:
$$r = \frac{\sum (x - \bar{x})(y - \bar{y})}{n\sigma_x \sigma_y}$$
, $\sigma_x = \sqrt{\left(\frac{\sum (x - \bar{x})^2}{n}\right)}$, $\sigma_y = \sqrt{\frac{\sum (y - \bar{y})^2}{n}}$

$$0.5 = \frac{120}{n \times \sqrt{\left(\frac{90}{n}\right)} \times 8} \Rightarrow 0.5 = \frac{120}{\sqrt{n^2 \times \frac{90}{n}} \times 8}$$
$$0.5 = \frac{120}{\sqrt{90n \times 8}} \Rightarrow \sqrt{90n} = \frac{120}{0.5 \times 8} = 30$$
$$squaring we get, 90n = 900 \Rightarrow n = 10$$

Eg 10:Calculate the coefficient of correlation between x and y.

	Х	у
no.of observations	15	15
arithmetic mean	25	18
standard deviation	3.01	3.03
sum of squares of the	136	138
deviation from		
arithmetic mean		
$\sum (x - \bar{x})(y - \bar{y}) = 122$		

solution:
$$r = \frac{\sum (x - \bar{x})(y - \bar{y})}{n\sigma_x \sigma_y}$$
, $\sigma_x = \sqrt{\left(\frac{\sum (x - \bar{x})^2}{n}\right)}$, $\sigma_y = \sqrt{\frac{\sum (y - \bar{y})^2}{n}}$

$$r = \frac{122}{15 \times \sqrt{\frac{136}{15}} \times \sqrt{\frac{138}{15}}} = \frac{122}{15 \times 3.01 \times 3.03} = 0.89$$

Eg 11:n = 20, r = 0.3, $\bar{x} = 15$, $\bar{y} = 20$, $\sigma_x = 4$, $\sigma_y = 5$. One pair (27, 30) was wrongly taken as (17, 35). Find corrected value of r.

solution: $\overline{x} = 15, \frac{\Sigma x}{n} = 15 \Rightarrow \Sigma x = 15 \times 20 = 300$

corrected value $\bar{x} = \frac{300 - 17 + 27}{20} = \frac{310}{20} = 15.5$

$$\overline{y} = 20, \frac{\sum y}{n} = 20 \Rightarrow \sum y = 20 \times 20 = 400$$

corrected value $\bar{y} = \frac{400-35+30}{20} = \frac{395}{20} = 19.75$

$$\sigma_x = 4 \Rightarrow (\sigma_x)^2 = 16 \Rightarrow \frac{\sum x^2}{n} - (\bar{x})^2 = 16$$

 $\sum x^2 = (16 + 225) \times 20 = 4820$

corrected value of $\sum x^2 = 4820 + 27 \times 27 - 17 \times 17 = 5260$ corrected value of $\sigma_x = \sqrt{\frac{5260}{20} - (15.5)^2} = \sqrt{22.75} = 4.77$

$$\sigma_{y} = 5 \Rightarrow (\sigma_{y})^{2} = 25 \Rightarrow \frac{\sum y^{2}}{n} - (\bar{y})^{2} = 25$$

$$\sum y^{2} = (25 + 400) \times 20 = 8500$$

corrected value of $\sum y^{2} = 8500 + 30 \times 30 - 35 \times 35 = 8175$
corrected value of $\sigma_{y} = \sqrt{\frac{8500}{20} - (19.75)^{2}} = 5.91$

$$r = \frac{\sum (x - \bar{x})(y - \bar{y})}{n\sigma_{x}\sigma_{y}} \Rightarrow 0.3 = \frac{\sum (x - \bar{x})(y - \bar{y})}{20 \times 4 \times 5}$$

$$\Rightarrow \sum (x - \bar{x})(y - \bar{y}) = 120$$

corrected value of

$$\sum (x - \bar{x})(y - \bar{y}) = 120 + (27 - 17.5)(30 - 19.75) - (17 - 15)(35 - 20)$$

$$= 120 + 9.5 \times 10.25 - (2) \times (15) = 187.375 = 187.38$$

$$187.38$$

corrected value of
$$r = \frac{187.38}{20 \times 4.77 \times 5.91} = 0.33$$

13.2.3 Coefficient of Rank Correlation:

In certain types of characteristics it is not possible to get numerical measurements, but we can rank the individuals in order according to our own judgment, e.g. beauty, smartness. If two persons rank a given group of individuals and we have to find how far the two judges agree with each other, the technique of rank correlation can be used. In some cases though actual measurements are available we may be interested in only the ranks, that is the relative position of an individual in the group. Here also the rank correlation is used.

The formula for Spearman's rank correlation coefficient is

 $R = 1 - \frac{6\sum d^2}{N(N^2 - 1)}$ where d = difference between the ranks of the same

individual, N = number of individuals.

Remark: R follows the same property as r

Eg 1: The ranks according to judges in a beauty contest are

R ₁	1	2	3	4	5	6
R ₂	4	1	2	3	6	5

Find the coefficient of rank correlation.

Total

Solution:N = 6

	$R_1(d_1)$	1	2	3	4	5	6	
	$R_2(d_2)$	4	1	2	3	6	5	
	$\mathbf{d} = \mathbf{d}_1 - \mathbf{d}_2$	-3	1	1	1	-1	1	
	d^2	9	1	1	1	1	1	14
_ 1	$6\sum d^2$	_ 1		14	_ 1	14	21 3	

$$R = 1 - \frac{62a}{N(N^2 - 1)} = 1 - 6 \times \frac{14}{6(36 - 1)} = 1 - \frac{14}{35} = \frac{21}{35} = \frac{5}{5}$$
$$= 0.6.$$

Eg 2: Find spearman's rank correlation coefficient between cost and sales for the following

		0								
cost	39	65	62	90	82	75	25	98	36	78
sales	47	53	58	86	62	68	60	91	51	84

Solution: N = 10

cost (X)	39	65	62	90	82	75	25	98	36	78	
sales (Y)	47	53	58	86	62	68	60	91	51	84	
d ₁	8	6	7	2	3	5	10	1	9	4	
d ₂	10	8	7	2	5	4	6	1	9	3	
$\mathbf{d} = \mathbf{d}_1 - \mathbf{d}_2$	-2	-2	0	0	-2	1	4	0	0	1	
d^2	4	4	0	0	4	1	16	0	0	1	30

$$R = 1 - \frac{6\sum d^2}{N(N^2 - 1)} = 1 - 6 \times \frac{30}{10(100 - 1)} = 1 - \frac{18}{99} = \frac{81}{99} = \frac{9}{11}$$

= 0.8182

Coefficient of Rank correlation when ranks are repeated

In the above example, the ranks were different for all the but in some cases, two or more items may have the same numerical values and ranks should be the same for these values. Suppose we give ranks 1,2,3 and the next two values have to given same rank. In this case next two ranks are 4 and 5. These are to be distributed equally. Therefore both the individuals will get the rank $\frac{4+5}{2}$ and the next one will get the rank as 6.

When there are groups getting the same rank, there is some adjustment in the formula also. If $m_1, m_2, m_3, ...$ denotes the number of times the same rank appear, the coefficient of Rank Correlation will be

$$R = 1 - \frac{6\{\sum d^2 + CF\}}{N(N^2 - 1)}, \text{ where} = \frac{1}{12} [(m_1^3 - m_1) + (m_2^3 - m_2) + (m_3^2 - m_3) + \cdots], \text{ CF} - \text{Correction Factor}$$

Total

Eg 1: Marks of 10 students in two I.Q tests carried out in two successive months. Find the coefficient of correlation.

marks test 1	75	60	60	73	55	57	53	72	65	69
marks test 2	65	65	64	70	58	60	58	68	63	65

Solution: N = 10

Total

marks test 1	75	60	60	73	55	57	53	72	65	69	
marks test 2	65	65	64	70	58	60	58	68	63	65	
d ₁	1	6.5 { 6 }	6.5 { 7 }	2	9	8	10	3	5	4	
d ₂	4 { 3 }	4 { 4 }	6	1	9.5 { 9 }	8	9.5 { 10 }	2	7	4 {5}	
$d = d_1 - d_2$	-3	2.5	0.5	1	- 0.5	0	0.5	1	2	0	
d^2	9	6.2 5	0.2 5	1	0.2 5	0	0.25	1	4	0	22

 $m_1 = 2$, {no. of times marks 60 is repeated in test 1}

 $m_2 = 3$, {no. of times marks 65 is repeated in test 2}

 $m_3 = 2$, {no. of times marks 58 is repeated in test 2}

$$CF = \frac{1}{12} \{ (m_1^3 - m_1) + (m_2^3 - m_2) + (m_3^3 - m_3) \} = \frac{1}{12} \{ 6 + 24 + 6 \}$$
$$= \frac{1}{12} \{ 36 \} = 3$$
$$R = 1 - \frac{6\{ \sum d^2 + CF \}}{n(n^2 - 1)} = 1 - \frac{6\{22 + 3\}}{10(99)} = 1 - 6 \times \frac{25}{990} = 1 - \frac{15}{99}$$
$$= \frac{84}{99} = 0.8485$$

Eg 2: The coefficient of Rank correlation for certain data is found to 0.6. If the sum of the squares of the differences is given to be 66, find the number of observations.

Solution: Given R = 0.6, $\sum d^2 = 66$, To find number of observations i.e to find N

$$R = 1 - \frac{6\sum d^2}{N(N^2 - 1)} \Rightarrow 0.6 = 1 - \frac{6 \times 66}{N(N^2 - 1)}$$
$$\Rightarrow \frac{6 \times 66}{N(N^2 - 1)} = 1 - 0.6$$
$$\Rightarrow \frac{6 \times 66}{N(N^2 - 1)} = 0.4 \Rightarrow N(N^2 - 1) = \frac{6 \times 66}{0.4}$$
$$\Rightarrow (N - 1)N(N + 1) = 990$$
$$\Rightarrow (N - 1)N(N + 1) = 9 \times 10 \times 11 \Rightarrow N = 10$$

13.3 THE LEAST SQUARE REGRESSION LINES

When we know that two given variables are correlated we try to establish some relation between the two so that we can estimate the value of one of the variables given the value of other e.g If we know that there is positive correlation between the heights and weights of a group of individuals, we can find an equation between the height and the weight. We can estimate the weight of an individual belonging to the same population given his height.

Correlation coefficient only determines whether the variables are related and if so, how strong is the relationship. But it is not useful for prediction. The equations used for prediction or estimation are known as **regression equations.**With the help of regression analysis we establish a model which expresses the functional relationship between the two variables. These are also known as estimation equations.

13.3.1 Regression Lines:

We can find two straight lines which will be useful for estimating Y when X is given. It is known as regression of Yon X. Here X is considered as independent variable. For estimating X when Y is given. It is known as regression of X on Y. Here Y is considered as independent variable. We fit a straight line for the set of points given in the bivariate data.

13.3.2 Least Square Method:

We know that a degree one equation represents a straight line. Therefore we take the equation of the regression line of Y on X as Y = a + bX where a and b are constants. The constant 'a' determines the point where the line cuts the Y – axis and constant 'b' determines the slope of the line. The method of *least squares* is used to determine the constants and we get a and b by solving the following two equations simultaneously which are known as *normal equations*.

$$\sum Y = Na + b \sum X$$
, $\sum XY = a \sum X + b \sum X^2$

Similarly, we take the regression of X on Y as X = c + dY where c and d are constants which can be determined by solving the following normal equations.

$$\sum X = Nc + d\sum Y$$
, $\sum XY = c\sum Y + d\sum Y^2$

Eg 1: Find the regression of Y on X and X on Y for the following data. Also estimate X when X = 7 and estimate X when Y = 16

Also esti	male r v	vnen $\Lambda - /$	and estim	ate A whe	n r - 10
Х	1	2	3	4	5
Y	10	12	15	14	15

Solution: N = 5

N = 5						Total
Х	1	2	3	4	5	15
Y	10	12	15	14	15	66
XY	10	24	45	56	75	210
X^2	1	4	9	16	25	55
Y^2	100	144	225	196	225	890

Normal Equation for regression of Y on X are given by

$$\sum y = Na + b \sum X i.e \ 66 = 5a + 15b$$

$$\sum XY = a \sum X + b \sum X^2$$
 i.e 210 = 15a + 55b

Solving the two equations simultaneously we get a = 9.6 and b = 1.2 \therefore The regression line of Y on X is given by Y = 9.6 + 1.2X - (1)For estimating value of Y when X = 7, we put X = 7 in eq (1) \therefore The estimate will be Y = 9.6 + 1.2(7) = 18

Normal equations for regression of X on Y are given by

$\sum X = Nc + d\sum Y$	$i.e \ 15 = 5c + 66d$
$\sum XY = c\sum Y + d\sum Y^2$	i.e 210 = 66c + 890d

Solving the two equations simultaneously we get c = -5.4255, d = 0.6383 \therefore The regression line of X on Y is given by X = -5.4255 + 0.6383Y ---(2)

For estimating value of X when Y = 16, we put Y = 16 in eq (2) \therefore The estimate will be X = -5.4255 + 0.6383(16) = 4.7873

13.3.3 Regression Lines and Regression Coefficients:

The regression of Y on X is given by

$$Y - \overline{Y} = b_{YX}(X - \overline{X}), b_{YX} = r \frac{\sigma_Y}{\sigma_X}$$

The regression of X on Y is given by

$$X - \overline{X} = b_{XY}(Y - \overline{Y}), b_{XY} = r \frac{\sigma_X}{\sigma_Y}$$

where b_{XY} and b_{YX} are known as regression coefficients.

Properties:

1)
$$b_{YX} = r \frac{\sigma_Y}{\sigma_X}$$
, $b_{XY} = r \frac{\sigma_X}{\sigma_Y}$
 $r^2 = b_{YX} \times b_{XY} \Rightarrow r = \pm \sqrt{b_{YX} \times b_{XY}}$
 $r > 0 \text{ if } b_{XY} \& b_{YX} > 0$
 $r < 0 \text{ if } b_{XY} \& b_{YX} < 0$

2) if u = X - a, v = Y - b then $b_{XY} = b_{uv} \& b_{YX} = b_{vu}$

3)
$$b_{YX} = r \frac{\sigma_Y}{\sigma_X} = \frac{cov(X,Y)}{\sigma_X^2} = \frac{\sum XY - \frac{\sum XY}{N}}{\sum X^2 - \frac{(\sum X)^2}{N}}$$

4)
$$b_{xy} = r \frac{\sigma_X}{\sigma_Y} = \frac{cov(X,Y)}{\sigma_Y^2} = \frac{\left(\sum XY - \frac{\sum X\sum Y}{N}\right)}{\sum Y^2 - \frac{\left(\sum YY - \frac{\sum X\sum Y}{N}\right)}{N}}$$

Eg: 1) The following data are given about the expenditure on clothes and expenditure on entertainment. Average expenditure on clothes Rs. 300, average expenditure on entertainment Rs. 100, S.D of expenditure on clothes Rs. 20, S.D of expenditure on entertainment Rs. 15, coefficient of correlation 0.78. Find the two regression equations.

Solution: Let the expenditure on clothes be denoted by x and the expenditure on entertainment be denoted by y.

Given
$$\bar{x} = 300$$
, $\bar{y} = 100$, $\sigma_x = 20$, $\sigma_y = 15$, $r = 0.78$
 $\therefore b_{xy} = \frac{r\sigma_x}{\sigma_y} = \frac{0.78 \times 20}{15} = 0.78 \times \frac{4}{3} = 1.04$
 $\& b_{yx} = \frac{r\sigma_y}{\sigma_x} = \frac{0.78 \times 15}{20} = 0.78 \times \frac{3}{4} = 0.585$
Regression equation of y on x is given by
 $(y - \bar{y}) = b_{yx}(x - \bar{x}) \Rightarrow y - 100 = 0.585(x - 300)$
 $-0.585x + y = 100 - 175.5 \Rightarrow y = 0.585x - 75.5$
Regression equation of x on y is given by
 $(x - \bar{x}) = b_{xy}(y - \bar{y}) \Rightarrow x - 300 = 1.04(y - 100)$
 $x = 1.04y + 300 - 104 \Rightarrow x = 1.04y + 196$

Eg 2: If $\sum x = 37$, $\sum y = 71$, $\sum xy = 563$, $\sum x^2 = 297$, $\sum y^2 = 1079$, n = 5. Find the two regression equations.

Solution:
$$\bar{x} = \frac{37}{5} = 7.4$$
, $\bar{y} = \frac{71}{5} = 14.2$, $b_{yx} = \frac{\sum xy - \frac{\sum xy}{n}}{\sum x^2 - \frac{(\sum x)^2}{n}} = \frac{(563 - \frac{37 \times 71}{5})}{297 - \frac{(37)^2}{5}}$
 $b_{yx} = \frac{563 \times 5 - 37 \times 71}{297 \times 5 - 37 \times 37} = \frac{2815 - 2627}{1485 - 1369} = \frac{188}{116} = 1.62$
 $b_{xy} = \frac{\sum xy - \frac{\sum xy}{n}}{\sum y^2 - \frac{(\sum y)^2}{n}} = \frac{(563 - \frac{37 \times 71}{5})}{1079 - \frac{(71)^2}{5}}$
 $b_{xy} = \frac{563 \times 5 - 37 \times 71}{1079 \times 5 - 71 \times 71} = \frac{188}{354} = 0.53$

regression equation of y on x

$$y - \overline{y} = b_{yx}(x - \overline{x}) \Rightarrow y - 14.2 = 1.62 \times (x - 7.4)$$

$$y = 1.62x + 2.21$$

regression equation of x on y

$$x - \overline{x} = b_{xy}(y - \overline{y}) \Rightarrow x - 7.4 = 0.53 \times (y - 14.2)$$

$$x = 0.53y - 0.126$$

Eg 3: Find the two regression lines for the following data

Price in Rs.	100	120	110	110	160	150
Demand(in units)	40	38	43	45	37	23

Also estimate the demand when price is 130 and the price when demand is 30 units.

Solution: N=6							Total
Price in Rs. (X)	100	120	110	110	160	150	750
Demand(in units) (Y)	40	38	43	45	37	23	226
XY	4000	4560	4730	4950	5920	3450	27610
X^2	10000	14400	12100	12100	25600	22500	96700
Y^2	1600	1444	1849	2025	1369	529	8816

$$\bar{x} = \frac{\sum x}{n} = \frac{750}{6} = 125, \ \bar{y} = \frac{\sum y}{n} = \frac{226}{6} = 37.67$$

$$b_{yx} = \frac{\sum xy - \frac{\sum x \sum y}{n}}{\sum x^2 - \frac{(\sum x)^2}{n}} = \frac{\left(27610 - \frac{750 \times 226}{6}\right)}{96700 - 750 \times \frac{750}{6}} = -\frac{3840}{17700} = -0.22$$

$$b_{xy} = \frac{\left(\sum xy - \frac{\sum x \sum y}{n}\right)}{\sum y^2 - \frac{(\sum y)^2}{n}} = -\frac{3840}{1820} = -2.11$$

regression equation of y on x

 $y - \bar{y} = b_{yx}(x - \bar{x}) \Rightarrow y - 37.67 = -0.22(x - 125)$ y = -0.22x + 65.17 - - - -(1)regression equation of x on y $x - \bar{x} = b_{xy}(y - \bar{y}) \Rightarrow x - 125 = -2.11(y - 37.67)$ x = -2.11y + 204.48 - - - - (2)

demand =? when price = 130, we put
$$x = 130$$
 in (1)
 $y = -0.22 \times 130 + 65.17 = 36.57$
price = ? when demand = 30, we put $y = 30$ in (2)
 $x = -2.11 \times 30 + 204.48 = 141.18$

Eg 4: The two regression lines are given by x + 2y = 5, 2x + 3y - 8 = 0, $(\sigma_x)^2 = 12$. Find the values of \overline{x} , \overline{y} , $(\sigma_y)^2$.

Solution: We solve the two regression equations simultaneously to find $\bar{x}\&\bar{v}$ x + 2y = 5 - -(1)2x + 3y - 8 = 0 - -(2)*Mul eq* (1) *by* 2 *and subtract from eq* (2) *gives* $\bar{y} = 2$ substitute y = 2 in eq (1) gives $\bar{x} = 1$ $x + 2y = 5 \Rightarrow x = -2y + 5, b_{xy} = -2$ $2x + 3y - 8 = 0 \Rightarrow y = \left(-\frac{2}{3}\right)x + 4, \qquad b_{yx} = -\frac{2}{3}$ $b_{xy} \times b_{yx} = \frac{4}{3} > 1$ not possible regression of y on x is given by x + 2y = 5 and x on y is *given by* 2x + 3y - 8 = 0 $x + 2y = 5 \Rightarrow y = -\frac{x}{2} + \frac{5}{2}, b_{yx} = -\frac{1}{2}$ $2x + 3y = 8 \Rightarrow x = \left(-\frac{3}{2}\right)y + 4$, $b_{xy} = -\frac{3}{2}$ $r^{2} = b_{yx} \times b_{xy} \Rightarrow r^{2} = \frac{3}{4} = 0.75 \Rightarrow r = -\sqrt{\frac{3}{4}} = -0.87$ $r^{2} = \left(b_{yx}\right)^{2} \times \frac{\left(\sigma_{y}\right)^{2}}{\left(\sigma_{y}\right)^{2}} \Rightarrow r^{2} = \frac{1}{4} \times \frac{\left(\sigma_{y}\right)^{2}}{12} \Rightarrow \frac{3}{4} = \frac{1}{4} \times \frac{\left(\sigma_{y}\right)^{2}}{12}$ $\Rightarrow (\sigma_v)^2 = 0.75 \times 48 = 36$

13.4 STANDARD ERROR OF ESTIMATE

Ley y_1 be the value of y for given value of x, a measure of the scatter about the regression line y on x is given by

$$S_{y,x} = \sqrt{\frac{\sum(y - y_1)^2}{n}}$$

which is called the standard error of estimate of y on x. Standard error of estimate of x on y

$$S_{x,y} = \sqrt{\frac{\sum (x - x_1)^2}{n}}$$

In general, $S_{y,x} \neq S_{x,y}$

Eg: 1) If the regression line of y on x is $y = a_0 + a_1 x$. Show that $S_{y,x}^2 = \frac{\sum y^2 - a_0 \sum y - a_1 \sum xy}{n}.$

solution: $y_1 = a_0 + a_1 x$

$$\begin{split} S_{y,x}^{2} &= \frac{\sum(y-y_{1})^{2}}{n} = \frac{\sum(y-y_{1})^{2}}{n} = \frac{\sum(y^{2}-2yy_{1}+y_{1}^{2})}{n} \\ &= \frac{\sum(y^{2}-2y(a_{0}+a_{1}x)+(a_{0}+a_{1}x)^{2})}{n} \\ &= \frac{\sum(y^{2}-2a_{0}y-2a_{1}xy+a_{0}^{2}+2a_{0}a_{1}x+a_{1}^{2}x^{2})}{n} \\ &= \frac{\sum(y^{2}-a_{0}y-a_{1}xy-a_{0}y-a_{1}xy+a_{0}^{2}+a_{0}a_{1}x+a_{1}^{2}x^{2}+a_{0}a_{1}x)}{n} \\ &= \frac{\left\{\sum y(y-a_{0}-a_{1}x)-a_{0}\sum(y-a_{1}x-a_{0})-a_{1}\sum x(y-a_{1}x-a_{0})\right\}}{n} \\ &= \frac{\left\{\sum y(y-a_{0}-a_{1}x)-a_{0}\sum(y-a_{1}x-a_{0})-a_{1}\sum x(y-a_{1}x-a_{0})\right\}}{n} \\ &= a_{0}+a_{1}x \ by \ least \ square \ method \ to \ find \ a_{0} \ and \ a_{1}, \\ &we \ solve \ \sum y = a_{0}n+a_{1}\sum x \ and \ \sum xy \ = a_{0}\sum x+a_{1}\sum x^{2} \end{split}$$

$$\Rightarrow \sum (y - a_0 - a_1 x) = 0$$
, $\sum (xy - a_0 x - a_1 x^2) = 0$

$$S_{y,x}^{2} = \frac{\sum y(y - a_{0} - a_{1}x)}{n} = \frac{\sum y^{2} - a_{0}\sum y - a_{1}\sum xy}{n}$$

Eg: 2) If the regression line of y on x is y = 35.82 + 0.476x and values of x and y are

Х	65	63	67	64	68	62	70	
у	68	66	68	65	69	66	68	

Find	$S_{y,x}^2$

solution:	y = 35.82 +	0.476x
-----------	-------------	--------

_

5				
Х	у	$y_1 = 35.82 + 0.476x$	$(y - y_1)$	$(y - y_1)^2$
65	68	66.76	1.24	1.538
63	66	65.808	0.192	0.037
67	68	67.712	0.288	0.083
64	65	66.284	-1.284	1.649
68	69	68.188	0.812	0.659
62	66	65.332	0.668	0.446

70	68	69.14	-1.14	1.30
total				5.712

$$S_{y,x}^2 = \frac{\sum (y - y_1)^2}{n} = \frac{5.712}{7} = 0.816$$

Eg: 3) Show that $\sum (y - \bar{y})^2 = \sum (y - y_1)^2 + \sum (y_1 - \bar{y})^2$ solution: $lhs = \sum (y - \bar{y})^2 = \sum (y - y_1 + y_1 - \bar{y})^2$ $= \sum \{(y - y_1)^2 + 2(y - y_1)(y_1 - \bar{y}) + (y_1 - \bar{y})^2\}$ $= \sum (y - y_1)^2 + \sum (y_1 - \bar{y})^2 + 2\sum (y - y_1)(y_1 - \bar{y})$ nowtoprove, $\sum (y - y_1)(y_1 - \bar{y}) = 0$ i. $e \sum (y - y_1)(y_1 - \bar{y}) = \sum (y - a_0 - a_1x)(a_0 + a_1x - \bar{y})$ $= a_0\sum (y - a_0 - a_1x) + a_1\sum x(y - a_0 - a_1x) - \bar{y}\sum (y - a_0 - a_1x) = 0$ $y = a_0 + a_1x$, using leasts quaremethod we find a_0 and a_1 , we solve $\sum y = a_0n + a_1\sum x$ and $\sum xy = a_0\sum x + a_1\sum x^2$ $\Rightarrow \sum (y - a_0 - a_1x) = 0$, $\sum (xy - a_0x - a_1x^2) = 0$ $\Rightarrow \sum (y - \bar{y})^2 = \sum (y - y_1)^2 + \sum (y_1 - \bar{y})^2$

13.5 EXPLAINED AND UNEXPLAINED VARIATION

In the previous section we had proved,

$$\sum (y - \bar{y})^2 = \sum (y - y_1)^2 + \sum (y_1 - \bar{y})^2$$

 $\sum (y - y_1)^2$ is called unexplained variation as the deviation $(y - y_1)$ behave randomly and have an unpredictable manner.

 $\sum (y_1 - \overline{y})^2$ is called explained variation as the deviation $(y_1 - \overline{y})$ have a definite pattern.

 $\sum (y - \bar{y})^2$ is called as total variation.

Eg: 1) For the following calculate total variation, explained variation and unexplained variation, given the regression of on x as y = 35.82 + 0.476x

	х	65	63	67	64	68	62	70	
	у	68	66	68	65	69	66	68	
in	$\overline{y} = 35.82 \pm 0.476$ $\overline{y} = 57.143$								

Solution: y = 35.82 + 0.476x, $y = \sum_{n=0}^{\infty} \frac{1}{n} = 07.145$

Х	у	$y_1 = 35.82 + 0.476x$	$(y - y_1)$	$(y - y_1)^2$	$(y_1 - \bar{y})^2$
65	68	66.76	1.24	1.538	0.147
63	66	65.808	0.192	0.037	1.306
67	68	67.712	0.288	0.083	0.734
64	65	66.284	-1.284	1.649	4.592
68	69	68.188	0.812	0.659	3.448

62	66	65.332	0.668	0.446	1.306
70	68	69.14	-1.14	1.30	0.734
total				5.712	12.267

Unexplained variation= $\sum (y - y_1)^2 = 5.712$

Explained variation = $\sum (y_1 - \overline{y})^2 = 12.267$

Total variation = $\sum (y - \bar{y})^2 = \sum (y - y_1)^2 + \sum (y_1 - \bar{y})^2 = 17.979$

13.6 COEFFICIENT OF DETERMINATION

The ratio of the explained variation to the total variation is called the coefficient of determination.

It is given by $\frac{\sum(y_1 - \bar{y})^2}{\sum(y - \bar{y})^2}$.

If there is zero explained variation (i.e the total variation is same as unexplained variation) then coefficient of determination will be 0. If there is zero unexplained variation (i.e the total variation is same as explained variation) then coefficient of determination will be 1. Since the ratio is always non negative, we denote it by r^2 . realled as coefficient of correlation is given by

$$r = \pm \sqrt{\frac{explained \ variation}{total \ variation}} = \pm \sqrt{\frac{\sum(y_1 - \bar{y})^2}{\sum(y - \bar{y})^2}}$$

varies from between -1 and +1. The + and - signs indicates positive and negative linear correlation respectively. Note *r* is a dimensionless quantity, i.e it does not depend on the units.

For the case of linear correlation, the quantity r is the same regardless of whether X or Y is considered the independent variables. Thus r is a good measure of the linear correlation between two variables.

Eg : 1)Show that $s_{y,x}^2 = s_y^2(1-r^2)$.

$$\begin{aligned} \text{Solution:} r &= \pm \sqrt{\frac{explained \ variation}{total \ variation}} = \pm \sqrt{\frac{\Sigma(y_1 - \bar{y})^2}{\Sigma(y - \bar{y})^2}} \\ r^2 &= \left(\sqrt{\frac{\Sigma(y_1 - \bar{y})^2}{\Sigma(y - \bar{y})^2}} \right)^2 = \left(\sqrt{\frac{\Sigma(y - \bar{y})^2 - \Sigma(y - y_1)^2}{\Sigma(y - \bar{y})^2}} \right)^2 \\ &\Rightarrow r^2 = \left(\sqrt{1 - \left(\frac{\Sigma(y - y_1)^2}{\Sigma(y - \bar{y})^2}\right)} \right)^2 = \left(\sqrt{1 - \frac{s_{y,x}^2}{s_y^2}} \right)^2 \\ &\Rightarrow s_{y,x}^2 = s_y^2 (1 - r^2) \end{aligned}$$

Eg: 2) Given the regression of y on x as y = 35.82 + 0.476x. Find the coefficient of determination and coefficient of correlation for the following data

	U							_
Х	65	63	67	64	68	62	70	
у	68	66	68	65	69	66	68	

	1			~	-
Х	У	$y_1 = 35.82 + 0.476x$	$(y - y_1)$	$(y - y_1)^2$	$(y_1 - \bar{y})^2$
65	68	66.76	1.24	1.538	0.147
63	66	65.808	0.192	0.037	1.306
67	68	67.712	0.288	0.083	0.734
64	65	66.284	-1.284	1.649	4.592
68	69	68.188	0.812	0.659	3.448
62	66	65.332	0.668	0.446	1.306
70	68	69.14	-1.14	1.30	0.734
total				5.712	12.267

Unexplained variation $=\sum (y - y_1)^2 = 5.712$ Explained variation $=\sum (y_1 - \overline{y})^2 = 12.267$

Total variation = $\sum (y - \bar{y})^2 = \sum (y - y_1)^2 + \sum (y_1 - \bar{y})^2 = 17.979$

Coefficient of determination $= \frac{\sum (y_1 - \bar{y})^2}{\sum (y - \bar{y})^2} = \frac{12.267}{17.979} = 0.682$

Coefficient of correlation $r = \pm \sqrt{\frac{explained variation}{total variation}} = \pm \sqrt{0.682}$

13.7 SUMMARY

In this chapter we have learnt about the scatter diagram which helps us to find the nature and extent of relationship between the variables, Coefficient of correlation which is a numerical measure of nature and extent of relationship between two given variables whose values lies between +1 and -1. We had also learnt about the coefficient of Rank correlation which is used in cases where it is not possible to get numerical measurements, but we can rank the individuals in order according to our judgment. This chapter also deals with the least square regression lines of y on x as well as x on y. This chapter also explains about the standard error of estimate, Explained and unexplained variation and correlation of time series.

13.8 EXERCISES

1) Following table gives the us the marks obtained by 6 students in mid term exam and final semester exam. Find the coefficient of correlation.

Mid term Exam	12	14	23	18	10	19
Final semester Exam	68	78	85	75	70	74

- 2) Given covariance = 27, r = 0.6, variance of y = 25. Find variance of x.
- 3) Find the coefficient of correlation between the heights of male students and female students

height of male students	65	66	67	68	69	70	71
height of female students	67	68	66	69	72	72	69

- 4) If r = 0.38, cov(x, y) = 10.2, $\sigma_x = 16$. Find σ_y
- 5) Find r, if cov(x, y) = 6, $\sigma_x = 2.45$, $\sigma_y = 3.41$.
- 6) Find r, given $\sum (x \bar{x})(y \bar{y}) = 29, \sum (x \bar{x})^2 = 28, \sum (y \bar{y})^2 = 42.$
- 7) Ten competitors in Miss Universe are ranked by three judges in the following order:

J1	1	6	5	10	3	2	4	9	7	8
J2	3	5	8	4	7	10	2	1	6	9
J3	6	4	9	8	1	2	3	10	5	7

Using rank correlation coefficient determine which pair of judges has the nearest approach to common tastes in beauty.

- 8) The coefficient of rank correlation for certain data is found to be 0.6. If the sum of the squares of the differences is given to be 66. Find the number of items in the group.
- 9) The ranks of 16 students in the subject of DBMS and CN are given as follows. Calculate the rank coefficient of correlation.

Rank in	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
DBMS																
Rank in	1	10	3	4	5	7	2	6	8	11	15	9	14	12	16	13

- 10) The coefficient of rank correlation of marks obtained by 9 students was calculated to be 0.4. It was later discovered that the value of the difference between ranks for one student was written wrongly as 6 instead of 8. Find the correct value of coefficient of rank correlation.
- 11) Find the two regression equations given $\bar{x} = 8, \bar{y} = 2000, \sigma_x = 2, \sigma_y = 80, r = 0.7$. Also estimate y when x = 10 and estimate x when y = 2500.

10

- 12) Find the two regression equations for the following data: $\sum (x - \bar{x})(y - \bar{y}) = 135, \sum (x - \bar{x})^2 = 96, \sum (y - \bar{y})^2 = 206, \sum x$ = 120, $\sum y = 180, n = 5$
- 13) n = 50, regression equation of marks in mathematics (Y) on the marks in English(X) was 4Y- 5X = 8.Mean marks in English are 40. The ratio of the two standard deviation $\sigma_y: \sigma_x = 5:2$. Find the average marks in mathematics and coefficient of correlation between the marks in the two subjects.
- 14) The two regression lines between x and y are given below 2x + 3y = 61, x + y = 25. Find \bar{x} , \bar{y} and r.
- 15) Find the regression line of profits on output from the following data using least square method.

Ouput(100 tons)	5	7	9	11	13	15
Profit per unit(Rs.)	1.7	2.4	2.8	3.4	3.7	4.4

16) For the following find total variation given the regression of y on x as y = 35.82 + 0.476x

Х	65	63	67	64	68	62	70	66	68	67	69	71
у	68	66	68	65	69	66	68	65	71	67	68	70

13.9 SOLUTION TO EXERCISE

Q.	Solution	Q.	Solution
No.		No.	
1	r = 0.81	2	variance of $x = 81$
3	r = 0.67	4	1.68
5	r = 0.72	6	r = 0.85
7	$R_{12} = -0.212, R_{13} = 0.636, R_{23}$	8	n= 10
	= -0.297		
9	R = 0.8	10	R = 0.17
11	y = 28x + 1776, x = 0.02y - 32,	12	y = 1.41x + 2.16,
	y = 2056, x = 18		x = 0.66y + 0.24
13	52, r = 0.5	14	14, 11, -0.82
15	y = 0.26x + 0.5	16	38.917

13.10 REFERENCES

Following books are recommended for further reading:

- Statistics by Murray R, Spiegel, Larry J. Stephens, Mcgraw Hill International Publisher, 4th edition
- Fundamental of Mathematical Statistics by S. C. Gupta and V. K. Kapoor, Sultan Chand and Sons publisher, 11th edition
- Mathematical Statistics by J. N. Kapur and H. C. Saxena, S. Chand publisher, 12th edition
