

16/12/19 M.

[This question paper contains 7 printed pages]

**Your Roll No.** : .....

**Sl. No. of Q. Paper** : 7465 **J**

**Unique Paper Code** : 32351303

**Name of the Course** : **B.Sc.(Hons.)**  
**Mathematics**

**Name of the Paper** : Multivariate Calculus

**Semester** : III

**Time : 3 Hours** **Maximum Marks : 75**

**Instructions for Candidates :**

- (i) Write your Roll No. on the top immediately on receipt of this question paper.
- (ii) **All** Sections are compulsory.
- (iii) Attempt any **five** questions from each **Section**.
- (iv) All questions carry equal marks.

P.T.O.

7465

**Section- I**

1. Given that the function

$$f(x,y) = \begin{cases} \frac{3x^3 - 3y^3}{x^2 - y^2} & \text{for } x^2 \neq y^2 \\ B & \text{otherwise} \end{cases}$$

is continuous at the origin, what is B ?

2. In physics, the *wave equation* is :

$$\frac{\partial^2 z}{\partial t^2} = c^2 \frac{\partial^2 z}{\partial x^2}$$

and the *heat equation* is :

$$\frac{\partial z}{\partial t} = c^2 \frac{\partial^2 z}{\partial x^2}$$

Determine whether  $z = \sin 5ct \cos 5x$  satisfies the wave equation, the heat equation, or neither.

3. The radius and height of a right circular cone are measured with errors of at most 3% and 2%, respectively. Use increments to approximate the maximum possible percentage error in computing the volume of the cone using these measurements and the formula  $V = \frac{1}{3}\pi R^2 H$ .
4. If  $f(x, y, z) = xy^2e^{xz}$  and  $x = 2 + 3t$ ,  $y = 6 - 4t$ ,  $z = t^2$ . Compute  $\frac{df}{dt}(1)$ .
5. Sketch the level curve corresponding to  $C = 1$  for the function  $f(x, y) = \frac{x^2}{a^2} - \frac{y^2}{b^2}$  and find a unit normal vector at the point  $P_0(2\sqrt{3})$ .
6. Find the point on the plane  $2x + y - z = 5$  that is closest to the origin.



7465

**Section - II**

7. Find the volume of the solid bounded above by the plane  $z = y$  and below in the  $xy$ -plane by the part of the disk  $x^2 + y^2 \leq 1$  in the first quadrant.
8. Sketch the region of integration and then compute the integral  $\int_0^1 \int_x^{2x} e^{y-x} dy dx$  in 2 ways :
- (a) with the given order of integration
  - (b) with the order of integration reversed
9. Evaluate  $\int_0^2 \int_0^{\sqrt{2x-x^2}} y\sqrt{x^2+y^2} dy dx$  by converting to polar coordinates.
10. Find the volume of the tetrahedron bounded by the plane  $2x + y + 3z = 6$  and the coordinate planes  $x = 0$ ,  $y = 0$  and  $z = 0$ .

11. Compute  $\iiint_D \frac{dx dy dz}{\sqrt{x^2 + y^2 + z^2}}$  where D is the solid sphere  $x^2 + y^2 + z^2 \leq 3$ .

12. Use the change of variables to compute

$\iint_D \frac{(x-y)^4}{(x+y)^4} dy dx$ , where D is the triangular region bounded by the line  $x + y = 1$  and the coordinate axes.

### Section - III

13. Find the work done by the force field

$$\vec{F} = \frac{x}{\sqrt{x^2 + y^2}} \vec{i} - \frac{y}{\sqrt{x^2 + y^2}} \vec{j} \text{ when an object moves}$$

from  $(a, 0)$  to  $(0, a)$  on the path  $x^2 + y^2 = a^2$ .

14. Verify that the following line integral is independent of the path  $\oint (3x^2 + 2x + y^2) dx + (2xy + y^3) dy$  where C is any path from  $(0, 0)$  to  $(0, 1)$ .

7465

15. Use Green's theorem to evaluate

$$\oint_C (x \sin x dx - \exp(y^2) dy) \text{ where } C \text{ is the closed}$$

curve joining the points  $(1, -1)$ ,  $(2, 5)$  and  $(-1, -1)$  in counterclockwise direction.

16. State Stoke's theorem and use it to evaluate

$$\iint_S \text{curl} \vec{F} \cdot d\vec{S} \text{ where } \vec{F} = xz\vec{i} + yz\vec{j} + xy\vec{k} \text{ and } S \text{ is the}$$

part of the sphere  $x^2 + y^2 + z^2 = 4$  that lies inside the cylinder  $x^2 + y^2 = 1$  and above the  $xy$ -plane.

17. Use the divergence theorem to evaluate the

$$\text{surface integral } \iint_S \vec{F} \cdot \vec{N} dS, \text{ where } \vec{F} = (x^2 + y^2 - z^2)\vec{i} +$$

$$yx^2\vec{j} + 3z\vec{k}; S \text{ is the surface comprised of the five}$$

faces of the unit cube  $0 \leq x \leq 1, 0 \leq y \leq 1, 0 \leq z \leq 1$ ,

missing  $z = 0$ .



7465

18. Evaluate  $\iint_S 2x \, dS$  where  $S$  is the portion of the plane  $x + y + z = 1$  with  $x \geq 0, y \geq 0, z \geq 0$ .