

(M) Lib. 10-12-19

This question paper contains 4+1 printed pages]

Roll No.

--	--	--	--	--	--	--	--	--	--

S. No. of Question Paper : 8081

Unique Paper Code : 32357501

J

Name of the Paper : Numerical Methods

Name of the Course : B.Sc. (H) Mathematics : DSE-2

Semester : V

Duration : 3 Hours

Maximum Marks : 75

(Write your Roll No. on the top immediately on receipt of this question paper.)

All the six questions are compulsory.

Attempt any two parts from each question.

Marks are indicated against each question.

Use of Non-Programmable Scientific Calculator is allowed.

1. (a) A real root of the equation  $x^3 - 5x + 1 = 0$  lies in  $]0, 1[$ . Perform three iterations of Regula Falsi Method to obtain the root.

6

P.T.O.



- (b) Perform three iteration of Bisection Method to obtain root of the equation  $\cos(x) - xe^x$  in  $]0, 1[$ . 6
- (c) Discuss the order of convergence of the Secant method and give the geometrical interpretation of the method. 6
2. (a) Verify  $x = \sqrt{a}$  is a fixed point of the function  $h(x) = \frac{1}{2}\left(x + \frac{a}{x^2}\right)$ . Determine order of convergence of sequence  $p_n = h(p_{n-1})$  towards  $x = \sqrt{a}$ . 6.5
- (b) Use Secant method to find root of  $3x + \sin(x) - e^x = 0$  in  $]0, 1[$ . Perform three iterations. 6.5
- (c) Prove that Newton's Method is of order two using  $x^3 + 2x^2 - 3x - 1 = 0$  and initial approximation  $x_0 = 2$ . 6.5
3. (a) Define a lower and an upper triangular matrix. Solve the system of equations :

$$-3x_1 + 2x_2 - x_3 = -12$$

$$6x_1 + 8x_2 + x_3 = 1$$

$$4x_1 + 2x_2 + 7x_3 = 1$$

by obtaining an LU decomposition of the coefficient matrix

A of the above system.

6

- (b) For Jacobi method, calculate  $T_{jac}$ ,  $C_{jac}$  and spectral radius of the following matrix :

$$\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$$

6

- (c) Set up the Gauss-Seidel iteration scheme to solve the system of equations :

$$4x_1 + 2x_2 - x_3 = 1$$

$$2x_1 + 4x_2 + x_3 = -1$$

$$-x_1 + x_2 + 4x_3 = 1$$

Take the initial approximation as  $X^{(0)} = (0, 0, 0)$  and do three iterations. 6

4. (a) Construct the Lagrange form of the interpolating polynomial from the following data :

$x$	1	2	3
$f(x) = \ln x$	$\ln 1$	$\ln 2$	$\ln 3$

6.5

P.T.O.



(b) Prove that for  $n + 1$  distinct nodal points  $x_0, x_1, x_2, \dots, x_n$ , there exists a unique interpolating polynomial of at most degree  $n$ . 6.5

(c) Find the maximum value of the step size  $h$  that can be used in the tabulation of  $f(x) = e^x$  on the interval  $[0, 1]$  so that the error in the linear interpolation of  $f(x)$  is less than  $5 \times 10^{-4}$ . 6.5

5. (a) Define the backward difference operator  $\nabla$  and the Newton divided difference. Prove that :

$$f[x_0, x_1, \dots, x_n] = \frac{\nabla^n f_n}{n! h^n} \text{ where } h = x_{i+1} - x_i. \quad 6$$

(b) Construct the divided difference table for the following data set and then write out the Newton form of the interpolating polynomial :

$x$	-7	-5	-4	-1
$y$	10	5	2	10

Find the approximation of  $y$  for  $x = -3$ .

6

(c) Use the formula

$$f'(x_0) \approx \frac{f(x_0 + h) - f(x_0)}{h}$$

to approximate the derivative of  $f(x) = 1 + x + x^3$  at  $x_0 = 1$  taking  $h = 1, 0.1, 0.001$ . What is the order of approximation? 6

6. (a) Approximate the value of  $\int_0^1 e^{-x} dx$  using the Trapezoidal rule and verify that the theoretical error bound holds for the same. 6.5

(b) State Simpson's 1/3rd rule for the evaluation of  $\int_a^b f(x) dx$

and prove that it has degree of precision 3. 6.5

(c) Use Euler's method to approximate the solution of the initial value problem.

$$x' = (1 + x^2)/t, \quad x(1) = 0, \quad 1 \leq t \leq 4 \text{ taking 5 steps. } 6.5$$