

[This question paper contains 28 printed pages]

29 DEC 2022

Your Roll No.....

C

Sr. No. of Question Paper : 4515

Unique Paper Code : 12271303

Name of the Paper : Statistical Methods for Economics

Name of the Course : Core CBCS

Semester/Mode : CBCS Semester – III

Duration : 3 Hours

Maximum Marks : 75



### Instructions for Candidates

1. Write your Roll No. on the top immediately on receipt of this question paper.
2. All questions within each section are to be answered in a contiguous manner on the answer sheet. Start each question on a new page, and all subparts of a question should follow one after the other.
3. The use of a simple non-programmable calculator is allowed.
4. Statistical tables are attached for your reference.
5. In all calculations, figures should be rounded to two decimal places.
6. Answers may be written either in English or Hindi; but the same medium should be used throughout the paper.

### छात्रों के लिए निर्देश

1. इस प्रश्न-पत्र के मिलते ही ऊपर दिए गए निर्धारित स्थान पर अपना अनुक्रमांक लिखिए।
2. प्रत्येक खंड के सभी प्रश्नों के उत्तर पत्रक पर सन्निहित तरीके से दीजिये। प्रत्येक प्रश्न को एक नए पृष्ठ पर प्रारंभ कीजिये, और एक प्रश्न के सभी उपभागों को एक के बाद एक अनुसरण कीजिये।
3. एक साधारण गैर-प्रोग्रामेबल कैलकुलेटर के उपयोग की अनुमति है।
4. सास्विकीय टेबल आपके संदर्भ के लिए संलग्न हैं।
5. सभी गणनाओं में, आंकड़ों को दो दशमलव स्थानों पर गोल किया जाना चाहिए।
6. इस प्रश्न-पत्र का उत्तर अंग्रेजी या हिंदी किसी एक भाषा में दीजिए, लेकिन सभी उत्तरों का माध्यम एक ही होना चाहिए।

## SECTION I

*Do both the questions.*

दोनों प्रश्न कीजिये।

1. The following measurements were recorded for the drying time, in hours, of a certain brand of wall paint.

3.4, 2.5, 4.8, 2.9, 3.6, 2.8, 3.3, 5.6, 3.7, 2.8, 4.4, 4.0, 5.2, 3.0, 4.8

Assume the above data relates to the population of the drying time,

- (i) Calculate the population mean. Is it a random variable? Why/Why not?
  - (ii) A sample of three drying times is drawn from the above population. How many such samples can be drawn if it is a random sample?
  - (iii) Suppose two such samples are (3.4, 4.4, 5.2) and (3.6, 3.6, 2.8). What are the respective sample means? Is the sample mean a random variable and why?
- (5)

दीवार के पेंट के एक निश्चित ब्रांड में, सुखाने के समय के लिए निम्नलिखित माप, घंटों में, दर्ज किए गए थे।

3.4, 2.5, 4.8, 2.9, 3.6, 2.8, 3.3, 5.6, 3.7, 2.8, 4.4, 4.0, 5.2, 3.0, 4.8

मान लीजिये कि उपरोक्त डेटा सूखने के समय की आबादी से संबंधित है।

- (i) जनसंख्या औसत की गणना कीजिये। क्या यह एक यादृच्छिक चर है? क्यों/क्यों नहीं?
- (ii) उपरोक्त जनसंख्या में से तीन सुखाने का समय का एक नमूना तैयार किया गया है। यदि यह एक यादृच्छिक नमूना है तो ऐसे कितने नमूने खींचे जा सकते हैं?
- (iii) मान लीजिए कि दो ऐसे नमूने हैं (3.4, 4.4, 5.2) और (3.6, 3.6, 2.8)। संबंधित नमूना औसत क्या हैं? क्या नमूना का औसत एक यादृच्छिक चर है और क्यों?

2. A worker has asked her supervisor for a letter of recommendation for a new job. She estimates that there is an 80 percent chance that she will get the job if she receives a strong recommendation, a 40 percent chance if she receives a moderately

good recommendation, and a 10 percent chance if she receives a weak recommendation. She further estimates that the probabilities that the recommendation will be strong, moderate, and weak are 0.7, 0.2, and 0.1, respectively.

- (i) How certain is she that she will receive the new job offer?
- (ii) Given that she does receive the offer, how likely is it she received a strong recommendation?
- (iii) Given that she does not receive the job offer, how likely is it that she received a weak recommendation? (5)

एक कार्यकर्ता ने अपने पर्यवेक्षक से नई नौकरी के लिए सिफारिश पत्र मांगा है। वह अनुमान लगाती है कि अगर उसे एक मजबूत सिफारिश मिलती है तो उसे नौकरी मिलने की 80 प्रतिशत संभावना है, अगर उसे मामूली अच्छी सिफारिश मिलती है तो 40 प्रतिशत मौका और कमज़ोर सिफारिश मिलने पर 10 प्रतिशत मौका मिलता है। वह आगे अनुमान लगाती है कि सिफारिश के मजबूत, मध्यम और कमज़ोर होने की संभावनाएँ क्रमशः 0.7, 0.2 और 0.1 हैं।

- (i) वह कितनी निश्चित है कि उसे नई नौकरी का प्रस्ताव प्राप्त होगा?
- (ii) यह देखते हुए कि उसे प्रस्ताव प्राप्त होता है, उसे एक मजबूत सिफारिश प्राप्त होने की कितनी संभावना है?
- (iii) यह देखते हुए कि उसे नौकरी का प्रस्ताव नहीं मिला है, इस बात की कितनी संभावना है कि उसे कमज़ोर सिफारिश मिली?

## SECTION II

**Q3 is compulsory.**

*Attempt any two questions from Q4, Q5 and Q6.*

**Q3 अनिवार्य है। Q4, Q5 और Q6 में से किन्हीं दो प्रश्नों का उत्तर दीजिये।**

3. Let  $Y$  be the number of parcels mailed by a randomly selected customer through 'ABC Courier Service'. Suppose the probability mass function (pmf) of  $Y$  is as below

Y	1	2	3	4
P(y)	0.3	0.4	0.1	0.2

Consider a random sample of 2 customers. Derive the sampling distribution of the range of Y. (5)

मान लीजिए Y 'एबीसी कूरियर सेवा' के माध्यम से यादृच्छिक रूप से चुने गए ग्राहक द्वारा मेल किए गए पार्सल की संख्या है। मान लीजिए कि Y का प्रायिकता द्रव्यमान फलन (pmf) इस प्रकार है :

Y	1	2	3	4
P(y)	0.3	0.4	0.1	0.2

2 ग्राहकों के यादृच्छिक नमूने पर विचार करें। Y के परास का न्यादर्शन वितरण व्युत्पन्न कीजिए।

4. (a) A telephone exchange desk employs 5 operators who receive calls independently of one another, each according to a Poisson process at a rate of 4 calls per minute.
- (i) What is the probability that during a given 1-minute interval, the first operator receives no calls?
  - (ii) What is the probability that during a given 45-second interval, the second operator receives no calls?
  - (iii) What is the probability that during a given 1-minute interval, exactly 2 of the 5 operators receives no calls? (2+2+3)
- (b) Suppose that the number of miles that a car can run before its battery wears out is exponentially distributed with an average value of 10,000 miles. If a person desires to take a 5000-mile trip, what is the probability that he or she will be able to complete the trip without having to replace the car battery? (3)
- (क) एक टेलीफोन एक्सचेंज डेस्क 5 ऑपरेटरों को नियुक्त करता है जो पॉइसन प्रक्रिया के अनुसार 4 कॉल प्रति मिनट की दर से एक दूसरे से स्वतंत्र रूप से कॉल प्राप्त करते हैं।

- (i) इसकी क्या प्रायिकता है कि दिए गए 1 मिनट के अंतराल के दौरान, पहले ऑपरेटर को कोई कॉल न मिले?
- (ii) इसकी क्या प्रायिकता है कि दिए गए 45 - सेकंड के अंतराल के दौरान, दूसरे ऑपरेटर को कोई कॉल न मिले?
- (iii) इसकी क्या प्रायिकता है कि दिए गए 1 - मिनट के अंतराल के दौरान, 5 में से ठीक 2 ऑपरेटरों को कोई कॉल न मिले?
- (iv) मान लीजिए कि बैटरी खराब होने से पहले एक कार जितनी मील चल सकती है, वह 10,000 मील के औसत मूल्य के साथ तेजी से वितरित की जाती है। यदि कोई व्यक्ति 5000 मील की यात्रा करना चाहता है, तो क्या संभावना है कि वह कार की बैटरी को बदले बिना यात्रा पूरी कर पाएगा?
5. Suppose X and Y are two discrete random variables which have the joint probability mass function as given below

		Y				
		0	1	2	3	
X	0	$\frac{1}{8}$	$\frac{1}{8}$	$\frac{1}{16}$	$\frac{1}{16}$	
	1	0	$\frac{1}{4}$	$\frac{1}{4}$	$\frac{1}{8}$	

- (i) Find the marginal distribution of X and Y.
- (ii) Find Variance of Y.
- (iii) Find the conditional distribution of Y given X=0.
- (iv) Find the expected value of  $h(X, Y)$  where  $h(X, Y) = 2X - 5XY$
- (2+3+2+3)

मान लीजिए कि X और Y दो असतत यादृच्छिक चर हैं जिनका संयुक्त संभाव्यता द्रव्यमान फलन है जैसा कि नीचे दिया गया है

X	Y				
	0	1	2	3	
0	$\frac{1}{8}$	$\frac{1}{8}$	$\frac{1}{16}$	$\frac{1}{16}$	
1	0	$\frac{1}{4}$	$\frac{1}{4}$	$\frac{1}{8}$	

- (i) X और Y का सीमांत वितरण ज्ञात कीजिए।
- (ii) Y का प्रसरण ज्ञात कीजिए।
- (iii) दिए गए X=0 पर Y का सशर्त वितरण ज्ञात कीजिए।
- (iv) h(X,Y) का अपेक्षित मान ज्ञात कीजिए, जहाँ  $h(X,Y) = 2X - 5XY$  है।
6. (a) The amount of time, in minutes, that a person must wait for a bus is uniformly distributed between zero and 15 minutes, inclusive.
- (i) On the average, how long must a person wait? Also find the standard deviation,  $\sigma$ .
- (ii) What value c is such that 90% of waiting times are below it?
- (2+2)
- (b) 70% of all students in a certain University attend classes regularly (with a minimum attendance of 90%). 80% of these scores a CGPA of at least 9, while a student who is not regular scores a CGPA of 9 or more only 20% of the time. Consider randomly selecting 5 students.
- (i) What is the probability that exactly 3 of these are regular and score a CGPA of 9 or more?
- (ii) What is the probability that exactly 4 of these scores a CGPA of 9 or more?
- (3+3)
- (क) एक व्यक्ति को बस के लिए जितना समय, मिनटों में इंतजार करना पड़ता है, वह समान रूप से शून्य और 15 मिनट, जिसमें दोनों शामिल है, के बीच वितरित किया जाता है।

- (i) औसतन एक व्यक्ति को कितने समय तक प्रतीक्षा करनी चाहिए? मानक विचलन ०, भी खोजिये।
- (ii) c का क्या मान है कि 90% प्रतीक्षा समय इससे कम है?
- (ख) एक निश्चित विश्वविद्यालय में सभी छात्रों में से 70% नियमित रूप से कक्षाओं में भाग लेते हैं (न्यूनतम 90% उपस्थिति के साथ)। इनमें से 80% ने कम से कम ९ का सीजीपीए स्कोर किया है, जबकि एक छात्र जो नियमित रूप से ९ या उससे अधिक का सीजीपीए स्कोर नहीं करता है, वह केवल 20% नियमित है। यादृच्छिक रूप से ५ छात्रों का चयन करने पर विचार कीजिये।
- (i) इसकी क्या प्रायिकता है कि इनमें से ठीक ३ नियमित हैं और ९ या अधिक का सीजीपीए प्राप्त करते हैं?
- (ii) इसकी क्या प्रायिकता है कि इनमें से ठीक ४ का सीजीपीए ९ या अधिक है?

### SECTION III

*Attempt any two questions from Q7, Q8 and Q9.*

**Q7, Q8 और Q9 में से किन्हीं दो प्रश्नों का उत्तर दीजिये।**

7. (a) Suppose Y is a random variable with uniform distribution on the interval from  $\alpha$  to a known upper limit 20. It is desired to estimate  $\alpha$  on the basis of a random sample of size n. Explain intuitively, why the smallest value in the sample is not an unbiased estimator of  $\alpha$ . (4)
- (b) Suppose  $X_1, X_2, \dots, X_n$  is a random sample from an exponential distribution with parameter  $\lambda$ . Find the estimator of  $\lambda$  using maximum likelihood method? (4)
- (c) State the invariance principle of the Maximum Likelihood Estimators. Give an example. (2)
- (क) मान लीजिए कि Y एक यादृच्छिक चर है जिसमें  $\alpha$  से ज्ञात ऊपरी सीमा 20 के अंतराल पर समान वितरण होता है। आकार n के यादृच्छिक नमूने के आधार पर  $\alpha$  का अनुमान लगाना बाधित है। सहज रूप से समझाएं कि नमूने में सबसे छोटा मान  $\alpha$  का निष्पक्ष अनुमानक क्यों नहीं है?

- (ख) मान लीजिए  $X_1, X_2, \dots, X_n$   $\lambda$  पैरामीटर के साथ एक घातीय वितरण से एक यादृच्छिक नमूना है। अधिकतम प्रायिकता विधि का उपयोग करके  $\lambda$  का अनुमानक ज्ञात कीजिए?
- (ग) अधिकतम संभाव्यता अनुमानकों के अपरिवर्तनशीलता सिद्धांत का उल्लेख कीजिए। एक उदाहरण दीजिये।
8. (a) Assume that the time saved by travelling through metro in Delhi is normally distributed with true standard deviation 0.75.
- Compute 96% confidence interval for the true average time saved based on sample of 35 people , with a sample average time saved being 8.57 minutes.
  - Will your answer change if time saved by travelling through metro is not normally distributed?
  - How large must be the sample size if the width of the 92% interval is to be 0.50. (3+2+2)
- (b) Out of a random sample of 150 Britishers, 108 Britishers have blue eyes. With 94% confidence what can we say about the maximum error in the estimate of true proportion of Britishers who have blue eyes, irrespective of the sample size? (3)
- (क) मान लीजिये कि दिल्ली में मेट्रो से यात्रा करने से बचा हुआ समय सामान्य रूप से सही मानक विचलन 0.75 के साथ वितरित किया जाता है।
- 35 लोगों के नमूने के आधार पर बचाए गए वास्तविक औसत समय के लिए 96% विश्वास अंतराल की गणना कीजिये, जिसमें नमूना औसत समय 8.57 मिनट बचा हो।
  - यदि मेट्रो से यात्रा करके बचाए गए समय को सामान्य रूप से वितरित नहीं किया जाता है तो क्या आपका उत्तर बदल जाएगा?
  - यदि 92% के अंतराल की चौड़ाई 0.50 होनी है तो नमूना आकार कितना बड़ा होना चाहिए।
- (ख) 150 अंग्रेजों के यादृच्छिक नमूने में से 108 अंग्रेजों की आंखें नीली हैं। 94% विश्वास के साथ, नमूने के आकार की परवाह किए बिना, नीली आंखों वाले अंग्रेजों के वास्तविक अनुपात के अनुमान में अधिकतम त्रुटि के बारे में हम क्या कह सकते हैं?

9. (a) Let  $X$  denote the number of hours in a day that a randomly selected child watches TV. Suppose the probability density function of  $X$  is

$$f(x) = \begin{cases} \frac{\beta}{x^{\beta+1}} & x > 1 \\ 0 & \text{otherwise.} \end{cases}$$

A random sample of 5 children is taken with the following observations 1.3, 2.4, 1.1, 2.5 and 3.0. Use the method of moments to obtain an estimator of  $\beta$  and then compute the estimate of this data. (5)

- (b) A sample of 20 cigarettes is tested to determine nicotine content and the average value observed was 1.2 mg. Compute a 99 percent two-sided confidence interval for the mean nicotine content of a cigarette if the population variance is not known in advance of the experiment and the sample variance is 0.04. Also compute an upper bound at 99% confidence level for the true mean nicotine content of a cigarette. (3+2)

(क) मान लीजिए कि  $X$  एक दिन में उन घंटों की संख्या को दर्शाता है जो एक यादृच्छिक रूप से चयनित बच्चा टीवी देखता है। मान लीजिए कि  $X$  का प्रायिकता घनत्व फलन है

$$f(x) = \begin{cases} \frac{\beta}{x^{\beta+1}} & x > 1 \\ 0 & \text{अन्यथा.} \end{cases}$$

निम्नलिखित प्रेक्षणों 1.3, 2.4, 1.1, 2.5 और 3.0 के साथ 5 बच्चों का एक यादृच्छिक नमूना लिया जाता है।  $\beta$  का अनुमानक प्राप्त करने के लिए क्षणों की विधि का उपयोग कीजिए और फिर इस डेटा के अनुमान की गणना कीजिए।

- (ख) निकोटीन की मात्रा निर्धारित करने के लिए 20 सिगरेट के एक नमूने का परीक्षण किया गया और औसतन 1.2 मिलीग्राम निकोटीन देखा गया। एक सिगरेट की औसत निकोटीन सामग्री के लिए 99 प्रतिशत दो-तरफा आत्मविश्वास अंतराल की गणना कीजिए यदि जनसंख्या भिन्नता प्रयोग के पहले से ज्ञात नहीं है और नमूना भिन्नता 0.04 है। एक सिगरेट के वास्तविक माध्य निकोटीन सामग्री के लिए 99% आत्मविश्वास स्तर पर ऊपरी सीमा की गणना भी कीजिए।

#### SECTION IV

*Attempt any two questions of Q10, Q11 and Q12.*

**Q10, Q11 और Q12 में से किन्हीं दो प्रश्नों का उत्तर दीजिये।**

10. Up to 4 ppm (parts per million) of chlorine in drinking water is considered not only safe but useful for preventing contamination but higher than that is harmful. Following reports of a spike in the chlorine levels in drinking water in a municipality, the concerned regulatory agency wanted to ensure that the Jal Board is sticking to the guidelines and for this a random sample consisting of 100 collections from different parts of the municipality were tested.

- (i) Set up the appropriate null and alternative hypothesis. The mean chlorine content of the 100 samples collected was 3.8 ppm. Assuming that the distribution of the chlorine content is normal with  $\sigma = 0.9$ , carry out test at the 0.01 and 0.05 levels of significance.
- (ii) Clearly writing down the rejection rule for the 5% significance test, compute  $\beta (3.92)$  and the power of the test.
- (iii) Do the probabilities of Type I error and Type II error always add up to one?  
Explain your answer. (4+4+2)

पीने के पानी में क्लोरीन के 4 पीपीएम (प्रति मिलियन भाग) को न केवल सुरक्षित, बल्कि संदूषण को रोकने के लिए उपयोगी माना जाता है, लेकिन इससे अधिक मात्रा हानिकारक होती है। नगरपालिका में पीने के पानी में क्लोरीन के स्तर में एक बढ़ावत की रिपोर्ट के बाद, संबंधित नियामक एजेंसी यह सुनिश्चित करना चाहती थी कि जल बोर्ड दिशानिर्देशों का पालन कर रहे हैं या नहीं। इसके लिए नगरपालिका के विभिन्न हिस्सों से 100 संग्रह से युक्त एक यादृच्छिक नमूने का परीक्षण किया गया।

- (i) उपयुक्त अशक्त और वैकल्पिक परिकल्पना सेट कीजिये। एकत्र किए गए 100 नमूनों की औसत क्लोरीन सामग्री 3.8 पीपीएम (प्रति मिलियन भाग) थी। यह मानते हुए कि क्लोरीन सामग्री का वितरण 0.9 के साथ सामान्य है, 0.01 और 0.05 के महत्व के स्तरों पर परीक्षण कीजिये।

- (ii) स्पष्ट रूप से 5% महत्व परीक्षण के लिए अस्वीकृति नियम को लिखिए।  $\beta$  (3.92) और परीक्षण की शक्ति की गणना कीजिए।
- (iii) क्या टाइप I त्रुटि की संभावना और टाइप II त्रुटि की संभावना का जोड़ इकाई होता है? अपना जवाब समझाएं।
11. (a) The weight of biscuits mentioned on its packet may not be the true weight. Let the true average weight of all packets be 20 grams. However, sellers claim that the true average weight is more than 20 grams. To test this a sample of 4 packets of biscuits are selected and their average weight was found to be 20.264 grams with a standard deviation of 1.88 grams. How would you carry out the above test? Which distribution would you use and why? Compute the p-value of the test. (5)

- (b) A random variable with unknown mean may be assumed to have a normal distribution with variance 9. Given a sample of 81 observations derive the non-rejection and the rejection regions for  $X$  using 6% significance test. If the level of significance is increased to 10%, will the rejection region increase or decrease? Explain. (5)

(क) हो सकता है कि बिस्किट के पैकेट पर लिखे बिस्किट का वजन सही वजन न हो। मान लीजिए सभी पैकेटों का वास्तविक औसत वजन 20 ग्राम है। हालांकि, विक्रेताओं का दावा है कि वास्तविक औसत वजन 20 ग्राम से अधिक है। इसका परीक्षण करने के लिए बिस्कुट के 4 पैकेटों का एक नमूना चुना गया और उनका औसत वजन 1.88 ग्राम के मानक विचलन के साथ 20.264 ग्राम पाया गया। आप उपरोक्त परीक्षण कैसे करेंगे? आप किस वितरण का उपयोग करेंगे और क्यों? परीक्षण के पी-मान की गणना कीजिए।

(ख) अज्ञात माध्य के साथ एक यादृच्छिक चर को विचरण 9 के साथ एक सामान्य वितरण माना जा सकता है। 81 टिप्पणियों के एक नमूने को देखते हुए 6% महत्व परीक्षण का उपयोग करके  $X$  के लिए गैर-अस्वीकृति और अस्वीकृति क्षेत्रों को प्राप्त कीजिए।

12. (a) A Company produces coffee and sells it to consumers in jars of 250 grams. Answer the following, with suitable justification :

- (i) Set up the appropriate hypotheses to be tested by a consumer if he suspects the amount of coffee filled in the jar
- (ii) Set up the appropriate hypotheses to be tested by the owner of the coffee company if he is suspicious of the amount of coffee filled in the jars
- (iii) Set up the appropriate hypotheses to be tested by the inspecting officer if he wants to ensure of the accuracy of the amount of coffee filled in the jars. (6)
- (b) A manufacturer of crystal glasses randomly selects 100 and determines that 14 of them are scratched. Does this provide compelling evidence for concluding that more than 10% have a scratch? State and test the relevant hypotheses using a level of significance of 8%. In reaching your conclusion, what type of error might you have committed? (4)
- (क) एक कंपनी कॉफी का उत्पादन करती है और इसे 250 ग्राम के जार में उपभोक्ताओं को बेचती है। उपयुक्त औचित्य के साथ, निम्नलिखित उत्तर दीजिये :
- (i) यदि एक उपभोक्ता जार में भरी हुई कॉफी की मात्रा पर संदेह करता है, तो उपभोक्ता द्वारा परीक्षण किए जाने के लिए उपयुक्त परिकल्पना स्थापित कीजिये।
  - (ii) जार में भरी हुई कॉफी की मात्रा पर संदेह होने पर कंपनी के मालिक द्वारा परीक्षण किए जाने के लिए उपयुक्त परिकल्पनाएं निर्धारित कीजिये।
  - (iii) निरीक्षण अधिकारी द्वारा परीक्षण किए जाने के लिए उपयुक्त परिकल्पनाएं निर्धारित कीजिये यदि वह जार में भरी हुई कॉफी की मात्रा सुनिश्चित करना चाहता है।
- (ख) क्रिस्टल ग्लास का एक निर्माता यादृच्छिक रूप से 100 का चयन करता है और निर्धारित करता है कि उनमें से 14 में खरोंच हैं। क्या यह निष्कर्ष निकालने के लिए सम्मोहक साक्ष्य प्रदान करता है कि 10% से अधिक में खरोंच है? 8% के महत्व के स्तर का उपयोग करके प्रासंगिक परिकल्पनाओं को राज्य और परीक्षण कीजिये। अपने निष्कर्ष पर पहुँचने में, आपने किस प्रकार की त्रुटि की होगी?

## A-2 Appendix Tables

**Table A.1 Cumulative Binomial Probabilities**

$$B(x; n, p) = \sum_{y=0}^x b(y; n, p)$$

a.  $n = 5$ 

		<i>p</i>														
		0.01	0.05	0.10	0.20	0.25	0.30	0.40	0.50	0.60	0.70	0.75	0.80	0.90	0.95	0.99
<i>x</i>	0	.951	.774	.590	.328	.237	.168	.078	.031	.010	.002	.001	.000	.000	.000	.000
	1	.999	.977	.919	.737	.633	.528	.337	.188	.087	.031	.016	.007	.000	.000	.000
	2	1.000	.999	.991	.942	.896	.837	.683	.500	.317	.163	.044	.058	.009	.001	.000
	3	1.000	1.000	1.000	.993	.984	.969	.913	.812	.663	.472	.367	.263	.081	.023	.001
	4	1.000	1.000	1.000	1.000	.999	.998	.990	.969	.922	.832	.763	.672	.410	.226	.049

b.  $n = 10$ 

		<i>p</i>														
		0.01	0.05	0.10	0.20	0.25	0.30	0.40	0.50	0.60	0.70	0.75	0.80	0.90	0.95	0.99
<i>x</i>	0	.904	.599	.349	.107	.056	.028	.006	.001	.000	.000	.000	.000	.000	.000	.000
	1	.996	.914	.736	.376	.244	.149	.046	.011	.002	.000	.000	.000	.000	.000	.000
	2	1.000	.988	.930	.678	.526	.383	.167	.055	.012	.002	.000	.000	.000	.000	.000
	3	1.000	.999	.987	.879	.776	.650	.382	.172	.055	.011	.004	.001	.000	.000	.000
	4	1.000	1.000	.998	.967	.922	.850	.633	.377	.166	.047	.020	.006	.000	.000	.000
	5	1.000	1.000	1.000	.994	.980	.953	.834	.623	.367	.150	.078	.033	.002	.000	.000
	6	1.000	1.000	1.000	.999	.996	.989	.945	.828	.618	.350	.224	.121	.013	.001	.000
	7	1.000	1.000	1.000	1.000	1.000	.998	.988	.945	.833	.617	.474	.322	.070	.012	.000
	8	1.000	1.000	1.000	1.000	1.000	1.000	.998	.989	.954	.851	.756	.624	.264	.086	.004
	9	1.000	1.000	1.000	1.000	1.000	1.000	1.000	.999	.994	.972	.944	.893	.651	.401	.096

c.  $n = 15$ 

		<i>p</i>														
		0.01	0.05	0.10	0.20	0.25	0.30	0.40	0.50	0.60	0.70	0.75	0.80	0.90	0.95	0.99
<i>x</i>	0	.860	.463	.206	.035	.013	.005	.000	.000	.000	.000	.000	.000	.000	.000	.000
	1	.990	.829	.549	.167	.080	.035	.005	.000	.000	.000	.000	.000	.000	.000	.000
	2	1.000	.964	.816	.398	.236	.127	.027	.004	.000	.000	.000	.000	.000	.000	.000
	3	1.000	.995	.944	.648	.461	.297	.091	.018	.002	.000	.000	.000	.000	.000	.000
	4	1.000	.999	.987	.836	.686	.515	.217	.059	.009	.001	.000	.000	.000	.000	.000
	5	1.000	1.000	.998	.939	.852	.722	.403	.151	.034	.004	.001	.000	.000	.000	.000
	6	1.000	1.000	1.000	.982	.943	.869	.610	.304	.095	.015	.004	.001	.000	.000	.000
	7	1.000	1.000	1.000	.996	.983	.950	.787	.500	.213	.050	.017	.004	.000	.000	.000
	8	1.000	1.000	1.000	.999	.996	.985	.905	.696	.390	.131	.057	.018	.000	.000	.000
	9	1.000	1.000	1.000	1.000	.999	.996	.966	.849	.597	.278	.148	.061	.002	.000	.000
<i>x</i>	10	1.000	1.000	1.000	1.000	1.000	.999	.991	.941	.783	.485	.314	.164	.013	.001	.000
	11	1.000	1.000	1.000	1.000	1.000	1.000	.998	.982	.909	.703	.539	.352	.056	.005	.000
	12	1.000	1.000	1.000	1.000	1.000	1.000	1.000	.996	.973	.873	.764	.602	.184	.036	.000
	13	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	.995	.965	.920	.833	.451	.171	.010
	14	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	.995	.987	.965	.794	.537	.140

(continued)

**Table A.1 Cumulative Binomial Probabilities (cont.)**d.  $n = 20$ 

$$B(x; n, p) = \sum_{y=0}^x b(y; n, p)$$

	<i>p</i>														
	0.01	0.05	0.10	0.20	0.25	0.30	0.40	0.50	0.60	0.70	0.75	0.80	0.90	0.95	0.99
0	.818	.358	.122	.012	.003	.001	.000	.000	.000	.000	.000	.000	.000	.000	.000
1	.983	.736	.392	.069	.024	.008	.001	.000	.000	.000	.000	.000	.000	.000	.000
2	.999	.925	.677	.206	.091	.035	.004	.000	.000	.000	.000	.000	.000	.000	.000
3	1.000	.984	.867	.411	.225	.107	.016	.001	.000	.000	.000	.000	.000	.000	.000
4	1.000	.997	.957	.630	.415	.238	.051	.006	.000	.000	.000	.000	.000	.000	.000
5	1.000	1.000	.989	.804	.617	.416	.126	.021	.002	.000	.000	.000	.000	.000	.000
6	1.000	1.000	.998	.913	.786	.608	.250	.058	.006	.000	.000	.000	.000	.000	.000
7	1.000	1.000	1.000	.968	.898	.772	.416	.132	.021	.001	.000	.000	.000	.000	.000
8	1.000	1.000	1.000	.990	.959	.887	.596	.252	.057	.005	.001	.000	.000	.000	.000
9	1.000	1.000	1.000	.997	.986	.952	.755	.412	.128	.017	.004	.001	.000	.000	.000
x	10	1.000	1.000	1.000	.999	.996	.983	.872	.588	.245	.048	.014	.003	.000	.000
11	1.000	1.000	1.000	1.000	.999	.995	.943	.748	.404	.113	.041	.010	.000	.000	.000
12	1.000	1.000	1.000	1.000	1.000	.999	.979	.868	.584	.228	.102	.032	.000	.000	.000
13	1.000	1.000	1.000	1.000	1.000	1.000	.994	.942	.750	.392	.214	.087	.002	.000	.000
14	1.000	1.000	1.000	1.000	1.000	1.000	.998	.979	.874	.584	.383	.196	.011	.000	.000
15	1.000	1.000	1.000	1.000	1.000	1.000	1.000	.994	.949	.762	.585	.370	.043	.003	.000
16	1.000	1.000	1.000	1.000	1.000	1.000	1.000	.999	.984	.893	.775	.589	.33	.016	.000
17	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	.996	.965	.909	.794	.523	.075	.001
18	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	.999	.992	.976	.931	.608	.264	.017
19	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	.999	.997	.988	.878	.642	.182

(continued)

## A-4 Appendix Tables

**Table A.1 Cumulative Binomial Probabilities (cont.)**

$$B(x; n, p) = \sum_{y=0}^x b(y; n, p)$$

e.  $n = 25$ 

	<i>p</i>														
	0.01	0.05	0.10	0.20	0.25	0.30	0.40	0.50	0.60	0.70	0.75	0.80	0.90	0.95	0.99
0	.778	.277	.072	.004	.001	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000
1	.974	.642	.271	.027	.007	.002	.000	.000	.000	.000	.000	.000	.000	.000	.000
2	.998	.873	.537	.098	.032	.009	.000	.000	.000	.000	.000	.000	.000	.000	.000
3	1.000	.966	.764	.234	.096	.033	.002	.000	.000	.000	.000	.000	.000	.000	.000
4	1.000	.993	.902	.421	.214	.090	.009	.000	.000	.000	.000	.000	.000	.000	.000
5	1.000	.999	.967	.617	.378	.193	.029	.002	.000	.000	.000	.000	.000	.000	.000
6	1.000	1.000	.991	.780	.561	.341	.074	.007	.000	.000	.000	.000	.000	.000	.000
7	1.000	1.000	.998	.891	.727	.512	.154	.022	.001	.000	.000	.000	.000	.000	.000
8	1.000	1.000	1.000	.953	.851	.677	.274	.054	.004	.000	.000	.000	.000	.000	.000
9	1.000	1.000	1.000	.983	.929	.811	.425	.115	.013	.000	.000	.000	.000	.000	.000
10	1.000	1.000	1.000	.994	.970	.902	.586	.212	.034	.002	.000	.000	.000	.000	.000
11	1.000	1.000	1.000	.998	.980	.956	.732	.345	.078	.006	.001	.000	.000	.000	.000
x	12	1.000	1.000	1.000	1.000	.997	.983	.846	.500	.154	.017	.003	.000	.000	.000
13	1.000	1.000	1.000	1.000	.999	.994	.922	.655	.268	.044	.020	.002	.000	.000	.000
14	1.000	1.000	1.000	1.000	.998	.966	.788	.414	.098	.030	.006	.000	.000	.000	.000
15	1.000	1.000	1.000	1.000	1.000	.987	.885	.575	.189	.071	.017	.000	.000	.000	.000
16	1.000	1.000	1.000	1.000	1.000	1.000	.996	.946	.726	.323	.149	.047	.000	.000	.000
17	1.000	1.000	1.000	1.000	1.000	1.000	.999	.978	.846	.488	.273	.109	.002	.000	.000
18	1.000	1.000	1.000	1.000	1.000	1.000	1.000	.993	.926	.659	.439	.220	.099	.000	.000
19	1.000	1.000	1.000	1.000	1.000	1.000	1.000	.998	.971	.807	.622	.383	.033	.001	.000
20	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	.991	.910	.786	.579	.098	.007	.000
21	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	.998	.967	.904	.766	.236	.034	.000
22	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	.991	.968	.902	.463	.127	.002
23	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	.998	.993	.973	.729	.358	.026
24	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	.999	.996	.928	.723	.222

**Table A.2 Cumulative Poisson Probabilities**

$$F(x; \mu) = \sum_{y=0}^x \frac{e^{-\mu} \mu^y}{y!}$$


	<i>μ</i>										
	.1	.2	.3	.4	.5	.6	.7	.8	.9	1.0	
0	.905	.819	.741	.670	.607	.549	.497	.449	.407	.368	
1	.995	.982	.963	.938	.910	.878	.844	.809	.772	.736	
x	2	1.000	.999	.996	.992	.986	.977	.966	.953	.937	.920
3		1.000	1.000	.999	.998	.997	.994	.991	.987	.981	
4				1.000	1.000	1.000	.999	.999	.998	.996	
5							1.000	1.000	1.000	.999	
6										1.000	

(continued)

## A-6 Appendix Tables

Table A.3 Standard Normal Curve Areas

$$(z) = P(Z \leq z)$$




$z$	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
-3.4	.0003	.0003	.0003	.0003	.0003	.0003	.0003	.0003	.0003	.0002
-3.3	.0005	.0005	.0005	.0004	.0004	.0004	.0004	.0004	.0004	.0003
-3.2	.0007	.0007	.0006	.0006	.0006	.0006	.0006	.0005	.0005	.0005
-3.1	.0010	.0009	.0009	.0009	.0008	.0008	.0008	.0008	.0007	.0007
-3.0	.0013	.0013	.0013	.0012	.0012	.0011	.0011	.0011	.0010	.0010
-2.9	.0019	.0018	.0017	.0017	.0016	.0016	.0015	.0015	.0014	.0014
-2.8	.0026	.0025	.0024	.0023	.0023	.0022	.0021	.0021	.0020	.0019
-2.7	.0035	.0034	.0033	.0032	.0031	.0030	.0029	.0028	.0027	.0026
-2.6	.0047	.0045	.0044	.0043	.0041	.0040	.0039	.0038	.0037	.0036
-2.5	.0062	.0060	.0059	.0057	.0055	.0054	.0052	.0051	.0049	.0038
-2.4	.0082	.0080	.0078	.0075	.0073	.0071	.0069	.0068	.0066	.0064
-2.3	.0107	.0104	.0102	.0099	.0096	.0094	.0091	.0089	.0087	.0084
-2.2	.0139	.0136	.0132	.0129	.0125	.0122	.0119	.0116	.0113	.0110
-2.1	.0179	.0174	.0170	.0166	.0162	.0158	.0154	.0150	.0146	.0143
-2.0	.0228	.0222	.0217	.0212	.0207	.0202	.0197	.0192	.0188	.0183
-1.9	.0287	.0281	.0274	.0268	.0262	.0256	.0250	.0244	.0239	.0233
-1.8	.0359	.0352	.0344	.0336	.0329	.0322	.0314	.0307	.0301	.0294
-1.7	.0446	.0436	.0427	.0418	.0409	.0401	.0392	.0384	.0375	.0367
-1.6	.0548	.0537	.0526	.0516	.0505	.0495	.0485	.0475	.0465	.0455
-1.5	.0668	.0655	.0643	.0630	.0618	.0606	.0594	.0582	.0571	.0559
-1.4	.0808	.0793	.0778	.0764	.0749	.0735	.0722	.0708	.0694	.0681
-1.3	.0968	.0951	.0934	.0918	.0901	.0885	.0869	.0853	.0838	.0823
-1.2	.1151	.1131	.1112	.1093	.1075	.1056	.1038	.1020	.1003	.0985
-1.1	.1357	.1335	.1314	.1292	.1271	.1251	.1230	.1210	.1190	.1170
-1.0	.1587	.1562	.1539	.1515	.1492	.1469	.1446	.1423	.1401	.1379
-0.9	.1841	.1814	.1788	.1762	.1736	.1711	.1685	.1660	.1635	.1611
-0.8	.2119	.2090	.2061	.2033	.2005	.1977	.1949	.1922	.1894	.1867
-0.7	.2420	.2389	.2358	.2327	.2296	.2266	.2236	.2206	.2177	.2148
-0.6	.2743	.2709	.2676	.2643	.2611	.2578	.2546	.2514	.2483	.2451
-0.5	.3085	.3050	.3015	.2981	.2946	.2912	.2877	.2843	.2810	.2776
-0.4	.3446	.3409	.3372	.3336	.3300	.3264	.3228	.3192	.3156	.3121
-0.3	.3821	.3783	.3745	.3707	.3669	.3632	.3594	.3557	.3520	.3482
-0.2	.4207	.4168	.4129	.4090	.4052	.4013	.3974	.3936	.3897	.3859
-0.1	.4602	.4562	.4522	.4483	.4443	.4404	.4364	.4325	.4286	.4247
-0.0	.5000	.4960	.4920	.4880	.4840	.4801	.4761	.4721	.4681	.4641

(continued)

Table A.3 Standard Normal Curve Areas (cont.)


 $\Phi(z) = P(Z \leq z)$ 

<i>z</i>	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
0.0	.5000	.5040	.5080	.5120	.5160	.5199	.5239	.5279	.5319	.5359
0.1	.5398	.5438	.5478	.5517	.5557	.5596	.5636	.5675	.5714	.5753
0.2	.5793	.5832	.5871	.5910	.5948	.5987	.6026	.6064	.6103	.6141
0.3	.6179	.6217	.6255	.6293	.6331	.6368	.6406	.6443	.6480	.6517
0.4	.6554	.6591	.6628	.6664	.6700	.6736	.6772	.6808	.6844	.6879
0.5	.6915	.6950	.6985	.7019	.7054	.7088	.7123	.7157	.7190	.7224
0.6	.7257	.7291	.7324	.7357	.7389	.7422	.7454	.7486	.7517	.7549
0.7	.7580	.7611	.7642	.7673	.7704	.7734	.7764	.7794	.7823	.7852
0.8	.7881	.7910	.7939	.7967	.7995	.8023	.8051	.8078	.8106	.8133
0.9	.8159	.8186	.8212	.8238	.8264	.8289	.8315	.8340	.8365	.8389
1.0	.8413	.8438	.8461	.8485	.8508	.8531	.8554	.8577	.8599	.8621
1.1	.8643	.8665	.8686	.8708	.8729	.8749	.8770	.8790	.8810	.8830
1.2	.8849	.8869	.8888	.8907	.8925	.8944	.8962	.8980	.8997	.9015
1.3	.9032	.9049	.9066	.9082	.9099	.9115	.9131	.9147	.9162	.9177
1.4	.9192	.9207	.9222	.9236	.9251	.9265	.9278	.9292	.9306	.9319
1.5	.9332	.9345	.9357	.9370	.9382	.9394	.9406	.9418	.9429	.9441
1.6	.9452	.9463	.9474	.9484	.9495	.9505	.9515	.9525	.9535	.9545
1.7	.9554	.9564	.9573	.9582	.9591	.9599	.9608	.9616	.9625	.9633
1.8	.9641	.9649	.9656	.9664	.9671	.9678	.9686	.9693	.9699	.9706
1.9	.9713	.9719	.9726	.9732	.9738	.9744	.9750	.9756	.9761	.9767
2.0	.9772	.9778	.9783	.9788	.9793	.9798	.9803	.9808	.9812	.9817
2.1	.9821	.9826	.9830	.9834	.9838	.9842	.9846	.9850	.9854	.9857
2.2	.9861	.9864	.9868	.9871	.9875	.9878	.9881	.9884	.9887	.9890
2.3	.9893	.9896	.9898	.9901	.9904	.9906	.9909	.9911	.9913	.9916
2.4	.9918	.9920	.9922	.9925	.9927	.9929	.9931	.9932	.9934	.9936
2.5	.9938	.9940	.9941	.9943	.9945	.9946	.9948	.9949	.9951	.9952
2.6	.9953	.9955	.9956	.9957	.9959	.9960	.9961	.9962	.9963	.9964
2.7	.9965	.9966	.9967	.9968	.9969	.9970	.9971	.9972	.9973	.9974
2.8	.9974	.9975	.9976	.9977	.9977	.9978	.9979	.9979	.9980	.9981
2.9	.9981	.9982	.9982	.9983	.9984	.9984	.9985	.9985	.9986	.9986
3.0	.9987	.9987	.9987	.9988	.9988	.9989	.9989	.9989	.9990	.9990
3.1	.9990	.9991	.9991	.9991	.9992	.9992	.9992	.9992	.9993	.9993
3.2	.9993	.9993	.9994	.9994	.9994	.9994	.9994	.9995	.9995	.9995
3.3	.9995	.9995	.9995	.9996	.9996	.9996	.9996	.9996	.9996	.9997
3.4	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9998

Table A.5 Critical Values for  $t$  Distributions

$v$	$\alpha$						
	.10	.05	.025	.01	.005	.001	.0005
1	3.078	6.314	12.706	31.821	63.657	318.31	636.62
2	1.886	2.920	4.303	6.965	9.925	22.326	31.598
3	1.638	2.353	3.182	4.541	5.841	10.213	12.924
4	1.533	2.132	2.776	3.747	4.604	7.173	8.610
5	1.476	2.015	2.571	3.365	4.032	5.893	6.869
6	1.440	1.943	2.447	3.143	3.707	5.208	5.959
7	1.415	1.895	2.365	2.998	3.499	4.785	5.408
8	1.397	1.860	2.306	2.896	3.355	4.501	5.041
9	1.383	1.833	2.262	2.821	3.250	4.297	4.781
10	1.372	1.812	2.228	2.764	3.169	4.144	4.587
11	1.363	1.796	2.201	2.718	3.106	4.025	4.437
12	1.356	1.782	2.179	2.681	3.055	3.930	4.318
13	1.350	1.771	2.160	2.650	3.012	3.852	4.221
14	1.345	1.761	2.145	2.624	2.977	3.787	4.140
15	1.341	1.753	2.131	2.602	2.947	3.733	4.073
16	1.337	1.746	2.120	2.583	2.921	3.686	4.015
17	1.333	1.740	2.110	2.567	2.898	3.646	3.965
18	1.330	1.734	2.101	2.552	2.878	3.610	3.922
19	1.328	1.729	2.093	2.539	2.861	3.579	3.883
20	1.325	1.725	2.086	2.528	2.845	3.552	3.850
21	1.323	1.721	2.080	2.518	2.831	3.527	3.819
22	1.321	1.717	2.074	2.508	2.819	3.505	3.792
23	1.319	1.714	2.069	2.500	2.807	3.485	3.767
24	1.318	1.711	2.064	2.492	2.797	3.467	3.745
25	1.316	1.708	2.060	2.485	2.787	3.450	3.725
26	1.315	1.706	2.056	2.479	2.779	3.435	3.707
27	1.314	1.703	2.052	2.473	2.771	3.421	3.690
28	1.313	1.701	2.048	2.467	2.763	3.408	3.674
29	1.311	1.699	2.045	2.462	2.756	3.396	3.659
30	1.310	1.697	2.042	2.457	2.750	3.385	3.646
32	1.309	1.694	2.037	2.449	2.738	3.365	3.622
34	1.307	1.691	2.032	2.441	2.728	3.348	3.601
36	1.306	1.688	2.028	2.434	2.719	3.333	3.582
38	1.304	1.686	2.024	2.429	2.712	3.319	3.566
40	1.303	1.684	2.021	2.423	2.704	3.307	3.551
50	1.299	1.676	2.009	2.403	2.678	3.262	3.496
60	1.296	1.671	2.000	2.390	2.660	3.232	3.460
120	1.289	1.658	1.980	2.358	2.617	3.160	3.373
	1.282	1.645	1.960	2.326	2.576	3.090	3.291

Table A.7 Critical Values for Chi-Squared Distributions



$\nu$	.995	.99	.975	.95	.90	.10	.05	.025	.01	.005
1	0.000	0.000	0.001	0.004	0.016	2.706	3.843	5.025	6.637	7.882
2	0.010	0.020	0.051	0.103	0.211	4.605	5.992	7.378	9.210	10.597
3	0.072	0.115	0.216	0.352	0.584	6.251	7.815	9.348	11.344	12.837
4	0.207	0.297	0.484	0.711	1.064	7.779	9.488	11.143	13.277	14.860
5	0.412	0.554	0.831	1.145	1.610	9.236	11.070	12.832	15.085	16.748
6	0.676	0.872	1.237	1.635	2.204	10.645	12.592	14.440	16.812	18.548
7	0.989	1.239	1.690	2.167	2.833	12.017	14.067	16.012	18.474	20.276
8	1.344	1.646	2.180	2.733	3.490	13.362	15.507	17.534	20.090	21.954
9	1.735	2.088	2.700	3.325	4.168	14.684	16.919	19.022	21.665	23.587
10	2.156	2.558	3.247	3.940	4.865	15.987	18.307	20.483	23.209	25.188
11	2.603	3.053	3.816	4.575	5.578	17.275	19.675	21.920	24.724	26.755
12	3.074	3.571	4.404	5.226	6.304	18.549	21.026	23.337	26.217	28.300
13	3.565	4.107	5.009	5.892	7.041	19.812	22.362	24.735	27.687	29.817
14	4.075	4.660	5.629	6.571	7.790	21.064	23.685	26.119	29.141	31.319
15	4.600	5.229	6.262	7.261	8.547	22.307	24.996	27.488	30.577	32.799
16	5.142	5.812	6.908	7.962	9.312	23.542	26.296	28.845	32.000	34.267
17	5.697	6.407	7.564	8.682	10.085	24.769	27.587	30.190	33.408	35.716
18	6.265	7.015	8.231	9.390	10.865	25.989	28.869	31.526	34.805	37.156
19	6.843	7.632	8.906	10.117	11.651	27.203	30.143	32.852	36.190	38.580
20	7.434	8.260	9.591	10.851	12.443	28.412	31.410	34.170	37.566	39.997
21	8.033	8.897	10.283	11.591	13.240	29.615	32.670	35.478	38.930	41.399
22	8.643	9.542	10.982	12.338	14.042	30.813	33.924	36.781	40.289	42.796
23	9.260	10.195	11.688	13.090	14.848	32.007	35.172	38.075	41.637	44.179
24	9.886	10.856	12.401	13.848	15.659	33.196	36.415	39.364	42.980	45.558
25	10.519	11.523	13.120	14.611	16.473	34.381	37.652	40.646	44.313	46.925
26	11.160	12.198	13.844	15.379	17.292	35.563	38.885	41.923	45.642	48.290
27	11.807	12.878	14.573	16.151	18.114	36.741	40.113	43.194	46.962	49.642
28	12.461	13.565	15.308	16.928	18.939	37.916	41.337	44.461	48.278	50.993
29	13.120	14.256	16.147	17.708	19.768	39.087	42.557	45.772	49.586	52.333
30	13.787	14.954	16.791	18.493	20.599	40.256	43.773	46.979	50.892	53.672
31	14.457	15.655	17.538	19.280	21.433	41.422	44.985	48.231	52.190	55.000
32	15.134	16.362	18.291	20.072	22.271	42.585	46.194	49.480	53.486	56.328
33	15.814	17.073	19.046	20.866	23.110	43.745	47.400	50.724	54.774	57.646
34	16.501	17.789	19.806	21.664	23.952	44.903	48.602	51.966	56.061	58.964
35	17.191	18.508	20.569	22.465	24.796	46.059	49.802	53.203	57.340	60.272
36	17.887	19.233	21.336	23.269	25.643	47.212	50.998	54.437	58.619	61.581
37	18.584	19.960	22.105	24.075	26.492	48.363	52.192	55.667	59.891	62.880
38	19.289	20.691	22.878	24.884	27.343	49.513	53.384	56.896	61.162	64.181
39	19.994	21.425	23.654	25.695	28.196	50.660	54.572	58.119	62.426	65.473
40	20.706	22.164	24.433	26.509	29.050	51.805	55.758	59.342	63.691	66.766


$$\text{For } \nu > 40, \chi_{\alpha,\nu}^2 \approx \nu \left( 1 - \frac{2}{9\nu} + z_\alpha \sqrt{\frac{2}{9\nu}} \right)^3$$

Copyright 2010 Cengage Learning. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part. Due to electronic rights, some third party content may be suppressed from the eBook and/or eChapter(s). Editorial review has deemed that any suppressed content does not materially affect the overall learning experience. Cengage Learning reserves the right to remove additional content at any time if subsequent rights restrictions require.

## A-12 Appendix Tables

Table A.8 t Curve Tail Areas


<i>t</i>	<i>v</i>	Area to the right of <i>t</i>																	
		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
0.0	.500	.500	.500	.500	.500	.500	.500	.500	.500	.500	.500	.500	.500	.500	.500	.500	.500	.500	.500
0.1	.468	.465	.463	.463	.462	.462	.462	.461	.461	.461	.461	.461	.461	.461	.461	.461	.461	.461	.461
0.2	.437	.430	.427	.426	.425	.424	.424	.423	.423	.423	.423	.423	.422	.422	.422	.422	.422	.422	.422
0.3	.407	.396	.392	.390	.388	.387	.386	.386	.386	.385	.385	.385	.384	.384	.384	.384	.384	.384	.384
0.4	.379	.364	.358	.355	.353	.352	.351	.350	.349	.349	.348	.348	.348	.347	.347	.347	.347	.347	.347
0.5	.352	.333	.326	.322	.319	.317	.316	.315	.315	.314	.313	.313	.313	.312	.312	.312	.312	.312	.312
0.6	.328	.305	.295	.290	.287	.285	.284	.283	.282	.281	.280	.280	.279	.279	.279	.278	.278	.278	.278
0.7	.306	.278	.267	.261	.258	.255	.253	.252	.251	.250	.249	.249	.248	.247	.247	.247	.247	.247	.246
0.8	.285	.254	.241	.234	.230	.227	.225	.223	.222	.221	.220	.220	.219	.218	.218	.218	.217	.217	.217
0.9	.267	.232	.217	.210	.205	.201	.199	.197	.196	.195	.194	.194	.193	.192	.191	.191	.190	.190	.190
1.0	.250	.211	.196	.187	.182	.178	.175	.173	.172	.170	.169	.169	.168	.167	.167	.166	.166	.165	.165
1.1	.235	.193	.176	.167	.162	.157	.154	.152	.150	.149	.147	.146	.146	.146	.144	.144	.143	.143	.143
1.2	.221	.177	.158	.148	.142	.138	.135	.132	.130	.129	.128	.127	.126	.124	.124	.124	.123	.123	.123
1.3	.209	.162	.142	.132	.125	.121	.117	.115	.113	.111	.110	.109	.108	.107	.107	.106	.105	.105	.105
1.4	.197	.148	.128	.117	.110	.106	.102	.100	.098	.096	.095	.093	.092	.091	.091	.090	.090	.089	.089
1.5	.187	.136	.115	.104	.097	.092	.089	.086	.084	.082	.081	.080	.079	.077	.077	.077	.076	.075	.075
1.6	.178	.125	.104	.092	.085	.080	.077	.074	.072	.070	.069	.068	.067	.065	.065	.065	.064	.064	.064
1.7	.169	.116	.094	.082	.075	.070	.065	.064	.062	.060	.059	.057	.056	.055	.055	.054	.054	.053	.053
1.8	.161	.107	.085	.073	.066	.061	.057	.055	.053	.051	.050	.049	.048	.046	.046	.045	.045	.044	.044
1.9	.154	.099	.077	.065	.058	.053	.050	.047	.045	.043	.042	.041	.040	.038	.038	.038	.037	.037	.037
2.0	.148	.092	.070	.058	.051	.046	.043	.040	.038	.037	.035	.034	.033	.032	.032	.031	.031	.030	.030
2.1	.141	.085	.063	.052	.045	.040	.037	.034	.033	.031	.030	.029	.028	.027	.027	.026	.025	.025	.025
2.2	.136	.079	.058	.046	.040	.035	.032	.029	.028	.026	.025	.024	.023	.022	.022	.021	.021	.021	.021
2.3	.131	.074	.052	.041	.035	.031	.027	.025	.023	.022	.021	.020	.019	.018	.018	.018	.017	.017	.017
2.4	.126	.069	.048	.037	.031	.027	.024	.022	.020	.019	.018	.017	.016	.015	.015	.014	.014	.014	.014
2.5	.121	.065	.044	.033	.027	.023	.020	.018	.017	.016	.015	.014	.013	.012	.012	.012	.011	.011	.011
2.6	.117	.061	.040	.030	.024	.020	.018	.016	.014	.013	.012	.012	.011	.010	.010	.010	.009	.009	.009
2.7	.113	.057	.037	.027	.021	.018	.015	.014	.012	.011	.010	.010	.009	.008	.008	.008	.008	.008	.007
2.8	.109	.054	.034	.024	.019	.016	.013	.012	.010	.009	.009	.008	.008	.007	.007	.006	.006	.006	.006
2.9	.106	.051	.031	.022	.017	.014	.011	.010	.009	.008	.007	.007	.006	.005	.005	.005	.005	.005	.005
3.0	.102	.048	.029	.020	.015	.012	.010	.009	.007	.007	.006	.006	.005	.004	.004	.004	.004	.004	.004
3.1	.099	.045	.027	.018	.013	.011	.009	.007	.006	.006	.005	.005	.004	.004	.004	.003	.003	.003	.003
3.2	.096	.043	.025	.016	.012	.009	.008	.006	.005	.005	.005	.004	.004	.003	.003	.003	.003	.002	.002
3.3	.094	.040	.023	.015	.011	.008	.007	.005	.005	.004	.004	.003	.003	.002	.002	.002	.002	.002	.002
3.4	.091	.038	.021	.014	.010	.007	.006	.005	.004	.003	.003	.002	.002	.002	.002	.002	.001	.001	.001
3.5	.089	.036	.020	.012	.009	.006	.005	.004	.003	.003	.002	.002	.002	.002	.002	.001	.001	.001	.001
3.6	.086	.035	.018	.011	.008	.006	.004	.004	.003	.002	.002	.002	.002	.001	.001	.001	.001	.001	.001
3.7	.084	.033	.017	.010	.007	.005	.004	.003	.002	.002	.002	.002	.001	.001	.001	.001	.001	.001	.001
3.8	.082	.031	.016	.010	.006	.004	.003	.003	.002	.002	.001	.001	.001	.001	.001	.001	.001	.001	.001
3.9	.080	.030	.015	.009	.006	.004	.003	.002	.002	.001	.001	.001	.001	.001	.001	.001	.000	.000	.000
4.0	.078	.029	.014	.008	.005	.004	.003	.002	.002	.001	.001	.001	.001	.001	.001	.001	.000	.000	.000



(continued)

Table A.8 t Curve Tail Areas (cont.)

$t$	$v$	19	20	21	22	23	24	25	26	27	28	29	30	35	40	60	120	$\infty (= z)$
0.0		.500	.500	.500	.500	.500	.500	.500	.500	.500	.500	.500	.500	.500	.500	.500	.500	.500
0.1		.461	.461	.461	.461	.461	.461	.461	.461	.461	.461	.461	.461	.460	.460	.460	.460	.460
0.2		.422	.422	.422	.422	.422	.422	.422	.421	.421	.421	.421	.421	.421	.421	.421	.421	.421
0.3		.384	.384	.384	.383	.383	.383	.383	.383	.383	.383	.383	.383	.383	.383	.383	.382	.382
0.4		.347	.347	.347	.347	.346	.346	.346	.346	.346	.346	.346	.346	.346	.346	.345	.345	.345
0.5		.311	.311	.311	.311	.311	.311	.311	.311	.311	.310	.310	.310	.310	.310	.309	.309	.309
0.6		.278	.278	.278	.277	.277	.277	.277	.277	.277	.277	.277	.276	.276	.275	.275	.274	
0.7		.246	.246	.246	.246	.245	.245	.245	.245	.245	.245	.245	.244	.244	.243	.243	.242	
0.8		.217	.217	.216	.216	.216	.216	.216	.215	.215	.215	.215	.215	.214	.214	.213	.213	
0.9		.190	.189	.189	.189	.189	.189	.188	.188	.188	.188	.188	.187	.187	.187	.186	.185	
1.0		.165	.165	.164	.164	.164	.164	.163	.163	.163	.163	.163	.162	.162	.162	.161	.160	
1.1		.143	.142	.142	.142	.141	.141	.141	.141	.141	.140	.140	.140	.139	.139	.138	.137	
1.2		.122	.122	.122	.121	.121	.121	.121	.120	.120	.120	.120	.120	.119	.119	.117	.116	
1.3		.105	.104	.104	.104	.103	.103	.103	.103	.102	.102	.102	.102	.101	.101	.099	.098	
1.4		.089	.089	.088	.088	.087	.087	.087	.087	.086	.086	.086	.086	.085	.085	.083	.082	
1.5		.075	.075	.074	.074	.074	.073	.073	.073	.073	.072	.072	.072	.071	.071	.069	.068	
1.6		.063	.063	.062	.062	.062	.061	.061	.061	.060	.060	.060	.059	.059	.059	.057	.056	
1.7		.053	.052	.052	.052	.051	.051	.051	.051	.050	.050	.050	.049	.049	.048	.047	.046	
1.8		.044	.043	.043	.043	.042	.042	.042	.042	.042	.041	.041	.041	.040	.040	.038	.037	
1.9		.036	.036	.036	.035	.035	.035	.035	.034	.034	.034	.034	.034	.033	.032	.031	.030	
2.0		.030	.030	.029	.029	.029	.028	.028	.028	.028	.028	.027	.027	.027	.026	.025	.024	
2.1		.025	.024	.024	.024	.023	.023	.023	.023	.022	.022	.022	.022	.021	.020	.019	.018	
2.2		.020	.020	.020	.019	.019	.019	.019	.018	.018	.018	.018	.017	.017	.016	.015	.014	
2.3		.016	.016	.016	.016	.015	.015	.015	.015	.015	.014	.014	.014	.013	.012	.012	.011	
2.4		.013	.013	.013	.013	.012	.012	.012	.012	.012	.012	.011	.011	.011	.010	.009	.008	
2.5		.011	.011	.010	.010	.010	.010	.010	.009	.009	.009	.009	.009	.008	.008	.007	.006	
2.6		.009	.009	.008	.008	.008	.008	.008	.008	.007	.007	.007	.007	.007	.007	.006	.005	
2.7		.007	.007	.007	.007	.006	.006	.006	.006	.006	.006	.006	.005	.005	.005	.004	.003	
2.8		.006	.006	.005	.005	.005	.005	.005	.005	.005	.005	.004	.004	.004	.004	.003	.003	
2.9		.005	.004	.004	.004	.004	.004	.004	.004	.004	.004	.004	.003	.003	.003	.002	.002	
3.0		.004	.004	.003	.003	.003	.003	.003	.003	.003	.003	.003	.003	.002	.002	.002	.002	
3.1		.003	.003	.003	.003	.003	.002	.002	.002	.002	.002	.002	.002	.002	.002	.001	.001	
3.2		.002	.002	.002	.002	.002	.002	.002	.002	.002	.002	.002	.002	.001	.001	.001	.001	
3.3		.002	.002	.002	.002	.002	.001	.001	.001	.001	.001	.001	.001	.001	.001	.001	.000	
3.4		.002	.001	.001	.001	.001	.001	.001	.001	.001	.001	.001	.001	.001	.001	.001	.000	
3.5		.001	.001	.001	.001	.001	.001	.001	.001	.001	.001	.001	.001	.001	.001	.000	.000	
3.6		.001	.001	.001	.001	.001	.001	.001	.001	.001	.001	.001	.001	.000	.000	.000	.000	
3.7		.001	.001	.001	.001	.001	.001	.001	.000	.000	.000	.000	.000	.000	.000	.000	.000	
3.8		.001	.001	.001	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	
3.9		.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	
4.0		.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	



**A-14** Appendix Tables**Table A.9 Critical Values for F Distributions**

		$\nu_1 = \text{numerator df}$								
		1	2	3	4	5	6	7	8	9
$\nu_2 = \text{denominator df}$	.100	39.86	49.50	53.59	55.83	57.24	58.20	58.91	59.44	59.86
	.050	161.45	199.50	215.71	224.58	230.16	233.99	236.77	238.88	240.54
	.010	4052.20	4999.50	5403.40	5624.60	5763.60	5859.00	5928.40	5981.10	6022.50
	.001	405,284	500,000	540,379	562,500	576,405	585,937	592,873	598,144	602,284
	.100	8.53	9.00	9.16	9.24	9.29	9.33	9.35	9.37	9.38
	.050	18.51	19.00	19.16	19.25	19.30	19.33	19.35	19.37	19.38
	.010	98.50	99.00	99.17	99.25	99.30	99.33	99.36	99.37	99.39
	.001	998.50	999.00	999.17	999.25	999.30	999.33	999.36	999.37	999.39
	.100	5.54	5.46	5.39	5.34	5.31	5.28	5.27	5.25	5.24
	.050	10.13	9.55	9.28	9.12	9.01	8.94	8.89	8.85	8.81
	.010	34.12	30.82	29.46	28.71	28.24	27.91	27.67	27.49	27.35
	.001	167.03	148.50	141.11	137.10	134.58	132.85	131.58	130.62	129.86
$\nu_2 = \text{denominator df}$	.100	4.54	4.32	4.19	4.11	4.05	4.01	3.98	3.95	3.94
	.050	7.71	6.94	6.59	6.39	6.26	6.16	6.09	6.04	6.00
	.010	21.20	18.00	16.69	15.98	15.52	15.21	14.98	14.80	14.66
	.001	74.14	61.25	56.18	53.44	51.71	50.53	49.66	49.00	48.47
	.100	4.06	3.78	3.62	3.52	3.45	3.40	3.37	3.34	3.32
	.050	6.61	5.79	5.41	5.19	5.05	4.95	4.88	4.82	4.77
	.010	16.26	13.27	12.06	11.39	10.97	10.67	10.46	10.29	10.16
	.001	47.18	37.12	33.20	31.09	29.75	28.83	28.16	27.65	27.24
	.100	3.78	3.46	3.29	3.18	3.11	3.05	3.01	2.98	2.96
	.050	5.99	5.14	4.76	4.53	4.39	4.28	4.21	4.15	4.10
	.010	13.75	10.92	9.78	9.15	8.75	8.47	8.26	8.10	7.98
	.001	35.51	27.00	23.70	21.92	20.80	20.03	19.46	19.03	18.69
$\nu_2 = \text{denominator df}$	.100	3.59	3.26	3.07	2.96	2.88	2.83	2.78	2.75	2.72
	.050	5.59	4.74	4.35	4.12	3.97	3.87	3.79	3.73	3.68
	.010	12.25	9.55	8.45	7.85	7.46	7.19	6.99	6.84	6.72
	.001	29.25	21.69	18.77	17.20	16.21	15.52	15.02	14.63	14.33
	.100	3.46	3.11	2.92	2.81	2.73	2.67	2.62	2.59	2.56
	.050	5.32	4.46	4.07	3.84	3.69	3.58	3.50	3.44	3.39
	.010	11.26	8.65	7.59	7.01	6.63	6.37	6.18	6.03	5.91
	.001	25.41	18.49	15.83	14.39	13.48	12.86	12.40	12.05	11.77
	.100	3.36	3.01	2.81	2.69	2.61	2.55	2.51	2.47	2.44
	.050	5.12	4.26	3.86	3.63	3.48	3.37	3.29	3.23	3.18
	.010	10.56	8.02	6.99	6.42	6.06	5.80	5.61	5.47	5.35
	.001	22.86	16.39	13.90	12.56	11.71	11.13	10.70	10.37	10.11
$\nu_2 = \text{denominator df}$	.100	3.29	2.92	2.73	2.61	2.52	2.46	2.41	2.38	2.35
	.050	4.96	4.10	3.71	3.48	3.33	3.22	3.14	3.07	3.02
	.010	10.04	7.56	6.55	5.99	5.64	5.39	5.20	5.06	4.94
	.001	21.04	14.91	12.55	11.28	10.48	9.93	9.52	9.20	8.96
	.100	3.23	2.86	2.66	2.54	2.45	2.39	2.34	2.30	2.27
	.050	4.84	3.98	3.59	3.36	3.20	3.09	3.01	2.95	2.90
	.010	9.65	7.21	6.22	5.67	5.32	5.07	4.89	4.74	4.63
	.001	19.69	13.81	11.56	10.35	9.58	9.05	8.66	8.35	8.12
	.100	3.18	2.81	2.61	2.48	2.39	2.33	2.28	2.24	2.21
	.050	4.75	3.89	3.49	3.26	3.11	3.00	2.91	2.85	2.80
	.010	9.33	6.93	5.95	5.41	5.06	4.82	4.64	4.50	4.39
	.001	18.64	12.97	10.80	9.63	8.89	8.38	8.00	7.71	7.48

(continued)

Table A.9 Critical Values for F Distributions (cont.)

$\nu_1 = \text{numerator df}$										
10	12	15	20	25	30	40	50	60	120	1000
60.19	60.71	61.22	61.74	62.05	62.26	62.53	62.69	62.79	63.06	63.30
241.88	243.91	245.95	248.01	249.26	250.10	251.14	251.77	252.20	253.25	254.19
6055.80	6106.30	6157.30	6208.70	6239.80	6260.60	6286.80	6302.50	6313.00	6339.40	6362.70
605,621	610,668	615,764	620,908	624,017	626,099	628,712	630,285	631,337	633,972	636,301
9.39	9.41	9.42	9.44	9.45	9.46	9.47	9.47	9.47	9.48	9.49
19.40	19.41	19.43	19.45	19.46	19.46	19.47	19.48	19.48	19.49	19.49
99.40	99.42	99.43	99.45	99.46	99.47	99.47	99.48	99.48	99.49	99.50
999.40	999.42	999.43	999.45	999.46	999.47	999.47	999.48	999.48	999.49	999.50
5.23	5.22	5.20	5.18	5.17	5.17	5.16	5.15	5.15	5.14	5.13
8.79	8.74	8.70	8.66	8.63	8.62	8.59	8.58	8.57	8.55	8.53
27.23	27.05	26.87	26.69	26.58	26.50	26.41	26.35	26.32	26.22	26.14
129.25	128.32	127.37	126.42	125.84	125.45	124.96	124.66	124.47	123.97	123.53
3.92	3.90	3.87	3.84	3.83	3.82	3.80	3.80	3.79	3.78	3.76
5.96	5.91	5.86	5.80	5.77	5.75	5.72	5.70	5.69	5.66	5.63
14.55	14.37	14.20	14.02	13.91	13.84	13.75	13.69	13.65	13.56	13.47
48.05	47.41	46.76	46.10	45.70	45.43	45.09	44.88	44.75	44.40	44.09
3.30	3.27	3.24	3.21	3.19	3.17	3.16	3.15	3.14	3.12	3.11
4.74	4.68	4.62	4.56	4.52	4.50	4.46	4.44	4.43	4.40	4.37
10.05	9.89	9.72	9.55	9.45	9.38	9.29	9.24	9.20	9.11	9.03
26.92	26.42	25.91	25.39	25.08	24.87	24.60	24.44	24.33	24.06	23.82
2.94	2.90	2.87	2.84	2.81	2.80	2.78	2.77	2.76	2.74	2.72
4.06	4.00	3.94	3.87	3.83	3.81	3.77	3.75	3.74	3.70	3.67
7.87	7.72	7.56	7.40	7.30	7.23	7.14	7.09	7.06	6.97	6.89
18.41	17.99	17.56	17.12	16.85	16.67	16.44	16.31	16.21	15.98	15.77
2.70	2.67	2.63	2.59	2.57	2.56	2.54	2.52	2.51	2.49	2.47
3.64	3.57	3.51	3.44	3.40	3.38	3.34	3.32	3.30	3.27	3.23
6.62	6.47	6.31	6.16	6.06	5.99	5.91	5.86	5.82	5.74	5.66
14.08	13.71	13.32	12.93	12.69	12.53	12.33	12.20	12.12	11.91	11.72
2.54	2.50	2.46	2.42	2.40	2.38	2.36	2.35	2.34	2.32	2.30
3.35	3.28	3.22	3.15	3.11	3.08	3.04	3.02	3.01	2.97	2.93
5.81	5.67	5.52	5.36	5.26	5.20	5.12	5.07	5.03	4.95	4.87
11.54	11.19	10.84	10.48	10.26	10.11	9.92	9.80	9.73	9.53	9.36
2.42	2.38	2.34	2.30	2.27	2.25	2.23	2.22	2.21	2.18	2.16
3.14	3.07	3.01	2.94	2.89	2.86	2.83	2.80	2.79	2.75	2.71
5.26	5.11	4.96	4.81	4.71	4.65	4.57	4.52	4.48	4.40	4.32
9.89	9.57	9.24	8.90	8.69	8.55	8.37	8.26	8.19	8.00	7.84
2.32	2.28	2.24	2.20	2.17	2.16	2.13	2.12	2.11	2.08	2.06
2.98	2.91	2.85	2.77	2.73	2.70	2.66	2.64	2.62	2.58	2.54
4.85	4.71	4.56	4.41	4.31	4.25	4.17	4.12	4.08	4.00	3.92
8.75	8.45	8.13	7.80	7.60	7.47	7.30	7.19	7.12	6.94	6.78
2.25	2.21	2.17	2.12	2.10	2.08	2.05	2.04	2.03	2.00	1.98
2.85	2.79	2.72	2.65	2.60	2.57	2.53	2.51	2.49	2.45	2.41
4.54	4.40	4.25	4.10	4.01	3.94	3.86	3.81	3.78	3.69	3.61
7.92	7.63	7.32	7.01	6.81	6.68	6.52	6.42	6.35	6.18	6.02
2.19	2.15	2.10	2.06	2.03	2.01	1.99	1.97	1.96	1.93	1.91
2.75	2.69	2.62	2.54	2.50	2.47	2.43	2.40	2.38	2.34	2.30
4.30	4.16	4.01	3.86	3.76	3.70	3.62	3.57	3.54	3.45	3.37
7.29	7.00	6.71	6.40	6.22	6.09	5.93	5.83	5.76	5.59	5.44

(continued)

Copyright 2010 Cengage Learning. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part. Due to electronic rights, some third party content may be suppressed from the eBook and/or eChapter(s). Editorial review has deemed that any suppressed content does not materially affect the overall learning experience. Cengage Learning reserves the right to remove additional content at any time if subsequent rights restrictions require.

## A-16 Appendix Tables

Table A.9 Critical Values for F Distributions (cont.)

		$\nu_1 = \text{numerator df}$									
		1	2	3	4	5	6	7	8	9	
$\alpha$		.100	3.14	2.76	2.56	2.43	2.35	2.28	2.23	2.20	2.16
13	.050	4.67	3.81	3.41	3.18	3.03	2.92	2.83	2.77	2.71	
	.010	9.07	6.70	5.74	5.21	4.86	4.62	4.44	4.30	4.19	
	.001	17.82	12.31	10.21	9.07	8.35	7.86	7.49	7.21	6.98	
	.100	3.10	2.73	2.52	2.39	2.31	2.24	2.19	2.15	2.12	
14	.050	4.60	3.74	3.34	3.11	2.96	2.85	2.76	2.70	2.65	
	.010	8.86	6.51	5.56	5.04	4.69	4.46	4.28	4.14	4.03	
	.001	17.14	11.78	9.73	8.62	7.92	7.44	7.08	6.80	6.58	
	.100	3.07	2.70	2.49	2.36	2.27	2.21	2.16	2.12	2.09	
15	.050	4.54	3.68	3.29	3.06	2.90	2.79	2.71	2.64	2.59	
	.010	8.68	6.36	5.42	4.89	4.56	4.32	4.14	4.00	3.89	
	.001	16.59	11.34	9.34	8.25	7.57	7.09	6.74	6.47	6.26	
	.100	3.05	2.67	2.46	2.33	2.24	2.18	2.13	2.09	2.06	
16	.050	4.49	3.63	3.24	3.01	2.85	2.74	2.66	2.59	2.54	
	.010	8.53	6.23	5.29	4.77	4.44	4.20	4.03	3.89	3.78	
	.001	16.12	10.97	9.01	7.94	7.27	6.80	6.46	6.19	5.98	
	.100	3.03	2.64	2.44	2.31	2.22	2.15	2.10	2.06	2.03	
17	.050	4.45	3.59	3.20	2.96	2.81	2.70	2.61	2.55	2.49	
	.010	8.40	6.11	5.19	4.67	4.34	4.10	3.93	3.79	3.68	
	.001	15.72	10.66	8.73	7.68	7.02	6.56	6.22	5.96	5.75	
	.100	3.01	2.62	2.42	2.29	2.20	2.13	2.08	2.04	2.00	
18	.050	4.41	3.55	3.16	2.93	2.77	2.66	2.58	2.51	2.46	
	.010	8.29	6.01	5.09	4.58	4.25	4.01	3.84	3.71	3.60	
	.001	15.38	10.39	8.49	7.46	6.81	6.35	6.02	5.76	5.56	
	.100	2.99	2.61	2.40	2.27	2.18	2.11	2.06	2.02	1.98	
19	.050	4.38	3.52	3.13	2.90	2.74	2.63	2.54	2.48	2.42	
	.010	8.18	5.93	5.01	4.50	4.17	3.94	3.77	3.63	3.52	
	.001	15.08	10.16	8.28	7.27	6.62	6.18	5.85	5.59	5.39	
	.100	2.97	2.59	2.38	2.25	2.16	2.09	2.04	2.00	1.96	
20	.050	4.35	3.49	3.10	2.87	2.71	2.60	2.51	2.45	2.39	
	.010	8.10	5.85	4.94	4.43	4.10	3.87	3.70	3.56	3.46	
	.001	14.82	9.95	8.10	7.10	6.46	6.02	5.69	5.44	5.24	
	.100	2.96	2.57	2.36	2.23	2.14	2.08	2.02	1.98	1.95	
21	.050	4.32	3.47	3.07	2.84	2.68	2.57	2.49	2.42	2.37	
	.010	8.02	5.78	4.87	4.37	4.04	3.81	3.64	3.51	3.40	
	.001	14.59	9.77	7.94	6.95	6.32	5.88	5.56	5.31	5.11	
	.100	2.95	2.56	2.35	2.22	2.13	2.06	2.01	1.97	1.93	
22	.050	4.30	3.44	3.05	2.82	2.66	2.55	2.46	2.40	2.34	
	.010	7.95	5.72	4.82	4.31	3.99	3.76	3.59	3.45	3.35	
	.001	14.38	9.61	7.80	6.81	6.19	5.76	5.44	5.19	4.99	
	.100	2.94	2.55	2.34	2.21	2.11	2.05	1.99	1.95	1.92	
23	.050	4.28	3.42	3.03	2.80	2.64	2.53	2.44	2.37	2.32	
	.010	7.88	5.66	4.76	4.26	3.94	3.71	3.54	3.41	3.30	
	.001	14.20	9.47	7.67	6.70	6.08	5.65	5.33	5.09	4.89	
	.100	2.93	2.54	2.33	2.19	2.10	2.04	1.98	1.94	1.91	
24	.050	4.26	3.40	3.01	2.78	2.62	2.51	2.42	2.36	2.30	
	.010	7.82	5.61	4.72	4.22	3.90	3.67	3.50	3.36	3.26	
	.001	14.03	9.34	7.55	6.59	5.98	5.55	5.23	4.99	4.80	

(continued)

**Table A.9 Critical Values for F Distributions (cont.)**

$v_1 = \text{numerator df}$										
10	12	15	20	25	30	40	50	60	120	1000
2.14	2.10	2.05	2.01	1.98	1.96	1.93	1.92	1.90	1.88	1.85
2.67	2.60	2.53	2.46	2.41	2.38	2.34	2.31	2.30	2.25	2.21
4.10	3.96	3.82	3.66	3.57	3.51	3.43	3.38	3.34	3.25	3.18
6.80	6.52	6.23	5.93	5.75	5.63	5.47	5.37	5.30	5.14	4.99
2.10	2.05	2.01	1.96	1.93	1.91	1.89	1.87	1.86	1.83	1.80
2.60	2.53	2.46	2.39	2.34	2.31	2.27	2.24	2.22	2.18	2.14
3.94	3.80	3.66	3.51	3.41	3.35	3.27	3.22	3.18	3.09	3.02
6.40	6.13	5.85	5.56	5.38	5.25	5.10	5.00	4.94	4.77	4.62
2.06	2.02	1.97	1.92	1.89	1.87	1.85	1.83	1.82	1.79	1.76
2.54	2.48	2.40	2.33	2.28	2.25	2.20	2.18	2.16	2.11	2.07
3.80	3.67	3.52	3.37	3.28	3.21	3.13	3.08	3.05	2.96	2.88
6.08	5.81	5.54	5.25	5.07	4.95	4.80	4.70	4.64	4.47	4.33
2.03	1.99	1.94	1.89	1.86	1.84	1.81	1.79	1.78	1.75	1.72
2.49	2.42	2.35	2.28	2.23	2.19	2.15	2.12	2.11	2.06	2.02
3.69	3.55	3.41	3.26	3.16	3.10	3.02	2.97	2.93	2.84	2.76
5.81	5.55	5.27	4.99	4.82	4.70	4.54	4.45	4.39	4.23	4.08
2.00	1.96	1.91	1.86	1.83	1.81	1.78	1.76	1.75	1.72	1.69
2.45	2.38	2.31	2.23	2.18	2.15	2.10	2.08	2.06	2.01	1.97
3.59	3.46	3.31	3.16	3.07	3.00	2.92	2.87	2.83	2.75	2.66
5.58	5.32	5.05	4.78	4.60	4.48	4.33	4.24	4.18	4.02	3.87
1.98	1.93	1.89	1.84	1.80	1.78	1.75	1.74	1.72	1.69	1.66
2.41	2.34	2.27	2.19	2.14	2.11	2.06	2.04	2.02	1.97	1.92
3.51	3.37	3.23	3.08	2.98	2.92	2.84	2.78	2.75	2.66	2.58
5.39	5.13	4.87	4.59	4.42	4.30	4.15	4.06	4.00	3.84	3.69
1.96	1.91	1.86	1.81	1.78	1.76	1.73	1.71	1.70	1.67	1.64
2.38	2.31	2.23	2.16	2.11	2.07	2.03	2.00	1.98	1.93	1.88
3.43	3.30	3.15	3.00	2.91	2.84	2.76	2.71	2.67	2.58	2.50
5.22	4.97	4.70	4.43	4.26	4.14	3.99	3.90	3.84	3.68	3.53
1.94	1.89	1.84	1.79	1.76	1.74	1.71	1.69	1.68	1.64	1.61
2.35	2.28	2.20	2.12	2.07	2.04	1.99	1.97	1.95	1.90	1.85
3.37	3.23	3.09	2.94	2.84	2.78	2.69	2.64	2.61	2.52	2.43
5.08	4.82	4.56	4.29	4.12	4.00	3.86	3.77	3.70	3.54	3.40
1.92	1.87	1.83	1.78	1.74	1.72	1.69	1.67	1.66	1.62	1.59
2.32	2.25	2.18	2.10	2.05	2.01	1.96	1.94	1.92	1.87	1.82
3.31	3.17	3.03	2.88	2.79	2.72	2.64	2.58	2.55	2.46	2.37
4.95	4.70	4.44	4.17	4.00	3.88	3.74	3.64	3.58	3.42	3.28
1.90	1.86	1.81	1.76	1.73	1.70	1.67	1.65	1.64	1.60	1.57
2.30	2.23	2.15	2.07	2.02	1.98	1.94	1.91	1.89	1.84	1.79
3.26	3.12	2.98	2.83	2.73	2.67	2.58	2.53	2.50	2.40	2.32
4.83	4.58	4.33	4.06	3.89	3.78	3.63	3.54	3.48	3.32	3.17
1.89	1.84	1.80	1.74	1.71	1.69	1.66	1.64	1.62	1.59	1.55
2.27	2.20	2.13	2.05	2.00	1.96	1.91	1.88	1.86	1.81	1.76
3.21	3.07	2.93	2.78	2.69	2.62	2.54	2.48	2.45	2.35	2.27
4.73	4.48	4.23	3.96	3.79	3.68	3.53	3.44	3.38	3.22	3.08
1.88	1.83	1.78	1.73	1.70	1.67	1.64	1.62	1.61	1.57	1.54
2.25	2.18	2.11	2.03	1.97	1.94	1.89	1.86	1.84	1.79	1.74
3.17	3.03	2.89	2.74	2.64	2.58	2.49	2.44	2.40	2.31	2.22
4.64	4.39	4.14	3.87	3.71	3.59	3.45	3.36	3.29	3.14	2.99

(continued)

Copyright 2010 Cengage Learning. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part. Due to electronic rights, some third party content may be suppressed from the eBook and/or eChapter(s). Editorial review has deemed that any suppressed content does not materially affect the overall learning experience. Cengage Learning reserves the right to remove additional content at any time if subsequent rights restrictions require.

A-18 Appendix Tables

Table A.9 Critical Values for *F* Distributions (cont.)

		$\nu_1 = \text{numerator df}$									
		1	2	3	4	5	6	7	8	9	
		.100	2.92	2.53	2.32	2.18	2.09	2.02	1.97	1.93	1.89
25	.050	4.24	3.39	2.99	2.76	2.60	2.49	2.40	2.34	2.28	
	.010	7.77	5.57	4.68	4.18	3.85	3.63	3.46	3.32	3.22	
	.001	13.88	9.22	7.45	6.49	5.89	5.46	5.15	4.91	4.71	
	.100	2.91	2.52	2.31	2.17	2.08	2.01	1.96	1.92	1.88	
26	.050	4.23	3.37	2.98	2.74	2.59	2.47	2.39	2.32	2.27	
	.010	7.72	5.53	4.64	4.14	3.82	3.59	3.42	3.29	3.18	
	.001	13.74	9.12	7.36	6.41	5.80	5.38	5.07	4.83	4.64	
	.100	2.90	2.51	2.30	2.17	2.07	2.00	1.95	1.91	1.87	
27	.050	4.21	3.35	2.96	2.73	2.57	2.46	2.37	2.31	2.25	
	.010	7.68	5.49	4.60	4.11	3.78	3.56	3.39	3.26	3.15	
	.001	13.61	9.02	7.27	6.33	5.73	5.31	5.00	4.76	4.57	
	.100	2.89	2.50	2.29	2.16	2.06	2.00	1.94	1.90	1.87	
28	.050	4.20	3.34	2.95	2.71	2.56	2.45	2.36	2.29	2.24	
	.010	7.64	5.45	4.57	4.07	3.75	3.53	3.36	3.23	3.12	
	.001	13.50	8.93	7.19	6.25	5.66	5.24	4.93	4.69	4.50	
	.100	2.89	2.50	2.28	2.15	2.06	1.99	1.93	1.89	1.86	
29	.050	4.18	3.33	2.93	2.70	2.55	2.43	2.35	2.28	2.22	
	.010	7.60	5.42	4.54	4.04	3.73	3.50	3.33	3.20	3.09	
	.001	13.39	8.85	7.12	6.19	5.59	5.18	4.87	4.64	4.45	
	.100	2.88	2.49	2.28	2.14	2.05	1.98	1.93	1.88	1.85	
30	.050	4.17	3.32	2.92	2.69	2.53	2.42	2.33	2.27	2.21	
	.010	7.56	5.39	4.51	4.02	3.70	3.47	3.30	3.17	3.07	
	.001	13.29	8.77	7.05	6.12	5.53	5.12	4.82	4.58	4.39	
	.100	2.88	2.44	2.23	2.09	2.00	1.93	1.87	1.83	1.79	
40	.050	4.08	3.23	2.84	2.61	2.45	2.34	2.25	2.18	2.12	
	.010	7.31	5.18	4.31	3.83	3.51	3.29	3.12	2.99	2.89	
	.001	12.61	8.25	6.59	5.70	5.13	4.73	4.44	4.21	4.02	
	.100	2.81	2.41	2.20	2.06	1.97	1.90	1.84	1.80	1.76	
50	.050	4.03	3.18	2.79	2.56	2.40	2.29	2.20	2.13	2.07	
	.010	7.17	5.06	4.20	3.72	3.41	3.19	3.02	2.89	2.78	
	.001	12.22	7.96	6.34	5.46	4.90	4.51	4.22	4.00	3.82	
	.100	2.79	2.39	2.18	2.04	1.95	1.87	1.82	1.77	1.74	
60	.050	4.00	3.15	2.76	2.53	2.37	2.25	2.17	2.10	2.04	
	.010	7.08	4.98	4.13	3.65	3.34	3.12	2.95	2.82	2.72	
	.001	11.97	7.77	6.17	5.31	4.76	4.37	4.09	3.86	3.69	
	.100	2.76	2.36	2.14	2.00	1.91	1.83	1.78	1.73	1.69	
100	.050	3.94	3.09	2.70	2.46	2.31	2.19	2.10	2.03	1.97	
	.010	6.90	4.82	3.98	3.51	3.21	2.99	2.82	2.69	2.59	
	.001	11.50	7.41	5.86	5.02	4.48	4.11	3.83	3.61	3.44	
	.100	2.73	2.33	2.11	1.97	1.88	1.80	1.75	1.70	1.66	
200	.050	3.89	3.04	2.65	2.42	2.26	2.14	2.06	1.98	1.93	
	.010	6.76	4.71	3.88	3.41	3.11	2.89	2.73	2.60	2.50	
	.001	11.15	7.15	5.63	4.81	4.29	3.92	3.65	3.43	3.26	
	.100	2.71	2.31	2.09	1.95	1.85	1.78	1.72	1.68	1.64	
1000	.050	3.85	3.00	2.61	2.38	2.22	2.11	2.02	1.95	1.89	
	.010	6.66	4.63	3.80	3.34	3.04	2.82	2.66	2.53	2.43	
	.001	10.89	6.96	5.46	4.65	4.14	3.78	3.51	3.30	3.13	

(continued)

Table A.9 Critical Values for F Distributions (cont.)

$v_1 = \text{numerator df}$										
10	12	15	20	25	30	40	50	60	120	1000
1.87	1.82	1.77	1.72	1.68	1.66	1.63	1.61	1.59	1.56	1.52
2.24	2.16	2.09	2.01	1.96	1.92	1.87	1.84	1.82	1.77	1.72
3.13	2.99	2.85	2.70	2.60	2.54	2.45	2.40	2.36	2.27	2.18
4.56	4.31	4.06	3.79	3.63	3.52	3.37	3.28	3.22	3.06	2.91
1.86	1.81	1.76	1.71	1.67	1.65	1.61	1.59	1.58	1.54	1.51
2.22	2.15	2.07	1.99	1.94	1.90	1.85	1.82	1.80	1.75	1.70
3.09	2.96	2.81	2.66	2.57	2.50	2.42	2.36	2.33	2.23	2.14
4.48	4.24	3.99	3.72	3.56	3.44	3.30	3.21	3.15	2.99	2.84
1.85	1.80	1.75	1.70	1.66	1.64	1.60	1.58	1.57	1.53	1.50
2.20	2.13	2.06	1.97	1.92	1.88	1.84	1.81	1.79	1.73	1.68
3.06	2.93	2.78	2.63	2.54	2.47	2.38	2.33	2.29	2.20	2.11
4.41	4.17	3.92	3.66	3.49	3.38	3.23	3.14	3.08	2.92	2.78
1.84	1.79	1.74	1.69	1.65	1.63	1.59	1.57	1.56	1.52	1.48
2.19	2.12	2.04	1.96	1.91	1.87	1.82	1.79	1.77	1.71	1.66
3.03	2.90	2.75	2.60	2.51	2.44	2.35	2.30	2.26	2.17	2.08
4.35	4.11	3.86	3.60	3.43	3.32	3.18	3.09	3.02	2.86	2.72
1.83	1.78	1.73	1.68	1.64	1.62	1.58	1.56	1.55	1.51	1.47
2.18	2.10	2.03	1.94	1.89	1.85	1.81	1.77	1.75	1.70	1.65
3.00	2.87	2.73	2.57	2.48	2.41	2.33	2.27	2.23	2.14	2.05
4.29	4.05	3.80	3.54	3.38	3.27	3.12	3.03	2.97	2.81	2.66
1.82	1.77	1.72	1.67	1.63	1.61	1.57	1.55	1.54	1.50	1.46
2.16	2.09	2.01	1.93	1.88	1.84	1.79	1.76	1.74	1.68	1.63
2.98	2.84	2.70	2.55	2.45	2.39	2.30	2.25	2.21	2.11	2.02
4.24	4.00	3.75	3.49	3.33	3.22	3.07	2.98	2.92	2.76	2.61
1.76	1.71	1.66	1.61	1.57	1.54	1.51	1.48	1.47	1.42	1.38
2.08	2.00	1.92	1.84	1.78	1.74	1.69	1.66	1.64	1.58	1.52
2.80	2.66	2.52	2.37	2.27	2.20	2.11	2.06	2.02	1.92	1.82
3.87	3.64	3.40	3.14	2.98	2.87	2.73	2.64	2.57	2.41	2.25
1.73	1.68	1.63	1.57	1.53	1.50	1.46	1.44	1.42	1.38	1.33
2.03	1.95	1.87	1.78	1.73	1.69	1.63	1.60	1.58	1.51	1.45
2.70	2.56	2.42	2.27	2.17	2.10	2.01	1.95	1.91	1.80	1.70
3.67	3.44	3.20	2.95	2.79	2.68	2.53	2.44	2.38	2.21	2.05
1.71	1.66	1.60	1.54	1.50	1.48	1.44	1.41	1.40	1.35	1.30
1.99	1.92	1.84	1.75	1.69	1.65	1.59	1.56	1.53	1.47	1.40
2.63	2.50	2.35	2.20	2.10	2.03	1.94	1.88	1.84	1.73	1.62
3.54	3.32	3.08	2.83	2.67	2.55	2.41	2.32	2.25	2.08	1.92
1.66	1.61	1.56	1.49	1.45	1.42	1.38	1.35	1.34	1.28	1.22
1.93	1.85	1.77	1.68	1.62	1.57	1.52	1.48	1.45	1.38	1.30
2.50	2.37	2.22	2.07	1.97	1.89	1.80	1.74	1.69	1.57	1.45
3.30	3.07	2.84	2.59	2.43	2.32	2.17	2.08	2.01	1.83	1.64
1.63	1.58	1.52	1.46	1.41	1.38	1.34	1.31	1.29	1.23	1.16
1.88	1.80	1.72	1.62	1.56	1.52	1.46	1.41	1.39	1.30	1.21
2.41	2.27	2.13	1.97	1.87	1.79	1.69	1.63	1.58	1.45	1.30
3.12	2.90	2.67	2.42	2.26	2.15	2.00	1.90	1.83	1.64	1.43
1.61	1.55	1.49	1.43	1.38	1.35	1.30	1.27	1.25	1.18	1.08
1.84	1.76	1.68	1.58	1.52	1.47	1.41	1.36	1.33	1.24	1.11
2.34	2.20	2.06	1.90	1.79	1.72	1.61	1.54	1.50	1.35	1.16
2.99	2.77	2.54	2.30	2.14	2.02	1.87	1.77	1.69	1.49	1.22

Copyright 2010 Cengage Learning. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part. Due to electronic rights, some third party content may be suppressed from the eBook and/or eChapter(s). Editorial review has determined that any suppressed content does not materially affect the overall learning experience. Cengage Learning reserves the right to remove additional content at any time if subsequent rights restrictions require.

## A-20 Appendix Tables

Table A.10 Critical Values for Studentized Range Distributions

<i>v</i>	$\alpha$	<i>m</i>											
		2	3	4	5	6	7	8	9	10	11	12	
5	.05	3.64	4.60	5.22	5.67	6.03	6.33	6.58	6.80	6.99	7.17	7.32	
	.01	5.70	6.98	7.80	8.42	8.91	9.32	9.67	9.97	10.24	10.48	10.70	
6	.05	3.46	4.34	4.90	5.30	5.63	5.90	6.12	6.32	6.49	6.65	6.79	
	.01	5.24	6.33	7.03	7.56	7.97	8.32	8.61	8.87	9.10	9.30	9.48	
7	.05	3.34	4.16	4.68	5.06	5.36	5.61	5.82	6.00	6.16	6.30	6.43	
	.01	4.95	5.92	6.54	7.01	7.37	7.68	7.94	8.17	8.37	8.55	8.71	
8	.05	3.26	4.04	4.53	4.89	5.17	5.40	5.60	5.77	5.92	6.05	6.18	
	.01	4.75	5.64	6.20	6.62	6.96	7.24	7.47	7.68	7.86	8.03	8.18	
9	.05	3.20	3.95	4.41	4.76	5.02	5.24	5.43	5.59	5.74	5.87	5.98	
	.01	4.60	5.43	5.96	6.35	6.66	6.91	7.13	7.33	7.49	7.65	7.78	
10	.05	3.15	3.88	4.33	4.65	4.91	5.12	5.30	5.46	5.60	5.72	5.83	
	.01	4.48	5.27	5.77	6.14	6.43	6.67	6.87	7.05	7.21	7.36	7.49	
11	.05	3.11	3.82	4.26	4.57	4.82	5.03	5.20	5.35	5.49	5.61	5.71	
	.01	4.39	5.15	5.62	5.97	6.25	6.48	6.67	6.84	6.99	7.13	7.25	
12	.05	3.08	3.77	4.20	4.51	4.75	4.95	5.12	5.27	5.39	5.51	5.61	
	.01	4.32	5.05	5.50	5.84	6.10	6.32	6.51	6.67	6.81	6.94	7.06	
13	.05	3.06	3.73	4.15	4.45	4.69	4.88	5.05	5.19	5.32	5.43	5.53	
	.01	4.26	4.96	5.40	5.73	5.98	6.19	6.37	6.53	6.67	6.79	6.90	
14	.05	3.03	3.70	4.11	4.41	4.64	4.83	4.99	5.13	5.25	5.36	5.46	
	.01	4.21	4.89	5.32	5.63	5.88	6.08	6.26	6.41	6.54	6.66	6.77	
15	.05	3.01	3.67	4.08	4.37	4.59	4.78	4.94	5.08	5.20	5.31	5.40	
	.01	4.17	4.84	5.25	5.56	5.80	5.99	6.16	6.31	6.44	6.55	6.66	
16	.05	3.00	3.65	4.05	4.33	4.56	4.74	4.90	5.03	5.15	5.26	5.35	
	.01	4.13	4.79	5.19	5.49	5.72	5.92	6.08	6.22	6.35	6.46	6.56	
17	.05	2.98	3.63	4.02	4.30	4.52	4.70	4.86	4.99	5.11	5.21	5.31	
	.01	4.10	4.74	5.14	5.43	5.66	5.85	6.01	6.15	6.27	6.38	6.48	
18	.05	2.97	3.61	4.00	4.28	4.49	4.67	4.82	4.96	5.07	5.17	5.27	
	.01	4.07	4.70	5.09	5.38	5.60	5.79	5.94	6.08	6.20	6.31	6.41	
19	.05	2.96	3.59	3.98	4.25	4.47	4.65	4.79	4.92	5.04	5.14	5.23	
	.01	4.05	4.67	5.05	5.33	5.55	5.73	5.89	6.02	6.14	6.25	6.34	
20	.05	2.95	3.58	3.96	4.23	4.45	4.62	4.77	4.90	5.01	5.11	5.20	
	.01	4.02	4.64	5.02	5.29	5.51	5.69	5.84	5.97	6.09	6.19	6.28	
24	.05	2.92	3.53	3.90	4.17	4.37	4.54	4.68	4.81	4.92	5.01	5.10	
	.01	3.96	4.55	4.91	5.17	5.37	5.54	5.69	5.81	5.92	6.02	6.11	
30	.05	2.89	3.49	3.85	4.10	4.30	4.46	4.60	4.72	4.82	4.92	5.00	
	.01	3.89	4.45	4.80	5.05	5.24	5.40	5.54	5.65	5.76	5.85	5.93	
40	.05	2.86	3.44	3.79	4.04	4.23	4.39	4.52	4.63	4.73	4.82	4.90	
	.01	3.82	4.37	4.70	4.93	5.11	5.26	5.39	5.50	5.60	5.69	5.76	
60	.05	2.83	3.40	3.74	3.98	4.16	4.31	4.44	4.55	4.65	4.73	4.81	
	.01	3.76	4.28	4.59	4.82	4.99	5.13	5.25	5.36	5.45	5.53	5.60	
120	.05	2.80	3.36	3.68	3.92	4.10	4.24	4.36	4.47	4.56	4.64	4.71	
	.01	3.70	4.20	4.50	4.71	4.87	5.01	5.12	5.21	5.30	5.37	5.44	
	.05	2.77	3.31	3.63	3.86	4.03	4.17	4.29	4.39	4.47	4.55	4.62	
	.01	3.64	4.12	4.40	4.60	4.76	4.88	4.99	5.08	5.16	5.23	5.29	

Copyright 2010 Cengage Learning. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part. Due to electronic rights, some third party content may be suppressed from the eBook and/or eChapter(s). Editorial review has deemed that any suppressed content does not materially affect the overall learning experience. Cengage Learning reserves the right to remove additional content at any time if subsequent rights restrictions require.