$\vec{F}(x,y) = (3y-4x)\hat{i} + (4x-y)\hat{j}$ 

when an object moves once counterclockwise around the ellipse  $4x^2 + y^2 = 4$ .

4. Use Stoke's theorem to evaluate the surface integral

$$\iint_S\!\left(\text{curl }\vec{F}.N\right)dS$$

where  $F = x i + y^2 j + z e^{xy} k$  and S is that part of surface  $z = 1 - x^2 - 2y^2$  with  $z \ge 0$ .

- 5. Use divergence theorem to evaluate the integral  $\iint_S \vec{F}.N \,dS \quad \text{where} \quad \vec{F}\big(x,y,z\big) = (\cos yz)\hat{i} + e^{xz}\hat{j} + 3z^2\,\hat{k} \;,$  where S is hemisphere surface  $z = \sqrt{4-x^2-y^2}$  together with the disk  $x^2 + y^2 \le 4$ , in x-yplane.
- 6. Evaluate the line integral  $\int_C \vec{F} . d\vec{R}$

Where  $\vec{F}(x, y) = [(2x - x^2y) e^{-xy} + tan^{-1}y]\hat{i} +$ 

$$\left[\frac{x}{y^2+1} - x^3 e^{-xy}\right] \hat{j} \text{ and C is the ellipse } 9x^2 + 4y^2 = 36.$$

[This question paper contains 4 printed pages.]

12 8 DEC 2022

Your Roll No.

Sr. No. of Question Paper: 1427

Unique Paper Code

: 32351303

Name of the Paper

: BMATH 307 - Multivariate

Calculus

Name of the Course

: B.Sc. (H) Mathematics

Semester

: III

Duration: 3 Hours

Maximum Marks: 75

## Instructions for Candidates

- 1. Write your Roll No. on the top immediately on receipt of this question paper.
- 2. All sections are compulsory
- 3. Attempt any Five questions from each section. All questions carry equal marks

## SECTION I

1. Let  $f(x,y) = \frac{xy(x^2 - y^2)x}{x^2 + y^2}$  if  $(x, y) \neq (0,0)$ 

= 0 otherwise

Show that f(0, y) = -y and f(x, 0) = x for all x and y.

2. Use incremental approximation to estimate the function  $f(x, y) = \sin(xy)$  at the point

$$\sqrt{\frac{\pi}{2}} + .01, \sqrt{\frac{\pi}{2}} - .01$$

- 3. If  $z = xy + f(x^2 + y^2)$ , show that  $y \frac{\partial z}{\partial x} x \frac{\partial z}{\partial y} = y^2 x^2$ .
- 4. Assume that maximum directional derivative of f at  $P_0(1,2)$  is equal to 50 and is attained in the direction towards Q(3,-4). Find  $\nabla f$  at  $P_0(1,2)$ .
- 5. Find the absolute extrema of  $f(x, y) = 2x^2 y^2$  on the disk  $x^2 + y^2 \le 1$ .
- 6. Use Lagrange multiplier to find the distance from (0,0,0) to plane Ax + By + Cz = D where at least one of A, B, C is nonzero.

## SECTION II

- 1. Compute the integral  $\int_0^1 \int_x^{2x} e^{y-x} dy dx$  with the order of integration reversed.
- 2. Use Polar double integral to show that a sphere of radius  $\alpha$  has volume  $\frac{4}{3}\pi$  a<sup>3</sup>.

3. Compute the area of region D bounded above by line y = x, and below by circle  $x^2 + y^2 - 2y = 0$ .

3

- 4. Find the volume of the solid bounded above by paraboloid  $z = 6 x^2 y^2$  and below by  $z = 2x^2 + y^2$ .
- 5. Evaluate  $\iiint_{D} \frac{dx \ dy \ dz}{\sqrt{x^2 + y^2 + z^2}}, \text{ where D is the solid}$  sphere  $x^2 + y^2 + z^2 \le 3$ .
- 6. Use a suitable change of variables to find the area of region R bounded by the hyperbolas xy=1 and xy=4 and the lines y=x and y=4x.

## SECTION III

- Find the mass of a wire in the shape of curve
  C: x = 3 sin t, y = 3 cos t, z = 2t for 0 ≤ t ≤ π and density at point (x, y, z) on the curve is δ(x, y, z) = x.
- 2. Find the work done by force

$$\vec{F}(x,y,z) = (y^2 - z^2)\hat{i} + (2yz)\hat{j} - (x^2)\hat{k}$$

on an object moving along the curve C given by x(t) = t,  $y(t) = t^2$ ,  $z(t) = t^3$ ,  $0 \le t \le 1$ .

3. Use Green's theorem to find the work done by the force field