

Duration:[2½Hours]

[Total Marks: 75]

- N.B. 1) All questions are compulsory.
2) Figures to the right indicate full marks.

1. (a) Attempt any **ONE** question: (8)

- i. Show that every planar graph is 5-vertex colorable.
- ii. State and prove Euler's formula for planar graphs. Hence or otherwise prove that the minimum degree of a simple planar graph is ≤ 5 .

(b) Attempt any **TWO** questions: (12)

- i. Define vertex chromatic number $\chi(G)$ of a simple finite graph G . Prove that $\chi(G) \leq \Delta(G) + 1$ where $\chi(G)$ represents vertex chromatic number of a graph G and $\Delta(G)$ denotes the maximum degree of G .
- ii. If G is a planar (p, q) graph in which every face is bounded by a cycle of length at least n , then show that $q \leq \frac{n(p-2)}{n-2}$.
- iii. For a simple graph G of order p and size q , prove that $\pi_k(G)$, the chromatic polynomial of the graph G , is monic polynomial of degree p in k with integer coefficients and constant term zero.
- iv. Show that for a cycle C_n of length n , the chromatic polynomial π_k is given by $\pi_k(C_n) = (k-1)^n + (-1)^n(k-1)$ for $n \geq 3$.

2. (a) Attempt any **ONE** question: (8)

- i. State and prove Max Flow-Min Cut Theorem.
- ii. State and prove the necessary and sufficient condition for a family of n sets to have System of Distinct Representative.

(b) Attempt any **TWO** questions: (12)

- i. Define $val(f)$, value of flow f . If f is flow in a network N and P is any f -augmenting path, then show that there exists a revised flow f' such that $val f' > val f$.
- ii. Show that the number of different system of distinct representatives for the family $A_i = \{1, 2, \dots, n\} - \{i\}$, $1 \leq i \leq n$ is D_n , the number of derangements on n symbols.
- iii. Prove that for any flow f and any cut (S, \bar{S}) , $val(f) = f^+(S) - f^-(S)$.
- iv. If $\{A_1, A_2, \dots, A_n\}$ be a family of set, then prove that the largest number of sets of the family which together have a system of distinct representative equals the minimum value of expression $|A_{i_1} \cup A_{i_2} \cup \dots \cup A_{i_k}| + (n - k)$ for all choices of $k = 1, 2, \dots, n$ and all choices of i_1, i_2, \dots, i_k with $1 \leq i_1 < i_2 < \dots < i_k \leq n$.

[TURN OVER

3. (a) Attempt any **ONE** question: (8)
- i. Derive the recurrence relation for number of ways of dividing a $(n + 1)$ - sided convex polygon into triangular regions by inserting diagonals that do not intersect in the interior and prove using generating function that the solution to this recurrence relation is a Catalan Number.
 - ii. Derive the recurrence relation to climb a staircase with n steps by taking 1 or 2 steps at a time and solve it using generating function.

- (b) Attempt any **TWO** questions: (12)
- i. Let B denotes a forbidden chess board in which a special square * has been identified. Let D denote the board obtained from the original board by deleting the row and column containing the special square and E denote the board obtained from the original board where only the special square * is removed from the board, then prove that $R(x, B) = xR(x, D) + R(x, E)$.
 - ii. Derive the recurrence relation for the number of regions into which the plane is divided by n straight lines, no two of which are parallel and no three of which are concurrent. Furthermore using generating function, show that the solution of the above recurrence is $\frac{n(n+1)}{2} + 1$
 - iii. Solve recurrence relation $a_n = a_{n-1} + 1$ for all $n \geq 2$ subject to initial conditions $a_1 = 1$ using generating function.
 - iv. Show that the number of nonnegative integer solutions of the equation $x_1 + x_2 + \dots + x_k = r$ is given by $\binom{r+k-1}{r}$.

4. Attempt any **THREE** questions: (15)
- (a) Show that a connected graph G on n vertices is a tree if and only if the chromatic polynomial of G is $k(k-1)^{n-1}$.
 - (b) Show that $\chi'(G) \geq \Delta(G)$ where $\chi'(G)$ denotes edge chromatic number and $\Delta(G)$ denotes the maximum degree of G . Give an example of the graph for which $\chi'(G) = \Delta(G)$.
 - (c) If f is any flow and K be any cut in a network N with $val(f) = cap(K)$ then show that f is maximum flow and K is minimum cut.
 - (d) Define System of distinct representatives for a family of sets. Let $A = (A_1, A_2, A_3, A_4, A_5, A_6)$, where $A_1 = \{1, 2, 3\}$, $A_2 = \{1, 2, 3, 4, 5\}$, $A_3 = \{1, 2\}$, $A_4 = \{2, 3\}$, $A_5 = \{1\}$, $A_6 = \{1, 3, 5\}$. Does family A have an System of Distinct Representative? If not, what is the largest number of sets in the family with an System of Distinct Representative?
 - (e) Prove that if B is a board of darkened squares that decomposes into the two disjoint sub boards B_1 and B_2 then prove that $R(x, B) = R(x, B_1)R(x, B_2)$, where $R(x, B)$ is a rook polynomial for board B .
 - (f) If h_n denote the number of nonnegative integral solutions of the equation $3e_1 + 4e_2 + 2e_3 + 5e_4 = n$, find generating function for h_n .
